

EFFECTIVE DATE: 09 | 01 | 2021

POLICY LAST UPDATED: 09 | 05 | 2021

OVERVIEW

Gene expression profile analysis and protein biomarkers have been proposed as a means to risk-stratify patients with prostate cancer to guide treatment decisions. These tests are intended to be used either on prostate needle biopsy tissue to guide management decisions for active surveillance or therapeutic intervention, to guide radiotherapy use after radical prostatectomy (RP), or to guide medication selection after progression in metastatic castration-resistant prostate cancer.

The following tests are addressed in this policy:

- Prolaris (Myriad)
- ProMark (Metamark Genetics)
- Decipher (Veracyte, Inc.)
- Oncotype DX Prostate (Genomic Health)
- Oncotype DX AR-V7 Nuclear Detect (Genomic Health)

MEDICAL CRITERIA

Medicare Advantage Plans

Prolaris

Prolaris is covered for the following indications. See detailed criteria below.

- A. *Early stage, needle biopsy proven prostate cancer, OR***
B. *Favorable intermediate risk, needle biopsy proven prostate cancer*

- A. The Prolaris™ assay is covered help determine which patients with early stage, needle biopsy proven prostate cancer, can be conservatively managed rather than treated with definitive surgery or radiation therapy, when the following clinical conditions are met:**

- Needle biopsy with localized adenocarcinoma of prostate (no clinical evidence of metastasis or lymph node involvement), **and**
- FFPE prostate biopsy specimen with at least 0.5 mm of cancer length, **and**
- Patient Stage as defined by the one of the following:
 - Very Low Risk Disease (T1c AND Gleason Score ≤ 6 AND PSA ≤ 10 ng/mL AND <3 prostate cores with tumor AND $\leq 50\%$ cancer in any core AND PSA density of < 0.15 ng/mL/g) **OR**
 - Low Risk Disease (T1-T2a AND Gleason Score ≤ 6 AND PSA ≤ 10 ng/mL), **and**
- Patient has an estimated life expectancy of greater than or equal to 10 years, **and**
- Patient is a candidate for and is considering conservative therapy and yet and would be eligible for definitive therapy (radical prostatectomy, radiation therapy or brachytherapy), **and**
- Result will be used to determine treatment between definitive therapy and conservative management, **and**
- Patient has not received pelvic radiation or androgen deprivation therapy prior to the biopsy, **and**
- Test is ordered by a physician certified in the Myriad PROLARIS™ Certification and Training Registry (CTR), **and**
- Patient is monitored for disease progression according to established standard of care, **and**
- Physician must report the development of metastasis or prostate cancer deaths in patients not treated definitively who were deemed low risk by the assay.

B. The Prolaris™ assay is covered help determine which patients with favorable intermediate risk, needle biopsy proven prostate cancer (as defined below), can be conservatively managed rather than treated with definitive surgery or radiation therapy, when the following clinical conditions are met:

- Needle biopsy with localized adenocarcinoma of prostate (no clinical evidence of metastasis or lymph node involvement), **and**
- FFPE prostate biopsy specimen with at least 0.5 mm of cancer length, **and**
- Patients with favorable intermediate-risk disease, defined by the NCCN as follows:
 - Predominant Gleason grade 3 (i.e. Gleason score 3+4=7), percentage of positive cores <50%, and no more than 1 NCCN intermediate-risk factor**DEFINITION:** NCCN intermediate risk factors include T2b-T2c, Gleason score 7, and PSA 10-20 ng/mL
- Patient has an estimated life expectancy of greater than or equal to 10 years, **and**
- Patient is a candidate for and is considering conservative therapy and yet and would be eligible for definitive
- therapy (radical prostatectomy, radiation therapy or brachytherapy), **and**
- Result will be used to determine treatment between definitive therapy and conservative management, **and**
- Patient has not received pelvic radiation or androgen deprivation therapy prior to the biopsy, **and**
- Patient is monitored for disease progression according to established standard of care.

Decipher

Decipher is covered for men with prostate cancer with:

- Localized or biochemically recurrent adenocarcinoma of the prostate (i.e., no clinical evidence of metastasis), **and**
- Who have a life expectancy of greater than or equal to 10 years, **and**
- The assay is performed on formalin-fixed paraffin embedded (FFPE) prostate biopsy tissue with at least 0.5 mm of linear tumor diameter or FFPE tissue from a prostate resection specimen, **and**
- Result will be used to determine treatment according to established practice guidelines, **and**
- Patient has not received pelvic radiation or androgen deprivation therapy (ADT) prior to the biopsy or prostate resection specimen, **and**
- Patient is monitored for disease progression according to established standards of care, **and**
- If they are a candidate and are considering (or being considered for) at least 1 of the following:
 - Conservative management and yet would be eligible for definitive therapy (radical prostatectomy (RP), radiation or brachytherapy), **or**
 - Radiation therapy and yet would be eligible for the addition of a brachytherapy boost, **or**
 - Radiation therapy and yet would be eligible for the addition of short-term (ADT), **or**
 - Radiation therapy with short-term ADT yet would be eligible for the use of long-term ADT, **or**
 - Radiation with standard ADT yet would be eligible for systemic therapy intensification using next generation androgen signaling inhibitors or chemotherapy, **or**
 - Observation post-prostatectomy yet would be eligible for the addition of post-operative adjuvant radiotherapy, **or**
 - Salvage radiotherapy post-prostatectomy yet would be eligible for the addition of ADT.

OncotypeDX Prostate

Oncotype DX Prostate is covered for the following indications. See detailed criteria below.

- A. Early stage, needle biopsy proven prostate cancer, OR**
- B. Favorable intermediate risk, needle biopsy proven prostate cancer**

A. When used to determine which patients with early stage, needle biopsy proven prostate cancer, can be conservatively managed rather than treated with definitive surgery or radiation therapy and the following clinical conditions are met:

- Needle biopsy with localized adenocarcinoma of prostate (no clinical evidence of metastasis or lymph node involvement), **and**
- Patient stage as defined by the one of the following:
 - Very Low Risk Disease (T1c AND Gleason Score = 6 AND PSA = 10 ng/mL
 - Low Risk Disease (T1-T2a AND Gleason Score = 6 AND PSA = 10 ng/mL), **and**
- Patient has an estimated life expectancy of ≥ 10 years, **and**
- Patient is a candidate for and is considering conservative therapy and yet would be eligible for definitive therapy (radical prostatectomy, radiation therapy or brachytherapy), **and**
- Patient has not received pelvic radiation or androgen deprivation therapy prior to the biopsy, **and**
- Test is ordered by a physician certified in the Genomic Health™ Oncotype DX® Prostate Cancer Assay Certification and Training Registry (CTR), **and**
- Patient is monitored for disease progression according to active surveillance guidelines as recorded in NCCN guidelines.

B. When used to determine which patients with favorable intermediate-risk, needle biopsy proven prostate cancer, can be conservatively managed rather than treated with definite surgery or radiation therapy and the following clinical conditions are met:

- Needle biopsy with localized adenocarcinoma of prostate (no clinical evidence of metastasis or lymph node involvement), **and**
- FFPE prostate biopsy specimen with at least 0.5 mm of cancer length, and NCCN Favorable Intermediate-risk disease defined as:
 - Gleason Grade Group 2 (Gleason Sum 3+4=7), **and**
- Patient has an estimated life expectancy of greater than or equal to 10 years, **and**
- Patient is a candidate for and is considering conservative therapy and yet would be eligible for definitive therapy (radical prostatectomy, radiation therapy or brachytherapy), **and**
- Result will be used to determine treatment between definitive therapy and conservative management, **and**
- Patient has not received pelvic radiation or androgen deprivation therapy prior to the biopsy, **and**
- Patient is monitored for disease progression according to established standard of care.

OncotypeDX AR-V7 Nuclear Detect

OncotypeDX AR-V7 Nuclear Detect assay is covered when the following clinical conditions are met:

- Patient has progressive mCRPC as defined by the Prostate Cancer Working Group 2 guidelines (a minimum of 2 rising prostate-specific antigen (PSA) levels 1 or more weeks apart, new lesions by bone scintigraphy, and/or new or enlarging soft tissue lesions by computed tomography (CT) or magnetic resonance imaging (MRI)).
- Patient has failed one androgen receptor signaling inhibitor (ARSi), (presently Enzalutamide (Xtandi), Apalutamide (Erleada), or Abiraterone (Zytiga) and future similar class drugs approved by the FDA).
- Patient is considered appropriate for treatment by their treating physician for the alternative ARSi as a single agent.
- Circulating tumor cells (CTC) with nuclear expression of AR-V7 protein have been assessed prior to initiation of therapy.

Commercial Products

Unless otherwise noted, for any test filed with an Unlisted CPT code, the medical necessity criteria in the Genetic Testing Services policy would be used. Please see the Related Policies section.

PRIOR AUTHORIZATION

Medicare Advantage Plans

Prior authorization is required for Prolaris, Decipher, Oncotype DX Prostate, and OncotypeDX AR-V7 Nuclear Detect and is obtained via the online tool for participating providers.

Medicare Advantage Plans and Commercial Products

There is no specific CPT coding for some testing referenced in this policy. Therefore, an Unlisted CPT code should be used (See Coding Section for details). All Unlisted genetic testing CPT codes require prior authorization to determine what service is being rendered and if the service is covered or not medically necessary. See the Related Policies section.

Prior authorization is required for Medicare Advantage Plans and recommended for Commercial products and is obtained via the online tool for participating providers. See the Related Policies section.

Note: Laboratories are not allowed to obtain clinical authorization or participate in the authorization process on behalf of the ordering physician. Only the ordering physician shall be involved in the authorization, appeal or other administrative processes related to prior authorization/medical necessity.

In no circumstance shall a laboratory or a physician/provider use a representative of a laboratory or anyone with a relationship to a laboratory and/or a third party to obtain authorization on behalf of the ordering physician, to facilitate any portion of the authorization process or any subsequent appeal of a claim where the authorization process was not followed and/or a denial for clinical appropriateness was issued, including any element of the preparation of necessary documentation of clinical appropriateness. If a laboratory or a third party is found to be supporting any portion of the authorization process, BCBSRI will deem the action a violation of this policy and severe action will be taken up to and including termination from the BCBSRI provider network. If a laboratory provides a laboratory service that has not been authorized, the service will be denied as the financial liability of the participating laboratory and may not be billed to the member.

POLICY STATEMENT

Medicare Advantage Plans

The following tests may be considered medically necessary when the medical criteria above are met:

- Prolaris
- Decipher
- Oncotype DX Prostate
- Oncotype DX AR-V7 Nuclear Detect

The following test is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes:

- ProMark

Commercial Products

The following tests are not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes:

- Prolaris
- Decipher
- Oncotype DX Prostate
- Oncotype DX AR-V7 Nuclear Detect
- ProMark

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable genetic testing and not medically necessary/not covered benefits/coverage.

BACKGROUND

Prostate cancer is the second most common noncutaneous cancer diagnosed among men in the United States. Autopsy studies in the era before the availability of prostate-specific antigen (PSA) screening have identified incidental cancerous foci in 30% of men 50 years of age, with incidence reaching 75% at age 80 years.

Localized prostate cancers may appear very similar clinically at diagnosis. However, they often exhibit diverse risk of progression that may not be captured by clinical risk categories (eg, D'Amico criteria) or prognostic tools based on clinical findings, including PSA titers, Gleason grade, or tumor stage. In studies of conservative management, the risk of localized disease progression based on prostate cancer-specific survival rates at 10 years may range from 15% to 20% to perhaps 27% at 20-year follow-up. Among older men (ages \geq 70 years) with low-risk disease, comorbidities typically supervene as a cause of death; these men will die with prostate cancer present, rather than from cancer itself. Other very similar appearing low-risk tumors may progress unexpectedly rapidly, quickly disseminating and becoming incurable.

Risk Stratification in Newly Diagnosed Disease

In the United States, most prostate cancers are clinically localized at diagnosis due in part to the widespread use of PSA testing. Clinicopathologic characteristics are used to stratify patients by risk based on the extent of the primary tumor (T category), nearby lymph node involvement (N category), metastasis (M category), PSA level and Gleason score. The National Comprehensive Cancer Network and American Urological Association risk categories for clinically localized prostate cancer are similar, derived from the D'Amico criteria and broadly include low-, intermediate-, or high-risk as follows as well as subcategories within these groups:

- Low: T1-T2a and Gleason score \leq 6/Gleason grade group 1 and PSA level \leq 10 ng/mL;
- Intermediate: T2b-T2c or Gleason score 3+4=7/Gleason grade group 2 or Gleason score 4+3=7/Gleason grade group 3 or PSA level 10-20 ng/mL;
- High: T3a or Gleason score 8/Gleason grade group 4 or Gleason score 9-10/Gleason grade group 5 or PSA level $>$ 20 ng/mL.

Risk stratification is combined with patient age, life expectancy, and treatment preferences to make initial therapy decisions.

Monitoring After Prostatectomy

All normal prostate tissue and tumor tissue is theoretically removed during radical prostatectomy (RP), so the serum level of PSA should be undetectable following RP. Detectable PSA post-RP indicates residual prostate tissue and presumably persistent or recurrent disease. PSA is serially measured following RP to detect early disease recurrence. The National Comprehensive Cancer Network recommends monitoring serum PSA every 6 to 12 months for the first 5 years and annually thereafter. Many recurrences following RP can be successfully treated. The American Urological Association has recommended that biochemical recurrence be defined as a serum PSA of 0.2 ng/mL or higher, which is confirmed by the second determination with a PSA level of 0.2 ng/mL or higher.

Castration-Resistant Prostate Cancer

Androgen deprivation therapy (ADT) is generally the initial treatment for patients with advanced prostate cancer. ADT can produce tumor response and improve quality of life but most patients will eventually progress on ADT. Disease that progresses while the patient is on ADT is referred to as castration-resistant prostate cancer. After progression, continued ADT is generally used in conjunction with other treatments. Androgen pathways are important in the progression of castration-resistant prostate cancer. Several drugs have been developed that either inhibit enzymes involved in androgen production or inhibit the androgen receptor, such as abiraterone and enzalutamide. Taxane chemotherapy with docetaxel or cabazitaxel may also be used after progression. Immunotherapy (sipuleucel-T) or radium 223 are options for select men.

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Prolaris® (Myriad Genetics), Oncotype DX® Prostate and Oncotype

DX AR-V7 Nuclear Detect (Genomic Health), Decipher® gene expression profiling test (Decipher Corp), and the ProMark™ protein biomarker test (Metamark Genetics) are available under the auspices of CLIA. Laboratories that offer laboratory-developed tests must be licensed by CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of this test.

Prolaris

Prolaris is an RNA based assay measuring the expression of 31 cell cycle progression (CCP) genes and 15 “housekeeping” genes that act as internal controls and normalization standards in each patient sample. The assay is performed on formalin fixed paraffin embedded (FFPE) prostate cancer blocks. The assay results are reported as a numerical score along with accompanying interpretive information.

Medicare Advantage Plans

The potential usefulness of this test is that it allows physicians to determine which patients with early prostate cancer are candidates for active surveillance or observation and are more likely to have a good outcome without needing to receive definitive treatment.

The clinical performance of this assay was assessed in several retrospective validation studies. These include two British cohorts of men diagnosed with prostate cancer on biopsy and then treated conservatively; and an additional cohort of men diagnosed by transurethral resection of the prostate (TURP) and conservatively managed. Further validation was performed in various other cohorts including men who underwent radical prostatectomy, and men treated with definitive radiotherapy. The Prolaris cell cycle progression score (CCP) was found to be an independent and more robust prognostic factor for disease related death than traditional clinicopathologic factors although disease stage and Gleason score consistently portended a more negative prognostic picture.

Commercial Products

For individuals who have low- or intermediate-risk clinically localized untreated prostate cancer who receive Prolaris, the evidence includes retrospective cohort studies of clinical validity using archived samples in patients of mixed risk categories. Relevant outcomes include overall survival, disease-specific survival, quality of life, and treatment-related morbidity. For the low-risk group, the Prostate Testing for Cancer and Treatment (ProtecT) trial showed 99% ten-year disease-specific survival in mostly low-risk patients receiving active surveillance. The low mortality rate estimated with tight precision makes it unlikely that a test intended to identify a subgroup of low-risk men with a net benefit from immediate treatment instead of active surveillance would find such a group. For the intermediate-risk group, the evidence of improved clinical validity or prognostic accuracy for prostate cancer death using Prolaris Cell Cycle Progression score in patients managed conservatively after needle biopsy has shown some improvement in areas under the receiver operating characteristic curve over clinicopathologic risk stratification tools. There is limited indirect evidence for potential clinical utility. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have localized prostate cancer treated with RP who receive Prolaris, the evidence includes retrospective cohort studies of clinical validity using archived samples. Relevant outcomes include overall survival, disease-specific survival, quality of life, and treatment-related morbidity. Evidence of improved clinical validity or prognostic accuracy for prostate cancer death using the Prolaris Cell Cycle Progression score in patients after prostatectomy has shown some improvement in areas under the receiver operating characteristic curve over clinicopathologic risk stratification tools. The evidence is insufficient to determine the effects of the technology on health outcomes.

Decipher Prostate Cancer Classifier Assay

The Decipher prostate cancer assay, a 22-biomarker expression signature using oligonucleotide microarray technology, interrogates 1.4 million RNAs extracted from a formalin-fixed paraffin embedded (FFPE) tissue block of the index lesion (defined by highest tumor stage or histological Gleason grade) from the RP specimen, or of the diagnostic biopsy core (defined by the highest Gleason grade). The biomarkers that

comprise the Decipher classifier include cell cycle progression, androgen signaling, cell adhesion, tumor cell motility, migration and immune evasion functions.

Medicare Advantage Plans

In the treatment of prostate cancer, a number of treatment approaches have shown potential benefits for survival, but also significant adverse events. Therefore, the risk of a patient having an unfavorable outcome due to prostate cancer within the patient's expected lifetime is at the core of the management of prostate cancer. Patients at a higher risk are recommended to have a greater intensity of treatment. For the treatment of localized prostate cancer there is a well-established risk stratification scheme based on clinical and pathological factors, though even within a single risk stratum there is significant variability, which Decipher® has shown the ability to further stratify. There are also numerous potential treatment options of varying levels of intensity with varying severities of side effects, so the potential benefit of treatment, which is heavily driven by risk stratum, must be weighed against the downsides of therapy. For patients with recurrent or metastatic disease there are also a number of effective treatment options of varying intensity, also associated with varying severities of adverse events. While there are no randomized controlled trials of outcomes using the Decipher® test, numerous studies from different institutions have all had similar and consistent findings, providing evidence that this test accurately risk stratifies patients based on genetic information and accurately predicts risk of biochemical recurrence, metastatic disease, or prostate-cancer specific mortality. Given that existing treatment paradigms are heavily reliant upon risk assessment, the ability to accurately risk stratify has potential utility in the management of prostate cancer. As such, this test provides clinically actionable incremental information that fits into existing evidence-based or consensus-recommended prostate cancer treatment paradigms.

Commercial Products

For individuals who have localized prostate cancer who are treated with RP and who receive the Decipher prostate cancer classifier, the evidence includes a study of analytic validity, prospective and retrospective studies of clinical validity using overlapping archived samples, decision-curve analyses examining indirect evidence of clinical utility, and prospective decision-impact studies without pathology or clinical outcomes. Relevant outcomes include overall survival, disease-specific survival, quality of life, and treatment-related morbidity. The clinical validity of the Decipher genomic classifier has been evaluated in samples of patients with high-risk prostate cancer undergoing different interventions following RP. Studies reported some incremental improvement in discrimination. However, it is unclear whether there is consistently improved reclassification—particularly to higher risk categories—or whether the test could be used to predict which men will benefit from radiotherapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

Oncotype DX Prostate

Oncotype DX® Prostate Cancer Assay is prostate biopsy-based 17-gene RT-PCR assay, representing four molecular pathways (androgen signaling, cellular organization, stromal response and proliferation), that provides a biologic measure of cancer aggressiveness. The assay is indicated for men who are considered candidates for active surveillance (AS) (those with NCCN® very low, low and favorable intermediate risk prostate cancer). The assay is designed to inform decisions between AS and immediate treatment.

Medicare Advantage Plans

The potential usefulness of this test is that it allows physicians to determine which patients with early prostate cancer are candidates for active surveillance and are more likely to have a good outcome without needing to receive definitive treatment.

Studies demonstrate that in men with NCCN very low-, low-, or intermediate-risk prostate cancer who were potential candidates for AS, Oncotype DX was prospectively validated as a biopsy-based predictor of adverse pathology, biochemical recurrence and metastasis, thus establishing the assay as a robust and independent measure of prostate cancer aggressiveness.

Commercial Products

For individuals who have low- or intermediate-risk clinically localized untreated prostate cancer who receive Oncotype DX Prostate, the evidence includes case-cohort and retrospective cohort studies of clinical validity using archived samples in patients of mixed risk categories, and a decision-curve analysis examining indirect evidence of clinical utility. Relevant outcomes include overall survival, disease-specific survival, quality of life, and treatment-related morbidity. Evidence for clinical validity and potential clinical utility of Oncotype DX Prostate in patients with clinically localized prostate cancer derives from a study predicting adverse pathology after RP. The validity of using tumor pathology as a surrogate for risk of progression and cancer-specific death is unclear. It is also unclear whether results from an RP population can be generalized to an active surveillance population. The evidence is insufficient to determine the effects of the technology on health outcomes.

Oncotype DX AR-V7 Nuclear Detect

Oncotype DX AR-V7 Nuclear Detect test is a commercially available, circulating tumor cell (CTC) based, liquid biopsy test. The test detects patients with CTCs who have nuclear expression of the AR-V7 truncated protein and provides information that can help guide treatment selection in patients with metastatic castrate resistant prostate cancer (mCRPC).

Medicare Advantage Plans

Men with metastatic castration resistant prostate cancer (mCRPC) have multiple life extending, FDA-approved therapeutics options. However, there is no clear consensus on the therapeutic sequencing after initial exposure to an androgen receptor signaling inhibitor (ARSi), such as apalutamide, abiraterone or enzalutamide. The response rate for a second ARSi, Abiraterone after Enzalutamide, or Enzalutamide after Abiraterone is lower than the initial exposure. Therefore, the most common clinical decision focuses on whether to start a second ARSi or taxane chemotherapy. It is therefore important to identify patients who will not respond to a 2nd ARSi in order to 1) avoid giving an ineffective therapy, and 2) delaying giving a more effective therapy, such as taxane chemotherapy, taxane combination therapies, Radium -223, PARP inhibitors, and platinum chemotherapy.

AR-V7 protein results from alternative androgen receptor (AR) mRNA splicing, which produces a constitutively active receptor that is associated with resistance to ARSi such as abiraterone and enzalutamide. A growing body of evidence suggests that patients with ARV7 positive mCRPC do not benefit from ARSi therapy but may respond to taxanes, such as docetaxel.³⁻⁵ Supporting observations include that 1) AR-V7 positivity is associated with resistance to androgen receptor-targeted therapies⁴; 2) taxanes are equally effective in AR-V7 –positive and AR-V7–negative mCRPC patients^{3,5} ; and 3) AR-V7 status may change during therapy. For these reasons, the National Comprehensive Cancer Network (NCCN) suggests that AR-V7 is a biomarker that may help guide therapy in mCRPC.

AR signaling requires that the AR transcriptional elements bind to DNA within the nucleus of the cancer cell. Therefore, the nuclear localization of the AR-V7 truncated protein may improve the clinical specificity of predicting ARSi resistance.⁸ Analysis of AR-V7 localization scoring guides has demonstrated that only nuclear AR-V7 protein expression improves the clinical specificity of predicting ARSi resistance, and importantly, is associated with improved overall survival with taxane chemotherapy.

Commercial Products

For individuals who have metastatic castration-resistant prostate cancer who receive the Oncotype DX AR-V7 Nuclear Detect, the evidence includes a retrospective cohort study of clinical validity using archived samples, and no studies of clinical utility. Relevant outcomes include overall survival, diseasespecific survival, quality of life, and treatment-related morbidity. Current evidence does not support improved outcomes with Oncotype DX AR-V7 Nuclear Detect, given that only a single clinical validity study, meeting inclusion criteria was available. The evidence is insufficient to determine the effects of the technology on health outcomes.

ProMark Protein Biomarker Test

The ProMark assay includes 8 biomarkers that predict prostate pathology aggressiveness and lethal outcomes: DERL1, PDSS2, pS6, YBX1, HSPA9, FUS, SMAD4, and CUL2. The assay results are combined using

predefined coefficients for each marker from a logistic regression model to calculate a risk score. The risk score is a continuous number between 0 and 1, which estimates the probability of “non-GS 6” pathology.

Medicare Advantage Plans and Commercial Products

For individuals who have low- or intermediate-risk clinically localized untreated prostate cancer who receive the ProMark protein biomarker test, the evidence includes a retrospective cohort study of clinical validity using archived samples and no studies of clinical utility. Relevant outcomes include overall survival, disease-specific survival, quality of life, and treatment-related morbidity. Current evidence does not support improved outcomes with ProMark given that only a single clinical validity study is available. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

The following CPT codes may be medically necessary for Medicare Advantage Plans when the medical criteria above is met and is not medically necessary for Commercial Products.

This code can be used for the Prolaris Assay.

81541 Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score

This code can be used for Decipher Prostate.

81542 Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score

This code can be used for the Oncotype DX® Prostate.

0047U Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score

The following Unlisted CPT code requires prior authorization for Medicare Advantage Plans and Commercial Products. The code can be used for any test identified in this policy that does not have a specific CPT code.

81479 Unlisted molecular pathology procedure

RELATED POLICIES

Genetic Testing Services

Proprietary Laboratory Analyses (PLA)

PUBLISHED

Provider Update, July 2021

Provider Update, May 2020

Provider Update, August 2019

Provider Update, November 2017

Provider Update, September 2016

REFERENCES

1. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: Prostate Cancer Genomic Classifier Assay for Men with Localized Disease (L38339)
2. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: MolDX: Prostate Cancer Genomic Classifier Assay for Men with Localized Disease (A57372)
3. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: Genomic Health™ ONCOTYPE DX® Prostate Cancer Assay (L36364)
4. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: ONCOTYPE DX® Genomic Prostate Score for Men with Favorable Intermediate Risk Prostate Cancer (L37305)Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding MolDX: Genomic Health™ ONCOTYPE DX® Prostate Cancer Assay (A56371)

5. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: ONCOTYPE DX AR-V7 Nucleus Detect for Men with Metastatic Castrate Resistant Prostate Cancer (MCRPC) (L37746)
6. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: MolDX: ONCOTYPE DX AR-V7 Nucleus Detect for Men with Metastatic Castrate Resistant Prostate Cancer (MCRPC) (A57601)
7. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: PROLARIS™ Prostate Cancer Genomic Assay (L36350)
8. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: MolDX: PROLARIS™ Prostate Cancer Genomic Assay (A57511)
9. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: PROLARIS™ Prostate Cancer Genomic Assay for Men with Favorable Intermediate Risk Disease (L37082)
10. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: MolDX: PROLARIS™ Prostate Cancer Genomic Assay for Men with Favorable Intermediate Risk Disease (A57691)
11. Dall'Era MA, Cooperberg MR, Chan JM, et al. Active surveillance for early-stage prostate cancer: review of the current literature. *Cancer*. Apr 15 2008; 112(8): 1650-9. PMID 18306379
12. Bangma CH, Roemeling S, Schroder FH. Overdiagnosis and overtreatment of early detected prostate cancer. *World J Urol*. Mar 2007; 25(1): 3-9. PMID 17364211
13. Johansson JE, Andren O, Andersson SO, et al. Natural history of early, localized prostate cancer. *JAMA*. Jun 09 2004; 291(22): 2713-9. PMID 15187052
14. Ploussard G, Epstein JI, Montironi R, et al. The contemporary concept of significant versus insignificant prostate cancer. *Eur Urol*. Aug 2011; 60(2): 291-303. PMID 21601982
15. Harnden P, Naylor B, Shelley MD, et al. The clinical management of patients with a small volume of prostatic cancer on biopsy: what are the risks of progression? A systematic review and meta-analysis. *Cancer*. Mar 01 2008; 112(5): 971-81. PMID 18186496
16. Brimo F, Montironi R, Egevad L, et al. Contemporary grading for prostate cancer: implications for patient care. *Eur Urol*. May 2013; 63(5): 892-901. PMID 23092544
17. Eylert MF, Persad R. Management of prostate cancer. *Br J Hosp Med (Lond)*. Feb 2012; 73(2): 95-9. PMID 22504752
18. Eastham JA, Kattan MW, Fearn P, et al. Local progression among men with conservatively treated localized prostate cancer: results from the Transatlantic Prostate Group. *Eur Urol*. Feb 2008; 53(2): 347-54. PMID 17544572
19. Bill-Axelsson A, Holmberg L, Ruutu M, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *N Engl J Med*. May 12 2005; 352(19): 1977-84. PMID 15888698
20. Thompson IM, Goodman PJ, Tangen CM, et al. Long-term survival of participants in the prostate cancer prevention trial. *N Engl J Med*. Aug 15 2013; 369(7): 603-10. PMID 23944298
21. Albertsen PC, Hanley JA, Fine J. 20-year outcomes following conservative management of clinically localized prostate cancer. *JAMA*. May 04 2005; 293(17): 2095-101. PMID 15870412
22. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer. Version 2.2020. https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed October 14, 2020.
23. American Urological Association (AUA). Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. 2017; [http://www.auanet.org/guidelines/clinically-localized-prostate-cancer-new-\(aua/astro/suo-guideline-2017\)](http://www.auanet.org/guidelines/clinically-localized-prostate-cancer-new-(aua/astro/suo-guideline-2017)). Accessed October 14, 2020.
24. Thompson IM, Valicenti RK, Albertsen P, et al. Adjuvant and salvage radiotherapy after prostatectomy: AUA/ASTRO Guideline. *J Urol*. Aug 2013; 190(2): 441-9. PMID 23707439
25. Food and Drug Administration (FDA). The Public Health Evidence for FDA Oversight of Laboratory Developed Tests: 20 Case Studies. 2015; [http://www.nila-usa.org/images/nila/The%20Public%20Health%20Case%20for%20FDA%20Oversight%20of%20LDTs%20110915\(2\)_508ed%20\(1\).pdf](http://www.nila-usa.org/images/nila/The%20Public%20Health%20Case%20for%20FDA%20Oversight%20of%20LDTs%20110915(2)_508ed%20(1).pdf). Accessed October 14, 2020.
26. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Gene Expression Analysis for Prostate Cancer Management. TEC Assessments. 2014; Volume 28: Tab 11.

27. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Gene Expression Profiling for Prostate Cancer Management. TEC Assessments. 2015;Volume 29:Tab 9.
28. Borley N, Feneley MR. Prostate cancer: diagnosis and staging. *Asian J Androl*. Jan 2009; 11(1): 74-80. PMID 19050692
29. Freedland SJ. Screening, risk assessment, and the approach to therapy in patients with prostate cancer. *Cancer*. Mar 15 2011; 117(6): 1123-35. PMID 20960523
30. Whitson JM, Carroll PR. Active surveillance for early-stage prostate cancer: defining the triggers for intervention. *J Clin Oncol*. Jun 10 2010; 28(17): 2807-9. PMID 20439633
31. Albertsen PC. Treatment of localized prostate cancer: when is active surveillance appropriate?. *Nat Rev Clin Oncol*. Jul 2010; 7(7): 394-400. PMID 20440282
32. Ip S, Dahabreh IJ, Chung M, et al. An evidence review of active surveillance in men with localized prostate cancer. *Evid Rep Technol Assess (Full Rep)*. Dec 2011; (204): 1-341. PMID 23126653
33. Nam RK, Cheung P, Herschorn S, et al. Incidence of complications other than urinary incontinence or erectile dysfunction after radical prostatectomy or radiotherapy for prostate cancer: a population-based cohort study. *Lancet Oncol*. Feb 2014; 15(2): 223-31. PMID 24440474
34. Hamdy FC, Donovan JL, Lane JA, et al. 10-Year Outcomes after Monitoring, Surgery, or Radiotherapy for Localized Prostate Cancer. *N Engl J Med*. Oct 13 2016; 375(15): 1415-1424. PMID 27626136
35. Tosoian JJ, Mamawala M, Epstein JI, et al. Intermediate and Longer-Term Outcomes From a Prospective Active-Surveillance Program for Favorable-Risk Prostate Cancer. *J Clin Oncol*. Oct 20 2015; 33(30): 3379-85. PMID 26324359
36. Klotz L, Vesprini D, Sethukavalan P, et al. Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. *J Clin Oncol*. Jan 20 2015; 33(3): 272-7. PMID 25512465
37. Wilt TJ, Brawer MK, Jones KM, et al. Radical prostatectomy versus observation for localized prostate cancer. *N Engl J Med*. Jul 19 2012; 367(3): 203-13. PMID 22808955
38. Wilt TJ, Jones KM, Barry MJ, et al. Follow-up of Prostatectomy versus Observation for Early Prostate Cancer. *N Engl J Med*. Jul 13 2017; 377(2): 132-142. PMID 28700844
39. van den Bergh RC, Korfage IJ, Roobol MJ, et al. Sexual function with localized prostate cancer: active surveillance vs radical therapy. *BJU Int*. Oct 2012; 110(7): 1032-9. PMID 22260273
40. Johansson E, Steineck G, Holmberg L, et al. Long-term quality-of-life outcomes after radical prostatectomy or watchful waiting: the Scandinavian Prostate Cancer Group-4 randomised trial. *Lancet Oncol*. Sep 2011; 12(9): 891-9. PMID 21821474
41. Wu CL, Schroeder BE, Ma XJ, et al. Development and validation of a 32-gene prognostic index for prostate cancer progression. *Proc Natl Acad Sci U S A*. Apr 09 2013; 110(15): 6121-6. PMID 23533275
42. Spans L, Clinckemalie L, Helsen C, et al. The genomic landscape of prostate cancer. *Int J Mol Sci*. May 24 2013; 14(6): 10822-51. PMID 23708091
43. Schoenborn JR, Nelson P, Fang M. Genomic profiling defines subtypes of prostate cancer with the potential for therapeutic stratification. *Clin Cancer Res*. Aug 01 2013; 19(15): 4058-66. PMID 23704282
44. Huang J, Wang JK, Sun Y. Molecular pathology of prostate cancer revealed by next-generation sequencing: opportunities for genome-based personalized therapy. *Curr Opin Urol*. May 2013; 23(3): 189-93. PMID 23385974
45. Yu YP, Song C, Tseng G, et al. Genome abnormalities precede prostate cancer and predict clinical relapse. *Am J Pathol*. Jun 2012; 180(6): 2240-8. PMID 22569189
46. Agell L, Hernandez S, Nonell L, et al. A 12-gene expression signature is associated with aggressive histological in prostate cancer: SEC14L1 and TCEB1 genes are potential markers of progression. *Am J Pathol*. Nov 2012; 181(5): 1585-94. PMID 23083832
47. Thompson I, Thrasher JB, Aus G, et al. Guideline for the management of clinically localized prostate cancer: 2007 update. *J Urol*. Jun 2007; 177(6): 2106-31. PMID 17509297
48. Kattan MW, Eastham JA, Wheeler TM, et al. Counseling men with prostate cancer: a nomogram for predicting the presence of small, moderately differentiated, confined tumors. *J Urol*. Nov 2003; 170(5): 1792-7. PMID 14532778
49. Cooperberg MR, Freedland SJ, Pasta DJ, et al. Multiinstitutional validation of the UCSF cancer of the prostate risk assessment for prediction of recurrence after radical prostatectomy. *Cancer*. Nov 15 2006; 107(10): 2384-91. PMID 17039503

50. Chen RC, Chang P, Vetter RJ, et al. Recommended patient-reported core set of symptoms to measure in prostate cancer treatment trials. *J Natl Cancer Inst.* Jul 2014; 106(7). PMID 25006192
51. Cuzick J, Berney DM, Fisher G, et al. Prognostic value of a cell cycle progression signature for prostate cancer death in a conservatively managed needle biopsy cohort. *Br J Cancer.* Mar 13 2012; 106(6): 1095-9. PMID 22361632
52. Cuzick J, Stone S, Fisher G, et al. Validation of an RNA cell cycle progression score for predicting death from prostate cancer in a conservatively managed needle biopsy cohort. *Br J Cancer.* Jul 28 2015; 113(3): 382-9. PMID 26103570
53. Lin DW, Crawford ED, Keane T, et al. Identification of men with low-risk biopsy-confirmed prostate cancer as candidates for active surveillance. *Urol Oncol.* Jun 2018; 36(6): 310.e7-310.e13. PMID 29655620
54. Montironi R, Mazzuccheli R, Scarpelli M, et al. Gleason grading of prostate cancer in needle biopsies or radical prostatectomy specimens: contemporary approach, current clinical significance and sources of pathology discrepancies. *BJU Int.* Jun 2005; 95(8): 1146-52. PMID 15877724
55. Sommariva S, Tarricone R, Lazzeri M, et al. Prognostic Value of the Cell Cycle Progression Score in Patients with Prostate Cancer: A Systematic Review and Meta-analysis. *Eur Urol.* Jan 2016; 69(1): 107-15. PMID 25481455
56. Crawford ED, Scholz MC, Kar AJ, et al. Cell cycle progression score and treatment decisions in prostate cancer: results from an ongoing registry. *Curr Med Res Opin.* Jun 2014; 30(6): 1025-31. PMID 24576172
57. Shore N, Concepcion R, Saltzstein D, et al. Clinical utility of a biopsy-based cell cycle gene expression assay in localized prostate cancer. *Curr Med Res Opin.* Apr 2014; 30(4): 547-53. PMID 24320750
58. Shore ND, Kella N, Moran B, et al. Impact of the Cell Cycle Progression Test on Physician and Patient Treatment Selection for Localized Prostate Cancer. *J Urol.* Mar 2016; 195(3): 612-8. PMID 26403586
59. Schaink A, Li C, Wells D, et al. Prolaris Cell Cycle Progression Test for Localized Prostate Cancer: A Health Technology Assessment. *Ont Health Technol Assess Ser.* 2017; 17(6): 1-75. PMID 28572867
60. Klein EA, Cooperberg MR, Magi-Galluzzi C, et al. A 17-gene assay to predict prostate cancer aggressiveness in the context of Gleason grade heterogeneity, tumor multifocality, and biopsy undersampling. *Eur Urol.* Sep 2014; 66(3): 550-60. PMID 24836057
61. Cullen J, Rosner IL, Brand TC, et al. A Biopsy-based 17-gene Genomic Prostate Score Predicts Recurrence After Radical Prostatectomy and Adverse Surgical Pathology in a Racially Diverse Population of Men with Clinically Low- and Intermediate-risk Prostate Cancer. *Eur Urol.* Jul 2015; 68(1): 123-31. PMID 25465337
62. Whalen MJ, Hackert V, Rothberg MB, et al. Prospective correlation between likelihood of favorable pathology on the 17-Gene Genomic Prostate Score and actual pathological outcomes at radical prostatectomy. *Urol Pract.* Sep 2016;3(5):379-386.
<https://www.sciencedirect.com/science/article/abs/pii/S2352077915002411>. Accessed October 14, 2020.
63. Van Den Eeden SK, Lu R, Zhang N, et al. A Biopsy-based 17-gene Genomic Prostate Score as a Predictor of Metastases and Prostate Cancer Death in Surgically Treated Men with Clinically Localized Disease. *Eur Urol.* Jan 2018; 73(1): 129-138. PMID 28988753
64. Salmasi A, Said J, Shindel AW, et al. A 17-Gene Genomic Prostate Score Assay Provides Independent Information on Adverse Pathology in the Setting of Combined Multiparametric Magnetic Resonance Imaging Fusion Targeted and Systematic Prostate Biopsy. *J Urol.* Sep 2018; 200(3): 564-572. PMID 29524506
65. Cooperberg MR, Simko JP, Cowan JE, et al. Validation of a cell-cycle progression gene panel to improve risk stratification in a contemporary prostatectomy cohort. *J Clin Oncol.* Apr 10 2013; 31(11): 1428-34. PMID 23460710
66. McShane LM, Altman DG, Sauerbrei W, et al. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol.* Dec 20 2005; 23(36): 9067-72. PMID 16172462
67. Epstein JI, Allsbrook WC, Amin MB, et al. The 2005 International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading of Prostatic Carcinoma. *Am J Surg Pathol.* Sep 2005; 29(9): 1228-42. PMID 16096414

68. Brand TC, Zhang N, Crager MR, et al. Patient-specific Meta-analysis of 2 Clinical Validation Studies to Predict Pathologic Outcomes in Prostate Cancer Using the 17-Gene Genomic Prostate Score. *Urology*. Mar 2016; 89: 69-75. PMID 26723180
69. Albala D, Kemeter MJ, Febbo PG, et al. Health Economic Impact and Prospective Clinical Utility of Oncotype DX(R) Genomic Prostate Score. *Rev Urol*. 2016; 18(3): 123-132. PMID 27833462
70. Eure G, Germany R, Given R, et al. Use of a 17-Gene Prognostic Assay in Contemporary Urologic Practice: Results of an Interim Analysis in an Observational Cohort. *Urology*. Sep 2017; 107: 67-75. PMID 28454985
71. Badani KK, Kemeter MJ, Febbo PG, et al. The impact of a biopsy based 17-Gene Genomic Prostate Score on treatment recommendations in men with newly diagnosed clinically prostate cancer who are candidates for active surveillance. *Urol Pract*. 2015;2(4):181-189. PMID not Indexed in Pubmed
72. Canfield SK, M.J.; Febbo, P.G.; Hornberger, J. Balancing confounding and generalizability using observational, real-world data: 17-gene genomic prostate score assay effect on active surveillance. *Rev Urol*. 2018;20(2):69-76.
73. Canfield S, Kemeter MJ, Hornberger J, et al. Active Surveillance Use Among a Low-risk Prostate Cancer Population in a Large US Payer System: 17-Gene Genomic Prostate Score Versus Other Risk Stratification Methods. *Rev Urol*. 2017; 19(4): 203-212. PMID 29472824
74. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Med Decis Making*. Nov-Dec 2006; 26(6): 565-74. PMID 17099194
75. Cooperberg MR, Broering JM, Carroll PR. Risk assessment for prostate cancer metastasis and mortality at the time of diagnosis. *J Natl Cancer Inst*. Jun 16 2009; 101(12): 878-87. PMID 19509351
76. Berlin A, Murgic J, Hosni A, et al. Genomic Classifier for Guiding Treatment of Intermediate-Risk Prostate Cancers to Dose-Escalated Image Guided Radiation Therapy Without Hormone Therapy. *Int J Radiat Oncol Biol Phys*. Jan 01 2019; 103(1): 84-91. PMID 30170099
77. Nguyen PL, Shin H, Yousefi K, et al. Impact of a Genomic Classifier of Metastatic Risk on Postprostatectomy Treatment Recommendations by Radiation Oncologists and Urologists. *Urology*. Jul 2015; 86(1): 35-40. PMID 26142578
78. Blume-Jensen P, Berman DM, Rimm DL, et al. Development and clinical validation of an in situ biopsy-based multimarker assay for risk stratification in prostate cancer. *Clin Cancer Res*. Jun 01 2015; 21(11): 2591-600. PMID 25733599
79. Fossati N, Karnes RJ, Boorjian SA, et al. Long-term Impact of Adjuvant Versus Early Salvage Radiation Therapy in pT3N0 Prostate Cancer Patients Treated with Radical Prostatectomy: Results from a Multi-institutional Series. *Eur Urol*. Jun 2017; 71(6): 886-893. PMID 27484843
80. Hwang WL, Tendulkar RD, Niemierko A, et al. Comparison Between Adjuvant and Early-Salvage Postprostatectomy Radiotherapy for Prostate Cancer With Adverse Pathological Features. *JAMA Oncol*. May 10 2018; 4(5): e175230. PMID 29372236
81. Buscariollo DL, Drumm M, Niemierko A, et al. Long-term results of adjuvant versus early salvage postprostatectomy radiation: A large single-institutional experience. *Pract Radiat Oncol*. Mar 2017; 7(2): e125-e133. PMID 28274403
82. Freedland SJ, Rumble RB, Finelli A, et al. Adjuvant and salvage radiotherapy after prostatectomy: American Society of Clinical Oncology clinical practice guideline endorsement. *J Clin Oncol*. Dec 01 2014; 32(34): 3892-8. PMID 25366677
83. Stephenson AJ, Scardino PT, Kattan MW, et al. Predicting the outcome of salvage radiation therapy for recurrent prostate cancer after radical prostatectomy. *J Clin Oncol*. May 20 2007; 25(15): 2035-41. PMID 17513807
84. Stephenson AJ, Scardino PT, Eastham JA, et al. Postoperative nomogram predicting the 10-year probability of prostate cancer recurrence after radical prostatectomy. *J Clin Oncol*. Oct 01 2005; 23(28): 7005-12. PMID 16192588
85. Cooperberg MR, Hilton JF, Carroll PR. The CAPRA-S score: A straightforward tool for improved prediction of outcomes after radical prostatectomy. *Cancer*. Nov 15 2011; 117(22): 5039-46. PMID 21647869
86. Cuzick J, Swanson GP, Fisher G, et al. Prognostic value of an RNA expression signature derived from cell cycle proliferation genes in patients with prostate cancer: a retrospective study. *Lancet Oncol*. Mar 2011; 12(3): 245-55. PMID 21310658

87. Bishoff JT, Freedland SJ, Gerber L, et al. Prognostic utility of the cell cycle progression score generated from biopsy in men treated with prostatectomy. *J Urol.* Aug 2014; 192(2): 409-14. PMID 24508632
88. Koch MO, Cho JS, Kaimakliotis HZ, et al. Use of the cell cycle progression (CCP) score for predicting systemic disease and response to radiation of biochemical recurrence. *Cancer Biomark.* Jun 07 2016; 17(1): 83-8. PMID 27314296
89. Freedland SJ, Gerber L, Reid J, et al. Prognostic utility of cell cycle progression score in men with prostate cancer after primary external beam radiation therapy. *Int J Radiat Oncol Biol Phys.* Aug 01 2013; 86(5): 848-53. PMID 23755923
90. Klein EA, Yousefi K, Haddad Z, et al. A genomic classifier improves prediction of metastatic disease within 5 years after surgery in node-negative high-risk prostate cancer patients managed by radical prostatectomy without adjuvant therapy. *Eur Urol.* Apr 2015; 67(4): 778-86. PMID 25466945
91. Den RB, Yousefi K, Trabulsi EJ, et al. Genomic classifier identifies men with adverse pathology after radical prostatectomy who benefit from adjuvant radiation therapy. *J Clin Oncol.* Mar 10 2015; 33(8): 944-51. PMID 25667284
92. Den RB, Feng FY, Showalter TN, et al. Genomic prostate cancer classifier predicts biochemical failure and metastases in patients after postoperative radiation therapy. *Int J Radiat Oncol Biol Phys.* Aug 01 2014; 89(5): 1038-1046. PMID 25035207
93. Cooperberg MR, Davicioni E, Crisan A, et al. Combined value of validated clinical and genomic risk stratification tools for predicting prostate cancer mortality in a high-risk prostatectomy cohort. *Eur Urol.* Feb 2015; 67(2): 326-33. PMID 24998118
94. Ross AE, Feng FY, Ghadessi M, et al. A genomic classifier predicting metastatic disease progression in men with biochemical recurrence after prostatectomy. *Prostate Cancer Prostatic Dis.* Mar 2014; 17(1): 64-9. PMID 24145624
95. Karnes RJ, Bergstralh EJ, Davicioni E, et al. Validation of a genomic classifier that predicts metastasis following radical prostatectomy in an at risk patient population. *J Urol.* Dec 2013; 190(6): 2047-53. PMID 23770138
96. Erho N, Crisan A, Vergara IA, et al. Discovery and validation of a prostate cancer genomic classifier that predicts early metastasis following radical prostatectomy. *PLoS One.* 2013; 8(6): e66855. PMID 23826159
97. Ross AE, Johnson MH, Yousefi K, et al. Tissue-based Genomics Augments Post-prostatectomy Risk Stratification in a Natural History Cohort of Intermediate- and High-Risk Men. *Eur Urol.* Jan 2016; 69(1): 157-65. PMID 26058959
98. Freedland SJ, Choerung V, Howard L, et al. Utilization of a Genomic Classifier for Prediction of Metastasis Following Salvage Radiation Therapy after Radical Prostatectomy. *Eur Urol.* Oct 2016; 70(4): 588-596. PMID 26806658
99. Glass AG, Leo MC, Haddad Z, et al. Validation of a Genomic Classifier for Predicting Post-Prostatectomy Recurrence in a Community Based Health Care Setting. *J Urol.* Jun 2016; 195(6): 1748-53. PMID 26626216
100. Klein EA, Haddad Z, Yousefi K, et al. Decipher Genomic Classifier Measured on Prostate Biopsy Predicts Metastasis Risk. *Urology.* Apr 2016; 90: 148-52. PMID 26809071
101. Spratt DE, Dai DLY, Den RB, et al. Performance of a Prostate Cancer Genomic Classifier in Predicting Metastasis in Men with Prostate-specific Antigen Persistence Postprostatectomy. *Eur Urol.* Jul 2018; 74(1): 107-114. PMID 29233664
102. Karnes RJ, Choerung V, Ross AE, et al. Validation of a Genomic Risk Classifier to Predict Prostate Cancer-specific Mortality in Men with Adverse Pathologic Features. *Eur Urol.* Feb 2018; 73(2): 168-175. PMID 28400167
103. Ross AE, Den RB, Yousefi K, et al. Efficacy of post-operative radiation in a prostatectomy cohort adjusted for clinical and genomic risk. *Prostate Cancer Prostatic Dis.* Sep 2016; 19(3): 277-82. PMID 27136742
104. Spratt DE, Yousefi K, Deheshi S, et al. Individual Patient-Level Meta-Analysis of the Performance of the Decipher Genomic Classifier in High-Risk Men After Prostatectomy to Predict Development of Metastatic Disease. *J Clin Oncol.* Jun 20 2017; 35(18): 1991-1998. PMID 28358655
105. Lobo JM, Dicker AP, Buerki C, et al. Evaluating the clinical impact of a genomic classifier in prostate cancer using individualized decision analysis. *PLoS One.* 2015; 10(3): e0116866. PMID 25837660

106. West TA, Kiely BE, Stockler MR. Estimating scenarios for survival time in men starting systemic therapies for castration-resistant prostate cancer: a systematic review of randomised trials. *Eur J Cancer*. Jul 2014; 50(11): 1916-24. PMID 24825113
107. Scher HI, Graf RP, Schreiber NA, et al. Nuclear-specific AR-V7 Protein Localization is Necessary to Guide Treatment Selection in Metastatic Castration-resistant Prostate Cancer. *Eur Urol*. Jun 2017; 71(6): 874-882. PMID 27979426
108. Armstrong AJ, Halabi S, Luo J, et al. Prospective Multicenter Validation of Androgen Receptor Splice Variant 7 and Hormone Therapy Resistance in High-Risk Castration-Resistant Prostate Cancer: The PROPHECY Study. *J Clin Oncol*. May 01 2019; 37(13): 1120-1129. PMID 30865549
109. Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options. *J Urol*. Mar 2018; 199(3): 683-690. PMID 29203269
110. Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part II: Recommended Approaches and Details of Specific Care Options. *J Urol*. Apr 2018; 199(4): 990-997. PMID 29331546
111. Antonarakis ES, Lu C, Wang H, et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N Engl J Med*. Sep 11 2014; 371(11): 1028-38. PMID 25184630
112. Scher HI, Lu D, Schreiber NA, et al. Association of AR-V7 on Circulating Tumor Cells as a Treatment-Specific Biomarker With Outcomes and Survival in Castration-Resistant Prostate Cancer. *JAMA Oncol*. Nov 01 2016; 2(11): 1441-1449. PMID 27262168
113. Scher HI, Graf RP, Schreiber NA, et al. Assessment of the Validity of Nuclear-Localized Androgen Receptor Splice Variant 7 in Circulating Tumor Cells as a Predictive Biomarker for Castration-Resistant Prostate Cancer. *JAMA Oncol*. Sep 01 2018; 4(9): 1179-1186. PMID 29955787
114. Eggener SE, Rumble RB, Armstrong AJ, et al. Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. *J Clin Oncol*. May 01 2020; 38(13): 1474-1494. PMID 31829902
115. Lowrance WT, Murad MH, Oh WK, et al. Castration-Resistant Prostate Cancer: AUA Guideline Amendment 2018. *J Urol*. Dec 2018; 200(6): 1264-1272. PMID 30086276
116. National Institute for Health and Care Excellence (NICE). Prostate cancer: diagnosis and management [NG131]. 2019; <https://www.nice.org.uk/guidance/ng131>. Accessed October 14, 2020.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

