

**Medical Coverage Policy | Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer**



**EFFECTIVE DATE:** 10|01|2023

**POLICY LAST UPDATED:** 09|29|2023

## OVERVIEW

Various genetic and protein biomarkers are associated with prostate cancer. These tests have the potential to improve the accuracy of differentiating which men should undergo prostate biopsy or rebiopsy after a prior negative biopsy.

The following tests are addressed in this policy:

- ProgenSA PCA3 Assay (Hologic), CPT 81313
- 4Kscore (OPKO Health), CPT 81539
- ConfirmMDx (MDx Health), CPT 81551
- ExoDx Prostate IntelliScore (EPI) (Exosome Diagnostics), CPT 0005U
- Apify (Exact Sciences), CPT 0021U
- MyProstateScore (MPS) (Lynx Dx), CPT 0113U
- PanGIA Prostate (Genetics Institute of America), CPT 0228U
- SelectMDx (MDx Health), CPT 0339U
- Prostate Health Index (phi) (Beckman Coulter), CPT 84153, 84154, 86316
- Prostate Core Mitomics Test (Mitomics), Unlisted CPT

## MEDICAL CRITERIA

### Medicare Advantage Plans

#### ProgenSA PCA3, CPT 81313

PCA3 testing may be considered medically necessary when the following criteria are met:

- The test is ordered only once per year by a physician or other qualified health care professional (i.e., NP, CNS, PA), and;
  - For those men who need a repeat biopsy in the setting of patients thought to be at a higher risk despite a prior negative biopsy, with confirmed \* moderately elevated PSA
    - >3ng/ml and <10ng/ml; **OR**
    - >/4ng/ml and < 10ng/ml in men > 75 years of age with **BOTH** of the following:
1. No other relative indication for prostate biopsy including **ANY** of the following:
    - a. DRE suspicious for cancer (e.g., nodules, induration, or asymmetry)
    - b. Positive multiparametric MRI (Prostate Imaging Reporting and Data System [PI-RADS]  $\geq 3$ ) (if available)
    - c. Positive prior biopsy (cancer Grade Group  $\geq 1$ , intraductal carcinoma (IDC), atypical intraductal proliferation (AIP))
    - d. Other major risk factor for prostate cancer including:
      - i. Ethnicity at higher risk for prostate cancer
      - ii. First-degree relative with prostate cancer
      - iii. High-penetrance prostate cancer risk gene(s) per NCCN (if known)
  2. No other relative contraindication for prostate biopsy including **ANY** of the following:
    - a. <10 year life expectancy, or otherwise not a candidate for prostate cancer treatment
    - b. Invasive treatment for benign prostatic disease or taking medications that influence serum PSA levels within 6 mo.
    - c. Active prostatitis on antibiotics

\*PSA elevation should be confirmed after a few weeks under standardized conditions (i.e., no ejaculation, manipulations, and urinary tract infections) in the same laboratory before considering a biopsy.

#### **4Kscore, CPT 81539**

The 4Kscore test will be considered medically reasonable and necessary when all the following are met:

1. When all of the components of the algorithm are present.
2. Testing of men 45 years of age and older, prior to an initial biopsy or following a negative biopsy, who have a confirmed\* moderately elevated PSA (greater than 3 and less than 10 ng/mL; greater than or equal to 4 and less than 10 ng/mL in men greater than 75 years of age) when BOTH of the following are present:
  - No other relative indication\*\* for prostate biopsy including ANY of the following: (this may not be an all-inclusive list)
    - DRE suspicious for cancer should be encouraged to undergo biopsy
    - Persistent and significant increase in PSA should be encouraged to undergo biopsy
    - Positive multiparametric magnetic resonance imaging (MRI) (if done)
    - Other major risk factor for prostate cancer including: (this may not be an all inclusive list)
      - Ethnicity at higher risk for prostate cancer
      - First-degree relative with prostate cancer
      - High-penetrance prostate cancer risk gene(s) per the National Comprehensive Cancer Network (NCCN) (if known)
  - No other relative contraindication\*\* for prostate biopsy including ANY of the following:
    - Less than a 10-year life expectancy
    - Benign disease not ruled out.
3. Presence of shared decision making between the ordering provider and the beneficiary concerning the 4Kscore testing.

\* PSA elevation should be verified after a few weeks under standardized conditions (e.g. no ejaculation, manipulations, and urinary tract infections, no medications such as 5 $\alpha$ -reductase) in the same laboratory or other CLIA approved laboratory before considering a biopsy.

\*\* The relative indications and contraindications are not absolute. When it is determined that the 4Kscore test is medically reasonable and necessary in a beneficiary with one of the relative indications or contraindications for prostate biopsy the medical record must support the medical necessity for the test and there must be documented evidence of shared decision making between the patient and provider. This supporting documentation must be provided to the laboratory at the time of ordering the test.

#### **ConfirmMDx, CPT 81551**

#### **SelectMDx, CPT 0339U**

ConfirmMDx and SelectMDx may be considered medically necessary when ALL of the following criteria are met:

1. The patient must not have an established diagnosis of prostate cancer.
2. The beneficiary is a candidate for prostate biopsy or repeat prostate biopsy, according to a consensus guideline [(i.e., National Comprehensive Cancer Network® (NCCN), American Society of Clinical Oncology®(ASCO), American Urological Association (AUA)].
  - a. For men  $\leq$  75 years of age – Prostate Specific Antigen (PSA) (or adjusted PSA in special populations, i.e., patients taking 5alpha-reductase inhibitors) OR repeat PSA are  $>3$  and  $<10$ ng/mL AND/OR Digital Rectal Exam (DRE) findings are very suspicious for cancer
  - b. For men  $>$  75 years of age – PSA (or adjusted PSA in special populations, i.e., patients taking 5-alpha-reductase inhibitors) OR repeat PSA are  $\geq 4$  and  $<10$ ng/mL AND/OR DRE findings are very suspicious for cancer.

EXCEPTION: a molecular biomarker test may be performed in men with PSA levels  $>10$  ng/mL who are being considered for repeat biopsy IF appropriate according to consensus guidelines AND according to the following: the specific biomarker test has been validated in men with PSA levels  $>10$  ng/mL AND a Multiparametric MRI (mpMRI) is negative, *if performed*.

3. The beneficiary has not had a prostate biopsy OR has had a previous negative or non-malignant but abnormal histopathology finding (i.e., atypical small acinar proliferation (ASAP) or high-grade prostatic intraepithelial neoplasia (HGPIN) on prostate biopsy).
  - o Patients under consideration for a repeat biopsy have first undergone repeat PSA and/or DRE testing as recommended by consensus guidelines
4. The beneficiary would benefit from treatment of prostate cancer and patient management will be impacted by use of a biomarker in a manner already demonstrated in the peer-reviewed published literature to improve patient outcomes.
5. The medical record supports the medical necessity for the biomarker test.
6. Testing is performed according to the intended use of the test in the intended patient population for which the test was developed and validated.
7. Testing must be performed according to Clinical Laboratory Improvement Amendments (CLIA) and/or Food and Drug Administration (FDA) regulations in an accredited laboratory.
8. For a given clinical indication (pre-OR post-biopsy), only one molecular biomarker may be performed UNLESS a second test, meeting all the criteria established herein, is reasonable and necessary as an adjunct to the first test, according to criteria established in this policy.
9. If the test relies on an algorithm which may range in complexity from a threshold determination of a single numeric value to a complex mathematical or computational function, the algorithm must be validated in a cohort that is not a development cohort for the algorithm.
10. The analytes measured have demonstrated clinical validity and clinical utility (i.e., improved detection or discrimination of cancer or high-grade cancer or reduction in the need for biopsy) in the peer-reviewed published literature, establishing a clear and significant biological/molecular basis for stratifying patients and subsequently selecting (either positively or negatively) their clinical management decision within a clearly defined population.
11. The test is ordered by a physician specialist in the management of prostate cancer, such as a urologist or oncologist. An exception may be made in geographic locations where the specialist(s) cannot be reasonably reached by the beneficiary and the ordering provider is located closer to the beneficiary's place of residence than the nearest specialist. We would generally expect that beneficiaries for whom the test is ordered under this exception to be living in rural locations, islands, or some other location where access to care is limited.

#### **ExoDx Prostate IntelliScore (EPI), CPT 0005U**

EPI may be considered medically necessary when the following criteria are met:

- The test is ordered only once per year by a physician or other qualified health care professional (i.e., NP, CNS, PA), and;
- For men receiving testing prior to potential biopsy greater than or equal to 50 years of age, and;
- With a PSA of greater than 4ng/ml.

#### **Medicare Advantage Plans and Commercial Products**

Unless otherwise noted, for any test filed with an Unlisted CPT code, the medical necessity criteria in the Genetic Testing Services policy would be used. Please see the Related Policies section.

#### **PRIOR AUTHORIZATION**

##### **Medicare Advantage Plans**

Prior authorization is required for the following tests:

- PCA3 testing (eg, ProgenSA PCA3 Assay)
- 4Kscore
- ConfirmMDx
- ExoDx Prostate IntelliScore
- SelectMDx

#### **Medicare Advantage Plans and Commercial Products**

There is no specific CPT coding for some of the services referenced in this policy. Therefore, an Unlisted CPT code should be used (see Coding Section for details). All Unlisted genetic testing CPT codes require prior authorization to determine what service is being rendered and if the service is covered or not medically necessary. See the Related Policies section.

Prior authorization is required for Medicare Advantage Plans and recommended for Commercial Products and is obtained via the online tool for participating providers. See the Related Policies section.

**Note:** Laboratories are not allowed to obtain clinical authorization or participate in the authorization process on behalf of the ordering physician. Only the ordering physician shall be involved in the authorization, appeal or other administrative processes related to prior authorization/medical necessity.

In no circumstance shall a laboratory or a physician/provider use a representative of a laboratory or anyone with a relationship to a laboratory and/or a third party to obtain authorization on behalf of the ordering physician, to facilitate any portion of the authorization process or any subsequent appeal of a claim where the authorization process was not followed and/or a denial for clinical appropriateness was issued, including any element of the preparation of necessary documentation of clinical appropriateness. If a laboratory or a third party is found to be supporting any portion of the authorization process, BCBSRI will deem the action a violation of this policy and severe action will be taken up to and including termination from the BCBSRI provider network. If a laboratory provides a laboratory service that has not been authorized, the service will be denied as the financial liability of the participating laboratory and may not be billed to the member.

## **POLICY STATEMENT**

### **Medicare Advantage Plans**

The following test is covered but due to the instruction to file an Unlisted CPT code, prior authorization is required:

- Prostate Health Index (phi)

The following tests are considered medically necessary when the medical criteria above are met:

- PCA3 testing (eg, ProgenSA PCA3 Assay)
- 4Kscore
- ConfirmMDx
- ExoDx Prostate IntelliScore
- SelectMDx

The following genetic and protein biomarkers for the diagnosis of prostate cancer are not covered as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome:

- Apifyny
- MyProstateScore
- PanGIA Prostate
- Mitochondrial DNA variant testing (eg, Prostate Core Mitomics Test)
- Candidate gene panels

Single nucleotide variant testing for cancer risk assessment of prostate cancer is not covered as the evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

### **Commercial Products**

The following test is covered but due to the instruction to file an Unlisted CPT code, prior authorization is required:

- Prostate Health Index (phi)

The following genetic and protein biomarkers for the diagnosis of prostate cancer are considered not medically necessary as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome:

- PCA3 testing (eg, ProgenSA PCA3 Assay)
- Kallikrein markers (eg, 4Kscore® Test)
- Gene hypermethylation testing (eg, ConfirmMDx)
- PCA3, ERG, and SPDEF RNA expression in exosomes (eg, ExoDx Prostate IntelliScore)
- Autoantibodies ARF 6, NKX3-1, 5¢-UTR-BMI1, CEP 164, 3¢-UTR-Ropporin, Desmocollin, AURKAIP-1, and CSNK2A2 (eg, Apifyny)
- TMPRSS:ERG fusion genes (eg, MyProstateScore {MPS})
- PanGIA Prostate
- HOXC6 and DLX1 testing (eg, SelectMDx)
- Mitochondrial DNA variant testing (eg, Prostate Core Mitomics Test)
- Candidate gene panels

Single nucleotide variant testing for cancer risk assessment of prostate cancer is considered not medically necessary as the evidence is insufficient to determine that the technology results in an improvement in the net health outcomes.

## **COVERAGE**

Benefits may vary between groups and contracts. Please refer to the appropriate section of the Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable laboratory testing and not medically necessary/not covered benefits/coverage.

## **BACKGROUND**

Prostate cancer is the most common cancer, and the second most common cause of cancer death in men. Prostate cancer is a complex, heterogeneous disease, ranging from microscopic tumors unlikely to be life-threatening to aggressive tumors that can metastasize, leading to morbidity or death. Early localized disease can usually be treated with surgery and radiotherapy, although active surveillance may be adopted in men whose cancer is unlikely to cause major health problems during their lifespan or for whom the treatment might be dangerous. In patients with inoperable or metastatic disease, treatment consists of hormonal therapy and possibly chemotherapy. The lifetime risk of being diagnosed with prostate cancer for men in the United States is approximately 16%, while the risk of dying of prostate cancer is 3%. African-American men have the highest prostate cancer risk in the United States; the incidence of prostate cancer is about 60% higher and the mortality rate is more than 2 to 3 times greater than that of white men. Autopsy results have suggested that about 30% of men age 55 and 60% of men age 80 who die of other causes have incidental prostate cancer, indicating that many cases of cancer are unlikely to pose a threat during a man's life expectancy.

### **Grading**

The most widely used grading scheme for prostate cancer is the Gleason system. It is an architectural grading system ranging from 1 (well differentiated) to 5 (undifferentiated); the score is the sum of the primary and secondary patterns. A Gleason score of 6 or less is low-grade prostate cancer that usually grows slowly; 7 is an intermediate grade; 8 to 10 is high-grade cancer that grows more quickly. A revised prostate cancer grading system has been adopted by the National Cancer Institute and the World Health Organization.

Numerous genetic alterations associated with development or progression of prostate cancer have been described, with the potential for the use of these molecular markers to improve the selection process of men who should undergo prostate biopsy or rebiopsy after an initial negative biopsy.

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Laboratories that offer laboratory-developed tests must be licensed under the CLIA for high-complexity testing. The following laboratories are certified under the CLIA: BioReference Laboratories

and GenPath Diagnostics (subsidiaries of OPKO Health; 4Kscore®), ARUP Laboratories, Mayo Medical Laboratories, LabCorp, BioVantra, others (PCA3 assay), Clinical Research Laboratory (Prostate Core Mitomic Test™), MDx Health (SelectMDx, ConfirmMDx), Innovative Diagnostics (phi™), and ExoDx® Prostate (Exosome Diagnostics). To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of these tests.

In February 2012, the Progenesa® PCA3 Assay (Gen-Probe; now Hologic) was approved by the FDA through the premarket approval process. The Progenesa PCA3 Assay (Hologic Gen-Probe) has been approved by the FDA to aid in the decision for repeat biopsy in men 50 years or older who have had one or more negative prostate biopsies and for whom a repeat biopsy would be recommended based on current standard of care. The Progenesa PCA3 Assay should not be used for men with atypical small acinar proliferation on their most recent biopsy.

In June 2012, proPSA, a blood test used to calculate the Prostate Health Index (phi; Beckman Coulter) was approved by the FDA through the premarket approval process. The phi test is indicated as an aid to distinguish prostate cancer from a benign prostatic condition in men ages 50 and older with prostate-specific antigen levels of 4 to 10 ng/mL and with digital rectal exam findings that are not suspicious. According to the manufacturer, the test reduces the number of prostate biopsies.

For individuals who are being considered for an initial prostate biopsy who receive testing for genetic and protein biomarkers of prostate cancer (eg, kallikreins biomarkers and 4Kscore Test, proPSA and Prostate Health Index, TMRSS fusion genes and MyProstateScore, SelectMDx for Prostate Cancer, ExoDx Prostate, Apify, PCA3 score, and PanGIA Prostate), the evidence includes systematic reviews, meta-analyses, and primarily observational studies. Relevant outcomes are overall survival, disease-specific survival, test validity, resource utilization, and quality of life. The evidence supporting clinical utility varies by test but has not been directly shown for any biomarker test. Absent direct evidence of clinical utility, a chain of evidence might be constructed. However, the performance of biomarker testing for directing biopsy referrals is uncertain. While some studies have shown a reduction or delay in biopsy based on testing, a chain of evidence for clinical utility cannot be constructed due to limitations in clinical validity. Test validation populations have included men with a positive digital rectal exam, a prostate-specific antigen level outside of the gray zone (between 3 or 4 ng/mL and 10 ng/mL), or older men for whom the information from test results are less likely to be informative. Many biomarker tests do not have standardized cutoffs to recommend a biopsy. In addition, comparative studies of the many biomarkers are lacking. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are being considered for repeat biopsy who receive testing for genetic and protein biomarkers of prostate cancer (eg, PCA3 score, Gene Hypermethylation and ConfirmMDx test, Prostate Core Mitomics Test), the evidence includes systematic reviews and meta-analyses and primarily observational studies. Relevant outcomes are overall survival, disease-specific survival, test validity, resource utilization, and quality of life. The performance of biomarker testing for guiding rebiopsy decisions is lacking. The tests are associated with a diagnosis of prostate cancer and aggressive prostate cancer, but studies on clinical validity are limited and do not compare performance characteristics with standard risk prediction models. Direct evidence supporting clinical utility has not been shown. No data are currently available on physician decisions on rebiopsy or on the longer-term clinical outcomes of men who did not have a biopsy based on test results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Progenesa PCA3 Assay**

PCA3 is an mRNA expression assay tested from post-DRE urine. The FDA has approved the PCA3 assay to help decide whether a repeat biopsy in men aged 50 years or older with one or more previous negative prostate biopsies is necessary. In a prospective multicenter study of 466 men with a least one prior negative prostate biopsy scheduled for repeat biopsy based on best clinical judgement, a PCA3 score cutoff of 25 showed a sensitivity of 77.5%, specificity of 57.1%, NPV of 90%, and PPV of 33.6%. Those with a score of <25 were 4.56 times more likely to have a negative repeat biopsy.



#### **4Kscore**

The 4Kscore test measures blood levels of four Kallikreins protein biomarkers (total prostate-specific antigen [tPSA], free PSA [fPSA], intact PSA [iPSA], and human Kallikrein-related peptidase 2 [hK2]) in addition to other clinical information, including age, digital rectal examination (DRE) and prior biopsy history. All of these components are placed into a proprietary algorithm to provide a percent risk for a high-grade Gleason score greater than or equal to 7 cancer on biopsy. The 4Kscore test algorithm's goal is to refine patient selection for biopsies to reduce unnecessary biopsies in men being considered for biopsy of the prostate for potential cancer. The clinical features of this group of men are poorly defined.

#### **ConfirmMDx**

ConfirmMDx assesses the methylation status of 3 biomarkers (GSTP1, RASSF1, APC) associated with prostate cancer. ConfirmMDx is intended for use in patients with high-risk factors such as elevated/rising prostate-specific antigen (PSA) or abnormal digital rectal examination (DRE), with a negative or non-malignant abnormal histopathology finding (e.g., atypical cell or high grade prostate intraepithelial neoplasia (HGPIN)) in the previous biopsy, and is being considered for repeat biopsy. Several case/control studies in archived biopsy core tissue blocks demonstrated the sensitivity, specificity and high negative predictive value (NPV) of these biomarkers to predict cancer detection in a repeat biopsy procedure. Single biopsy cores, using as little as 20 microns from formalin-fixed, paraffin embedded (FFPE) tissue blocks or sections cut from blocks fixed on glass slides are used in this assay.

#### **ExoDx Prostate IntelliScore (EPI)**

EPI is a urine-based 3-gene exosomal RNA expression assay. The EPI gene signature and score incorporates levels of PCA3 (PCa antigen 3), ERG (v-ets erythroblastosis virus E26 oncogene homologs) and SPDEF (SAM-pointed domain-containing Ets transcription factor). EPI uses a proprietary algorithm to translate the level of expression of these three genes into an individualized risk score that predicts the presence of HGPC, with a higher EPI score indicative of a higher probability of high-grade disease. EPI does not incorporate PSA and other SOC factors into the score, but is intended to be used in conjunction with SOC elements such as age, family history, PSA level and DRE results.

#### **Apifyn**

Apifyn uses an algorithm to score the detection of 8 autoantibodies (ARF 6, NKX3-1, 5' -UTR-BMI1, CEP 164, 3' -UTR-Ropporin, Desmocollin, AURKAIP-1, CSNK2A2) in serum. The identified biomarkers play a role in processes such as androgen response regulation and cellular structural integrity and are proteins that are thought to play a role in prostate tumorigenesis.

#### **MyProstateScore**

MyProstateScore measures TMPRSS2-ERG gene fusion and calculates a probability score that incorporates serum PSA or the PCPT, and urine TMPRSS2-ERG and PCA3 scores

#### **PanGIA Prostate**

PanGIA Prostate is a urine test that uses a device with binding pockets for small molecules, proteins, and cells. Results are uploaded to the cloud and a machine learning algorithm compares the results with a signature from patients who have had a positive biopsy and patients who have had a negative prostate biopsy. The report includes a diagnosis with the level of confidence in the diagnosis.

#### **SelectMDx**

SelectMDx for prostate cancer uses a model that combines HOXC6 and DLX1 gene expression with traditional risk assessment models. HOXC6 and DLX1 mRNA is measured in post-DRE urine against kallikrein-related peptidase 3 as an internal reference.

#### **Prostate Health Index**

The Prostate Health Index (phi; Beckman Coulter) is an assay that combines results of 3 blood serum immunoassays (total PSA, free PSA, [-2]proPSA [p2PSA]) numerically to produce a "phi score." This score is calculated with the phi algorithm using the following formula:  $([-2]proPSA/free PSA) \times \sqrt{total PSA}$ . The phi

score is indicated for men 50 years and older with above-normal total PSA readings between 4.0 ng/mL and 10 ng/mL who have had a negative DRE in order to distinguish prostate cancer from benign prostatic conditions.

## **Medicare Advantage Plans**

### **Progensa PCA3 Assay and ExoDx Prostate IntelliScore (EPI)**

While the results of the mostly industry-sponsored validation studies are promising, benefits remain theoretical, namely, that fewer biopsies of men with moderately elevated PSA is inherently a good thing. Certainly, it is good in the short term for men who avoid an “unnecessary” prostate biopsy. Not good, however, are necessary biopsies missed due to false negatives. Moreover, even the definition of “unnecessary” may be evolving. Also, some studies overrepresented men for whom the information is less likely to be helpful (a positive DRE, PSA levels outside the gray zone, or older men not candidates for surgery), or underrepresented others (e.g., high risk groups such as African Americans, etc.). Comparative studies of the many biomarkers are lacking and it is unclear how to use the tests in practice, particularly when test results are contradictory (1). For all these reasons, the long-term benefit of these tests to net health outcomes (i.e., mortality, morbidity, or quality of life) is not yet clear.

### **4Kscore**

In their guidelines, the American Urological Association (AUA) recognizes that the decision to undergo PSA screening in men ages 55 to 69 involves weighing the benefits of reducing the rate of metastatic prostate cancer and prevention of prostate cancer death against the known potential harms associated with screening and treatment. For this reason, shared decision-making is recommended for men 55 to 69 years of age that are considering PSA screening and proceeding based on a man’s values and preferences.

The greatest benefit of screening appears to be in men 55 to 69 years of age. Multiple approaches subsequent to a PSA test (e.g., urinary and serum biomarkers, imaging, risk calculators) are available for identifying men more likely to harbor a prostate cancer and/or one with an aggressive phenotype. The use of such tools can be considered in men with a suspicious PSA level to inform prostate biopsy decisions.

The NCCN updated guidelines states that the 4Kscore test may be considered prior to biopsy for those with prior negative biopsy who are thought to be at risk for clinically significant prostate cancer. The NCCN notes that it is important for patients and their urologists to understand that no optimal cutoff threshold has been established for the 4Kscore test. It is recommended that the 4Kscore test may be considered before biopsy in men with serum PSA levels greater than 3ng/ml who desire more specificity.

With a recommendation for specific clinical scenarios, coupled with documented shared decision making, the AUA and NCCN organizations both state that biomarkers specifically 4Kscore “can be considered in men with a suspicious PSA level to inform prostate biopsy decisions.”

### **ConfirmMDx**

Biomarkers can help stratify men who have an elevated PSA into those more likely versus less likely to have aggressive disease. These non-invasive biomarker tests have demonstrated that they can (1) reduce the need for unnecessary biopsies in men unlikely to have prostate cancer or high-grade prostate cancer and/or (2) better define men at risk for higher-grade prostate cancer. There is adequate evidence to show that the incremental information provided by validated molecular biomarker tests for prostate cancer in samples of patients whose findings can be generalized to the Medicare population, changes physician management in a way that improves outcomes.

### **CODING**

The following CPT codes are covered for Medicare Advantage Plans when medical criteria above are met and are not medically necessary for Commercial Products.

CPT code 81313 is generally used to represent the Progensa® PCA3 Assay but can also be used for non-brand



name testing.

**81313** PCA3/KLK3 (prostate specific antigen 3 [non-protein coding]/kallikrein-related peptidase 3 [prostate specific antigen]) ratio (eg, prostate cancer)

This code can be used for 4Kscore® Test:

**81539** Oncology (high-grade prostate cancer), biochemical assay of four proteins (total PSA, free PSA, intact PSA and human kallikrein 2 [hK2]) plus patient age, digital rectal examination status, and no history of positive prostate biopsy, utilizing plasma, prognostic algorithm reported as a probability score

This code can be used for the ConfirmMDx® gene hypermethylation test.

**81551** Oncology (prostate), promoter methylation profiling by real-time PCR of 3 genes (GSTP1, APC, RASSF1), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a likelihood of prostate cancer detection on repeat biopsy

This code can be used for ExoDx Prostate IntelliScore (EPI):

**0005U** Oncology (prostate) gene expression profile by real-time RT-PCR of 3 genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score

This code can be used for SelectMDx:

**0339U** Oncology (prostate), mRNA expression profiling of HOXC6 and DLX1, reverse transcription polymerase chain reaction (RT-PCR), first-void urine following digital rectal examination, algorithm reported as probability of high-grade cancer (New Code Effective 10/1/2022)  
**(For Dates of Service prior to 10/1/2022, an Unlisted CPT code must be used.)**

The following CPT code(s) are not covered for Medicare Advantage Plans and are not medically necessary for Commercial Products.

This code can be used for Apifyny:

**0021U** Oncology (prostate), detection of 8 autoantibodies (ARF 6, NKX3-1, 5'-UTR-BMI1, CEP 164, 3'-UTR-Ropporin, Desmocollin, AURKAIP-1, CSNK2A2), multiplexed immunoassay and flow cytometry serum, algorithm reported as risk score

This code can be used for MyProstateScore (MPS):

**0113U** Oncology (prostate), measurement of PCA3 and TMPRSS2-ERG in urine and PSA in serum following prostatic massage, by RNA amplification and fluorescence-based detection, algorithm reported as risk score

This code can be used for PanGIA Prostate:

**0228U** Oncology (prostate), multianalyte molecular profile by photometric detection of macromolecules adsorbed on nanosponge array slides with machine learning, utilizing first morning voided urine, algorithm reported as likelihood of prostate cancer

The following Unlisted CPT code requires prior authorization for Medicare Advantage Plans and Commercial Products. The code can be used for any test identified in this policy that does not have a specific CPT code.

**81479** Unlisted molecular pathology procedure

## RELATED POLICIES

Genetic Testing Services

Medical Necessity

Proprietary Laboratory Analyses (PLA)

## PUBLISHED

Provider Update, February 2023, June 2023, August 2023

Provider Update, March 2022  
Provider Update, May 2020  
Provider Update, August 2019  
Provider Update, April 2019

## REFERENCES

1. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): 4Kscore Test Algorithm (L37792)
2. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: 4Kscore Test Algorithm (A56653)
3. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): Biomarker Testing (Prior to Initial Biopsy) for Prostate Cancer Diagnosis (L37733)
4. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: Biomarker Testing (Prior to Initial Biopsy) for Prostate Cancer Diagnosis (A56609)
5. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): MolDX: MolDX: Molecular Biomarkers to Risk-Stratify Patients at Increased Risk for Prostate Cancer (L39005)
6. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: MolDX: Molecular Biomarkers to Risk-Stratify Patients at Increased Risk for Prostate Cancer (A58718)
7. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2014. Bethesda, MD: National Cancer Institute; 2017.
8. Odedina FT, Akinremi TO, Chinegwundoh F, et al. Prostate cancer disparities in Black men of African descent: a comparative literature review of prostate cancer burden among Black men in the United States, Caribbean, United Kingdom, and West Africa. *Infect Agent Cancer*. Feb 10 2009; 4 Suppl 1: S2. PMID 19208207
9. Bell KJ, Del Mar C, Wright G, et al. Prevalence of incidental prostate cancer: A systematic review of autopsy studies. *Int J Cancer*. Oct 01 2015; 137(7): 1749-57. PMID 25821151
10. Gleason DF. Classification of prostatic carcinomas. *Cancer Chemother Rep*. Mar 1966; 50(3): 125-8. PMID 5948714
11. National Cancer Institute. SEER Database. <https://seer.cancer.gov/seerinqury/index.php?page=view&id=20170036&type=q>. Accessed October 6, 2021.
12. Hoogendam A, Buntinx F, de Vet HC. The diagnostic value of digital rectal examination in primary care screening for prostate cancer: a meta-analysis. *Fam Pract*. Dec 1999; 16(6): 621-6. PMID 10625141
13. Gosselaar C, Roobol MJ, Roemeling S, et al. The role of the digital rectal examination in subsequent screening visits in the European randomized study of screening for prostate cancer (ERSPC), Rotterdam. *Eur Urol*. Sep 2008; 54(3): 581-8. PMID 18423977
14. Thompson IM, Pauler DK, Goodman PJ, et al. Prevalence of prostate cancer among men with a prostate-specific antigen level or =4.0 ng per milliliter. *N Engl J Med*. May 27 2004; 350(22): 2239-46. PMID 15163773
15. Catalona WJ, Smith DS, Ratliff TL, et al. Measurement of prostate-specific antigen in serum as a screening test for prostate cancer. *N Engl J Med*. Apr 25 1991; 324(17): 1156-61. PMID 1707140
16. Aus G, Bergdahl S, Lodding P, et al. Prostate cancer screening decreases the absolute risk of being diagnosed with advanced prostate cancer--results from a prospective, population-based randomized controlled trial. *Eur Urol*. Mar 2007; 51(3): 659-64. PMID 16934392
17. Buzzoni C, Auvinen A, Roobol MJ, et al. Metastatic Prostate Cancer Incidence and Prostate-specific Antigen Testing: New Insights from the European Randomized Study of Screening for Prostate Cancer. *Eur Urol*. Nov 2015; 68(5): 885-90. PMID 25791513
18. Arnsrud Godtman R, Holmberg E, Lilja H, et al. Opportunistic testing versus organized prostate-specific antigen screening: outcome after 18 years in the Goteborg randomized population-based prostate cancer screening trial. *Eur Urol*. Sep 2015; 68(3): 354-60. PMID 25556937
19. Hugosson J, Carlsson S, Aus G, et al. Mortality results from the Goteborg randomised population-based prostate-cancer screening trial. *Lancet Oncol*. Aug 2010; 11(8): 725-32. PMID 20598634
20. Schroder FH, Hugosson J, Roobol MJ, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med*. Mar 26 2009; 360(13): 1320-8. PMID 19297566

21. Wolf AM, Wender RC, Etzioni RB, et al. American Cancer Society guideline for the early detection of prostate cancer: update 2010. *CA Cancer J Clin.* Mar-Apr 2010; 60(2): 70-98. PMID 20200110
22. Rosario DJ, Lane JA, Metcalfe C, et al. Short term outcomes of prostate biopsy in men tested for cancer by prostate specific antigen: prospective evaluation within ProtecT study. *BMJ.* Jan 09 2012; 344: d7894. PMID 22232535
23. Liss M, Ehdaie B, Loeb S, et al. The Prevention and Treatment of the More Common Complications Related to Prostate Biopsy Update. 2012; updated 2016; <https://www.aunet.org//guidelines/prostate-needle-biopsy-complications>. Accessed October 6, 2021.
24. Lavalley LT, Binette A, Witiuk K, et al. Reducing the Harm of Prostate Cancer Screening: Repeated Prostate-Specific Antigen Testing. *Mayo Clin Proc.* Jan 2016; 91(1): 17-22. PMID 26688045
25. Ruiz-Aragon J, Marquez-Pelaez S. [Assessment of the PCA3 test for prostate cancer diagnosis: a systematic review and meta-analysis]. *Actas Urol Esp.* Apr 2010; 34(4): 346-55. PMID 20470697
26. Mackinnon AC, Yan BC, Joseph LJ, et al. Molecular biology underlying the clinical heterogeneity of prostate cancer: an update. *Arch Pathol Lab Med.* Jul 2009; 133(7): 1033-40. PMID 19642730
27. Partin AW, Brawer MK, Subong EN, et al. Prospective evaluation of percent free-PSA and complexed-PSA for early detection of prostate cancer. *Prostate Cancer Prostatic Dis.* Jun 1998; 1(4): 197-203. PMID 12496895
28. Thompson IM, Ankerst DP, Chi C, et al. Assessing prostate cancer risk: results from the Prostate Cancer Prevention Trial. *J Natl Cancer Inst.* Apr 19 2006; 98(8): 529-34. PMID 16622122
29. van Vugt HA, Roobol MJ, Kranse R, et al. Prediction of prostate cancer in unscreened men: external validation of a risk calculator. *Eur J Cancer.* Apr 2011; 47(6): 903-9. PMID 21163642
30. Rosenkrantz AB, Verma S, Choyke P, et al. Prostate Magnetic Resonance Imaging and Magnetic Resonance Imaging Targeted Biopsy in Patients with a Prior Negative Biopsy: A Consensus Statement by AUA and SAR. *J Urol.* Dec 2016; 196(6): 1613-1618. PMID 27320841
31. Mi C, Bai L, Yang Y, et al. 4Kscore diagnostic value in patients with high-grade prostate cancer using cutoff values of 7.5% to 10%: A meta-analysis. *Urol Oncol.* Jun 2021; 39(6): 366.e1-366.e10. PMID 33685800
32. Russo GI, Regis F, Castelli T, et al. A Systematic Review and Meta-analysis of the Diagnostic Accuracy of Prostate Health Index and 4-Kallikrein Panel Score in Predicting Overall and High-grade Prostate Cancer. *Clin Genitourin Cancer.* Aug 2017; 15(4): 429-439.e1. PMID 28111174
33. Parekh DJ, Punnen S, Sjoberg DD, et al. A multi-institutional prospective trial in the USA confirms that the 4Kscore accurately identifies men with high-grade prostate cancer. *Eur Urol.* Sep 2015; 68(3): 464-70. PMID 25454615
34. Punnen S, Freedland SJ, Polascik TJ, et al. A Multi-Institutional Prospective Trial Confirms Noninvasive Blood Test Maintains Predictive Value in African American Men. *J Urol.* Jun 2018; 199(6): 1459-1463. PMID 29223389
35. Bhattu AS, Zappala SM, Parekh DJ, et al. A 4Kscore Cut-off of 7.5% for Prostate Biopsy Decisions Provides High Sensitivity and Negative Predictive Value for Significant Prostate Cancer. *Urology.* Feb 2021; 148: 53-58. PMID 33217456
36. Stattin P, Vickers AJ, Sjoberg DD, et al. Improving the Specificity of Screening for Lethal Prostate Cancer Using Prostate-specific Antigen and a Panel of Kallikrein Markers: A Nested Case-Control Study. *Eur Urol.* Aug 2015; 68(2): 207-13. PMID 25682340
37. Loeb S, Shin SS, Broyles DL, et al. Prostate Health Index improves multivariable risk prediction of aggressive prostate cancer. *BJU Int.* Jul 2017; 120(1): 61-68. PMID 27743489
38. Konety B, Zappala SM, Parekh DJ, et al. The 4Kscore(R) Test Reduces Prostate Biopsy Rates in Community and Academic Urology Practices. *Rev Urol.* 2015; 17(4): 231-40. PMID 26839521
39. Pecoraro V, Roli L, Plebani M, et al. Clinical utility of the (-2)proPSA and evaluation of the evidence: a systematic review. *Clin Chem Lab Med.* Jul 01 2016; 54(7): 1123-32. PMID 26609863
40. Anyango R, Ojwando J, Mwita C, et al. Diagnostic accuracy of [-2]proPSA versus Gleason score and Prostate Health Index versus Gleason score for the determination of aggressive prostate cancer: a systematic review. *JBI Evid Synth.* Mar 17 2021; 19(6): 1263-1291. PMID 33741840
41. Catalona WJ, Partin AW, Sanda MG, et al. A multicenter study of [-2]pro-prostate specific antigen combined with prostate specific antigen and free prostate specific antigen for prostate cancer detection in the 2.0 to 10.0 ng/ml prostate specific antigen range. *J Urol.* May 2011; 185(5): 1650-5. PMID 21419439

42. Tosoian JJ, Druskin SC, Andreas D, et al. Use of the Prostate Health Index for detection of prostate cancer: results from a large academic practice. *Prostate Cancer Prostatic Dis.* Jun 2017; 20(2): 228-233. PMID 28117387
43. White J, Shenoy BV, Tutrone RF, et al. Clinical utility of the Prostate Health Index (phi) for biopsy decision management in a large group urology practice setting. *Prostate Cancer Prostatic Dis.* Apr 2018; 21(1): 78-84. PMID 29158509
44. Sanda MG, Feng Z, Howard DH, et al. Association Between Combined TMPRSS2:ERG and PCA3 RNA Urinary Testing and Detection of Aggressive Prostate Cancer. *JAMA Oncol.* Aug 01 2017; 3(8): 1085-1093. PMID 28520829
45. Tomlins SA, Day JR, Lonigro RJ, et al. Urine TMPRSS2:ERG Plus PCA3 for Individualized Prostate Cancer Risk Assessment. *Eur Urol.* Jul 2016; 70(1): 45-53. PMID 25985884
46. Ankerst DP, Goros M, Tomlins SA, et al. Incorporation of Urinary Prostate Cancer Antigen 3 and TMPRSS2:ERG into Prostate Cancer Prevention Trial Risk Calculator. *Eur Urol Focus.* Jan 2019; 5(1): 54-61. PMID 29422418
47. Tosoian JJ, Trock BJ, Morgan TM, et al. Use of the MyProstateScore Test to Rule Out Clinically Significant Cancer: Validation of a Straightforward Clinical Testing Approach. *J Urol.* Mar 2021; 205(3): 732-739. PMID 33080150
48. Newcomb LF, Zheng Y, Faino AV, et al. Performance of PCA3 and TMPRSS2:ERG urinary biomarkers in prediction of biopsy outcome in the Canary Prostate Active Surveillance Study (PASS). *Prostate Cancer Prostatic Dis.* Sep 2019; 22(3): 438-445. PMID 30664734
49. Van Neste L, Hendriks RJ, Dijkstra S, et al. Detection of High-grade Prostate Cancer Using a Urinary Molecular Biomarker-Based Risk Score. *Eur Urol.* Nov 2016; 70(5): 740-748. PMID 27108162
50. Haese A, Trooskens G, Steyaert S, et al. Multicenter Optimization and Validation of a 2-Gene mRNA Urine Test for Detection of Clinically Significant Prostate Cancer before Initial Prostate Biopsy. *J Urol.* Aug 2019; 202(2): 256-263. PMID 31026217
51. McKiernan J, Donovan MJ, O'Neill V, et al. A Novel Urine Exosome Gene Expression Assay to Predict High-grade Prostate Cancer at Initial Biopsy. *JAMA Oncol.* Jul 01 2016; 2(7): 882-9. PMID 27032035
52. Tutrone R, Donovan MJ, Torkler P, et al. Clinical utility of the exosome based ExoDx Prostate(IntelliScore) EPI test in men presenting for initial Biopsy with a PSA 2-10 ng/mL. *Prostate Cancer Prostatic Dis.* Dec 2020; 23(4): 607-614. PMID 32382078
53. Schipper M, Wang G, Giles N, et al. Novel prostate cancer biomarkers derived from autoantibody signatures. *Transl Oncol.* Apr 2015; 8(2): 106-11. PMID 25926076
54. Wysock JS, Becher E, Persily J, et al. Concordance and Performance of 4Kscore and SelectMDx for Informing Decision to Perform Prostate Biopsy and Detection of Prostate Cancer. *Urology.* Jul 2020; 141: 119-124. PMID 32294481
55. Cui Y, Cao W, Li Q, et al. Evaluation of prostate cancer antigen 3 for detecting prostate cancer: a systematic review and meta-analysis. *Sci Rep.* May 10 2016; 6: 25776. PMID 27161545
56. Rodriguez SVM, Garcia-Perdomo HA. Diagnostic accuracy of prostate cancer antigen 3 (PCA3) prior to first prostate biopsy: A systematic review and meta-analysis. *Can Urol Assoc J.* May 2020; 14(5): E214-E219. PMID 31793864
57. Nicholson A, Mahon J, Boland A, et al. The clinical effectiveness and cost-effectiveness of the PROGENSA(R) prostate cancer antigen 3 assay and the Prostate Health Index in the diagnosis of prostate cancer: a systematic review and economic evaluation. *Health Technol Assess.* Oct 2015; 19(87): i-xxxi, 1-191. PMID 26507078
58. Wei JT, Feng Z, Partin AW, et al. Can urinary PCA3 supplement PSA in the early detection of prostate cancer?. *J Clin Oncol.* Dec 20 2014; 32(36): 4066-72. PMID 25385735
59. Hennenlotter J, Neumann T, Alperowitz S, et al. Age-Adapted Prostate Cancer Gene 3 Score Interpretation - Suggestions for Clinical Use. *Clin Lab.* Mar 01 2020; 66(3). PMID 32162868
60. Vickers AJ, Gupta A, Savage CJ, et al. A panel of kallikrein marker predicts prostate cancer in a large, population-based cohort followed for 15 years without screening. *Cancer Epidemiol Biomarkers Prev.* Feb 2011; 20(2): 255-61. PMID 21148123
61. Ruffion A, Devonec M, Champetier D, et al. PCA3 and PCA3-based nomograms improve diagnostic accuracy in patients undergoing first prostate biopsy. *Int J Mol Sci.* Aug 29 2013; 14(9): 17767-80. PMID 23994838



62. Ruffion A, Perrin P, Devonec M, et al. Additional value of PCA3 density to predict initial prostate biopsy outcome. *World J Urol.* Aug 2014; 32(4): 917-23. PMID 24500192
63. Merdan S, Tomlins SA, Barnett CL, et al. Assessment of long-term outcomes associated with urinary prostate cancer antigen 3 and TMPRSS2:ERG gene fusion at repeat biopsy. *Cancer.* Nov 15 2015; 121(22): 4071-9. PMID 26280815
64. Djavan B, Waldert M, Zlotta A, et al. Safety and morbidity of first and repeat transrectal ultrasound guided prostate needle biopsies: results of a prospective European prostate cancer detection study. *J Urol.* Sep 2001; 166(3): 856-60. PMID 11490233
65. Lujan M, Paez A, Santonja C, et al. Prostate cancer detection and tumor characteristics in men with multiple biopsy sessions. *Prostate Cancer Prostatic Dis.* 2004; 7(3): 238-42. PMID 15289810
66. Stewart GD, Van Neste L, Delvenne P, et al. Clinical utility of an epigenetic assay to detect occult prostate cancer in histopathologically negative biopsies: results of the MATLOC study. *J Urol.* Mar 2013; 189(3): 1110-6. PMID 22999998
67. Partin AW, Van Neste L, Klein EA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol.* Oct 2014; 192(4): 1081-7. PMID 24747657
68. Waterhouse RL, Van Neste L, Moses KA, et al. Evaluation of an Epigenetic Assay for Predicting Repeat Prostate Biopsy Outcome in African American Men. *Urology.* Jun 2019; 128: 62-65. PMID 29660369
69. Van Neste L, Partin AW, Stewart GD, et al. Risk score predicts high-grade prostate cancer in DNA-methylation positive, histopathologically negative biopsies. *Prostate.* Sep 2016; 76(12): 1078-87. PMID 27121847
70. Partin AW, VAN Criekinge W, Trock BJ, et al. CLINICAL EVALUATION OF AN EPIGENETIC ASSAY TO PREDICT MISSED CANCER IN PROSTATE BIOPSY SPECIMENS. *Trans Am Clin Climatol Assoc.* 2016; 127: 313-327. PMID 28066067
71. Food and Drug Administration. Summary of Safety and Effectiveness Data. PMA P090026. Quantitative test for determination of [-2]proPSA levels. Silver Spring, MD: Food and Drug Administration; 2012.
72. Aubry W, Lieberthal R, Willis A, et al. Budget impact model: epigenetic assay can help avoid unnecessary repeated prostate biopsies and reduce healthcare spending. *Am Health Drug Benefits.* Jan 2013; 6(1): 15-24. PMID 24991343
73. Robinson K, Creed J, Reguly B, et al. Accurate prediction of repeat prostate biopsy outcomes by a mitochondrial DNA deletion assay. *Prostate Cancer Prostatic Dis.* Jun 2010; 13(2): 126-31. PMID 20084081
74. Legisi L, DeSa E, Qureshi MN. Use of the Prostate Core Mitomic Test in Repeated Biopsy Decision-Making: Real-World Assessment of Clinical Utility in a Multicenter Patient Population. *Am Health Drug Benefits.* Dec 2016; 9(9): 497-502. PMID 28465777
75. Leyten GH, Hessels D, Smit FP, et al. Identification of a Candidate Gene Panel for the Early Diagnosis of Prostate Cancer. *Clin Cancer Res.* Jul 01 2015; 21(13): 3061-70. PMID 25788493
76. Xiao K, Guo J, Zhang X, et al. Use of two gene panels for prostate cancer diagnosis and patient risk stratification. *Tumour Biol.* Aug 2016; 37(8): 10115-22. PMID 26820133
77. American Urological Association. Detection of prostate cancer. 2013 (Reviewed and Validity Confirmed 2018); <http://www.auanet.org/education/guidelines/prostate-cancer-detection.cfm>. Accessed October 6, 2021.
78. National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology: prostate cancer early detection. [http://www.nccn.org/professionals/physician\\_gls/pdf/prostate\\_detection.pdf](http://www.nccn.org/professionals/physician_gls/pdf/prostate_detection.pdf). Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for NCCN clinical practice guidelines in oncology: prostate cancer early detection V.2.2021 National Comprehensive Cancer Network rights reserved. Accessed October 6, 2021.
79. National Institute for Health and Care Excellence (NICE). Prostate cancer: diagnosis and management [NG131]. 2019; <https://www.nice.org.uk/guidance/ng131/chapter/Recommendations#assessment-and-diagnosis>. Accessed October 6, 2021.
80. U. S. Preventive Services Task Force. Prostate Cancer: Screening. 2018; <https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/prostate-cancer-screening1>. Accessed October 6, 2021.



**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

