

Medical Coverage Policy | Interspinous and Interlaminar Stabilization/Distraction Devices (Spacers)



EFFECTIVE DATE: 03|01|2018

POLICY LAST UPDATED: 02|05|2019

OVERVIEW

Interspinous and interlaminar implants (spacers) stabilize or distract the adjacent lamina and/or spinous processes and restrict extension to reduce pain in patients with lumbar spinal stenosis and neurogenic claudication. Interspinous spacers are small devices implanted between the vertebral spinous processes. After implantation, the device is opened or expanded to distract (open) the neural foramen and decompress the nerves. Interlaminar spacers are implanted midline between adjacent lamina and spinous processes to provide dynamic stabilization either following decompressive surgery or as an alternative to decompressive surgery.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

BlueCHiP for Medicare and Commercial Products

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare

Interspinous or interlaminar distraction devices as a stand-alone procedure are not covered as a treatment of spinal stenosis as the evidence is insufficient to determine the effects of the technology on health outcomes.

Use of an interlaminar stabilization device following decompression surgery is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

Removal for medical reasons (device failure, infection, etc.) is covered for all members. However, insertion of a replacement device after removal is not covered.

Commercial Products

Interspinous or interlaminar distraction devices as a stand-alone procedure are not medically necessary as a treatment of spinal stenosis as the evidence is insufficient to determine the effects of the technology on health outcomes.

Use of an interlaminar stabilization device following decompression surgery is not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

Removal for medical reasons (device failure, infection, etc.) is covered for all members. However, insertion of a replacement device after removal is not medically necessary.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Spinal Stenosis

Lumbar spinal stenosis (LSS), which affects over 200,000 people in the United States, involves a narrowed central spinal canal, lateral spinal recesses, and/or neural foramina, resulting in pain as well as limitation of activities such as walking, traveling, and standing. In adults over 60 in the United States, spondylosis (degenerative arthritis affecting the spine) is the most common cause. The primary symptom of LSS is neurogenic claudication with back and leg pain, sensory loss, and weakness in the legs. Symptoms are typically exacerbated by standing or walking and relieved with sitting or flexion at the waist.

Some sources describe the course of LSS as “progressive” or “degenerative,” implying that neurologic decline is the usual course. Longer term data from the control groups of clinical trials as well as from observational studies suggest that, over time, most patients remain stable, some improve, and some deteriorate.

The lack of a valid classification for LSS contributes to wide practice variation and uncertainty about who should be treated surgically and which surgical procedure is best for each patient. This uncertainty also complicates research on spinal stenosis, particularly the selection of appropriate eligibility criteria and comparators.

Treatment

Appropriate surgical treatments for patients with spinal stenosis not responding to conservative treatments include decompression with or without spinal fusion. There are many types of decompression surgery and types of fusion operations. In general, spinal fusion is associated with more complications and a longer recovery period and, in the past, was generally reserved for patients with spinal deformity or moderate grade spondylolisthesis.

Conservative treatment for spinal stenosis may include physical therapy, pharmacotherapy, epidural steroid injections, and many other modalities. The terms “nonsurgical” and “nonoperative” have also been used to describe conservative treatment. Professional societies recommend that surgery for LSS should be considered only after a patient fails to respond to conservative treatment, but there is no agreement about what constitutes an adequate course or duration of treatment.

The term “conservative management” may refer to “usual care” or to specific programs of nonoperative treatment, which use defined protocols for the components and intensity of conservative treatments, often in the context of an organized program of coordinated, multidisciplinary care. The distinction is important in defining what constitutes a failure of conservative treatment and what comparators should be used in trials of surgical vs nonsurgical management. The rationale for surgical treatment of symptomatic spinal stenosis rests on the Spine Patient Outcomes Research Trial (SPORT), which found that patients who underwent surgery for spinal stenosis and spondylolisthesis had better outcomes than those treated nonoperatively. The SPORT investigators did not require a specified program of nonoperative care but rather let each site decide what to offer. A subgroup analysis of the SPORT trial found that only 37% of nonsurgically treated patients received physical therapy in the first 6 weeks of the trial and that those who received physical therapy before 6 weeks had better functional outcomes and were less likely to cross over to surgery later. These findings provide some support for the view that, in clinical trials, patients who did not have surgery may have had suboptimal treatment, which can lead to a larger difference favoring surgery. The SPORT investigators asserted that their nonoperative outcomes represented typical results at a multidisciplinary spine center at the time, but recommended that future studies compare the efficacy of specific nonoperative programs to surgery.

A recent trial by Delitto et al (2015) compared surgical decompression with a specific therapy program emphasizing physical therapy and exercise. Patients with lumbar spinal stenosis and from 0 to 5 mm of slippage (spondylolisthesis) who were willing to be randomized to decompression surgery vs an intensive, organized program of nonsurgical therapy were eligible. Oswestry Disability Index scores were comparable to those in the SPORT trial. A high proportion of patients assigned to nonsurgical care (57%) crossed over to surgery (in SPORT the proportion was 43%), but crossover from surgery to nonsurgical care was minimal.

When analyzed by treatment assignment, Oswestry Disability Index scores were similar in the surgical and nonsurgical groups after 2 years of follow-up. The main implication is that about one-third of patients who were deemed candidates for decompression surgery but instead entered an intensive program of conservative care achieved outcomes similar to those of a successful decompression.

Diagnostic criteria for fusion surgery are challenging because patients without spondylolisthesis and those with grade 1 spondylolisthesis are equally likely to have predominant back pain or predominant leg pain. The SPORT trial did not provide guidance on which surgery is appropriate for patients who do not have spondylolisthesis, because nearly all patients with spondylolisthesis underwent fusion whereas nearly all those who did not have spondylolisthesis underwent decompression alone. In general, patients with predominant back pain have more severe symptoms, worse function, and less improvement with surgery (with or without fusion). Moreover, because back pain improved to the same degree for the fused spondylolisthesis patients as for the unfused spinal stenosis patients at 2 years, the SPORT investigators concluded that it was unlikely that fusion led to the better surgical outcomes in patients with spondylolisthesis than those with no spondylolisthesis.

Throughout the 2000s, decompression plus fusion became more widely used until, in 2011, it surpassed decompression alone as a surgical treatment for spinal stenosis. However, in 2016, findings from two randomized trials of decompression alone vs decompression plus fusion were published. The Swedish Spinal Stenosis Study (SSSS) found no benefit of fusion plus decompression compared with decompression alone in patients who had spinal stenosis with or without degenerative spondylolisthesis. The Spinal Laminectomy versus Instrumented Pedicle Screw (SLIP) trial found a small but clinically meaningful improvement in the Physical Component Summary score of the 36-Item Short-Form Health Survey but no change in Oswestry Disability Index scores at 2, 3, and 4 years in patients who had spinal stenosis with grade 1 spondylolisthesis (3-14 mm). The patients in SLIP who had laminectomy alone had higher reoperation rates than those in SSSS, and the patients who underwent fusion had better outcomes in SLIP than in SSSS. While some interpret the studies to reflect differences in patient factors—in particular, SSSS but not SLIP included patients with no spondylolisthesis, the discrepancy may also be influenced by factors such as time of follow-up or national practice patterns. As Pearson (2016) noted, it might have been helpful to have patient-reported outcome data on the patients before and after reoperation, to see whether the threshold for reoperation differed in the 2 settings. A small trial conducted in Japan, Inose et al (2018) found no difference in patient-reported outcomes between laminectomy alone and laminectomy plus posterolateral fusion in patients with 1-level spinal stenosis and grade 1 spondylolisthesis; about 40% of the patients also had dynamic instability. Certainty in the findings of this trial is limited because of its size and methodologic flaws.

Spacer Devices

Investigators have sought less invasive ways to stabilize the spine and reduce the pressure on affected nerve roots, including interspinous and interlaminar implants (spacers). These devices stabilize or distract the adjacent lamina and/or spinous processes and restrict extension in patients with lumbar spinal stenosis and neurogenic claudication.

Other types of dynamic posterior stabilization devices are pedicle screw/rod-based devices and total facet replacement systems; they are not discussed in this evidence review.

Interspinous Implants

Interspinous spacers are small devices implanted between the vertebral spinous processes. After implantation, the device is opened or expanded to distract the neural foramina and decompress the nerves. One type of interspinous implant is inserted between the spinous processes through a small (4-8 cm) incision and acts as a spacer between the spinous processes, maintaining flexion of that spinal interspace. The supraspinous ligament is maintained and assists in holding the implant in place. The surgery does not include any laminotomy, laminectomy, or foraminotomy at the time of insertion, thus reducing the risk of epidural

scarring and cerebrospinal fluid leakage. Other interspinous spacers require removal of the interspinous ligament and are secured around the upper and lower spinous processes.

Interlaminar Spacers

Interlaminar spacers are implanted midline between adjacent lamina and spinous processes to provide dynamic stabilization either following decompression surgery or as an alternative to decompression surgery. Interlaminar spacers have 2 sets of wings placed around the inferior and superior spinous processes. They may also be referred to as interspinous U. These implants aim to restrict painful motion while enabling normal motion. The devices (spacers) distract the laminar space and/or spinous processes and restrict extension. This procedure theoretically enlarges the neural foramen and decompresses the cauda equina in patients with spinal stenosis and neurogenic claudication.

The Superior® Indirect Decompression System (formerly InterSpinous Spacer) is indicated to treat skeletally mature patients suffering from pain, numbness, and/or cramping in the legs secondary to a diagnosis of moderate degenerative lumbar spinal stenosis, with or without grade 1 spondylolisthesis, confirmed by x-ray, magnetic resonance imaging, and/or computed tomography evidence of thickened ligamentum flavum, narrowed lateral recess, and/or central canal or foraminal narrowing. It is intended for patients with impaired physical function who experience relief in flexion from symptoms of leg/buttock/groin pain, numbness, and/or cramping, with or without back pain, and who have undergone at least 6 months of nonoperative treatment.

The coflex® Interlaminar Technology implant (Paradigm Spine) is a single-piece U-shaped titanium alloy dynamic stabilization device with pairs of wings that surround the superior and inferior spinous processes. The coflex® (previously called the Interspinous U) is indicated for use in 1- or 2- level lumbar stenosis from the L1 to L5 vertebrae in skeletally mature patients with at least moderate impairment in function, who experience relief in flexion from their symptoms of leg/buttocks/groin pain, with or without back pain, and who have undergone at least 6 months of nonoperative treatment. The coflex® "is intended to be implanted midline between adjacent lamina of 1 or 2 contiguous lumbar motion segments. Interlaminar stabilization is performed after decompression of stenosis at the affected level(s)."

The literature is dominated by reports from non-U.S. centers evaluating devices not approved by the U.S. Food and Drug Administration (FDA), although a number of them are in trials at U.S. centers. As of April 2018, only the X-STOP, coflex, and Superior Interspinous Spacer (ISS) devices had received FDA approval for use in the United States. Manufacturing of the X-STOP device stopped in 2015.

For individuals who have spinal stenosis and no spondylolisthesis or grade 1 spondylolisthesis who receive an interspinous or interlaminar spacer as a stand-alone procedure, the evidence includes 2 randomized controlled trials of 2 spacers (Superior Interspinous Spacer, coflex interlaminar implant). Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Overall, the use of interspinous or interlaminar distraction devices (spacers) as an alternative to spinal decompression has shown a high failure and complication rates. A pivotal trial compared the Superior Interspinous Spacer with the X-STOP (which is no longer marketed), without conservative care or standard surgery comparators. The trial reported significantly better outcomes with the Superior Interspinous Spacer on some measures. For example, the trial reported more than 80% of patients experienced improvements in certain quality of life outcome domains. Interpretation of this trial is limited by questions about the number of patients used to calculate success rates, the lack of efficacy of the comparator, and the lack of an appropriate control group treated by surgical decompression. The coflex interlaminar implant (formerly called the interspinous U) was compared with decompression in the multicenter, double-blind Foraminal Enlargement Lumbar Interspinous distraX ion trial. Functional outcomes and pain levels were similar in the 2 groups at 1-year follow-up, but reoperation rates due to the absence of recovery were substantially higher with the coflex implant (29%) than with bony decompression (8%). For patients with 2-level surgery, the reoperation rate was 38% for coflex and

6% for bony decompression. At 2 years, reoperations due to the absence of recovery had been performed in 33% of the coflex group and 8% of the bony decompression group. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have spinal stenosis and no spondylolisthesis or grade 1 spondylolisthesis who receive an interlaminar spacer with spinal decompression surgery, the evidence includes randomized controlled trials and nonrandomized comparative studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Use of the coflex interlaminar implant as a stabilizer after surgical decompression has been studied in 2 situations^{3/4}as an adjunct to decompression compared with decompression alone (superiority) and as an alternative to spinal fusion after decompression (noninferiority). In a randomized controlled trial conducted in a patient population with moderate-to-severe lumbar spinal stenosis with significant back pain and up to grade 1 spondylolisthesis, there was no difference in the primary outcome measure, the Oswestry Disability Index (ODI), between the patients treated with coflex plus decompression vs. decompression alone. "Composite clinical success" (CCS), defined as a minimum 15-point improvement in ODI score, no reoperations, no device-related complications, no epidural steroid injections in the lumbar spine, and no persistent new or worsening sensory or motor deficit, was used to assess superiority. A greater proportion of patients who received coflex plus decompression instead of decompression alone achieved the composite endpoint. However, the superiority of coflex plus decompression is uncertain because the difference in the CCS was primarily driven by a greater proportion of patients in the control arm who received a secondary rescue epidural steroid injection. Because the trial was open-label, surgeons' decision to use epidural steroid injection could have been affected by their knowledge of the patient's treatment. Consequently, including this component in the composite clinical success measure might have overestimated the potential benefit of treatment. This bias could have been mitigated using protocol-mandated standard objective clinical criteria to guide decisions about secondary interventions and subsequent adjudication of these events by an independent blinded committee. Greater certainty about the net health outcome of adding coflex to decompression surgery might be demonstrated when the 5-year follow-up results of these trials and an ongoing trial (NCT02555280) on decompression with and without the coflex implant in the United States are published. To be useful for clinical decision-making, this study should report the patient-reported effectiveness measures for both back pain (ODI and/or back visual analog scale) and the claudication (Zurich Claudication Questionnaire and/or leg visual analog scale) in all patients at 5 years. The evidence is insufficient to determine the effects of the technology on health outcomes.

For decompression with coflex vs decompression with spinal fusion, the pivotal randomized controlled trial, conducted in a patient population with spondylolisthesis no greater than grade 1 and significant back pain, showed that stabilization of decompression with the coflex implant was noninferior to decompression with spinal fusion for the composite clinical success measure. However, there is uncertainty about the net benefit of routinely adding spinal fusion to decompression in patients with no or low-grade spondylolisthesis. Therefore, demonstrating the noninferiority of coflex plus spinal decompression vs spinal decompression plus fusion, a comparator whose benefit on health outcomes is uncertain, makes it difficult to apply the results of the study.

Clinical input supplements and informs the interpretation of the published evidence. Clinical input respondents were mixed in the level of support of this indication. While some of the expert opinion supported a potential benefit in carefully selected individuals, other experts were not confident of a clinically meaningful benefit or use in generally accepted medical practice, citing long-term complications leading to removal of the device. Some clinical input suggested that spacers may have utility in patients who are high risk for general anesthesia. Consideration of existing studies as indirect evidence regarding the outcomes of using spacers in this subgroup is limited by substantial uncertainty regarding the balance of potential benefits and harms. The main source of uncertainty about the benefits versus risks of using coflex plus laminectomy in patients who are not able to have general anesthesia is whether revisions, removals, and other secondary

surgical procedures can be conducted safely if they are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

The following CPT codes are not covered for BlueCHiP for Medicare and not medically necessary for Commercial Products:

22867 Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level

22868 Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (list separately in addition to code for primary procedure)

22869 Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level

22870 Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (list separately in addition to code for primary procedure)

There is no specific CPT code for the removal of interlaminar/interspinous process stabilization/distraction devices, therefore, an appropriate Unlisted CPT code should be used.

RELATED POLICIES

Not applicable

PUBLISHED

Provider Update, April 2019

Provider Update, February 2019

Provider Update, January 2018

Provider Update, January 2017

Provider Update, June 2015

REFERENCES

1. Lurie J, Tomkins-Lane C. Management of lumbar spinal stenosis. *BMJ*. 2016;352:h6234.
2. Lurie JD, Tosteson TD, Tosteson A, et al. Long-term outcomes of lumbar spinal stenosis: eight-year results of the Spine Patient Outcomes Research Trial (SPORT). *Spine (Phila Pa 1976)*. 2015;40(2):63-76.
3. Schroeder GD, Kurd MF, Vaccaro AR. Lumbar spinal stenosis: How is it classified? *The Journal of the American Academy of Orthopaedic Surgeons*. 2016;24(12):843-852.
4. Haig AJ, Tomkins CC. Diagnosis and management of lumbar spinal stenosis. *Jama*. 2010;303(1):71-72.
5. Genevay S, Atlas SJ, Katz JN. Variation in eligibility criteria from studies of radiculopathy due to a herniated disc and of neurogenic claudication due to lumbar spinal stenosis: a structured literature review. *Spine (Phila Pa 1976)*. 2010;35(7):803-811.
6. Chou R, Deyo R, Friedly J, et al. Nonpharmacologic therapies for low back pain: a systematic review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med*. 2017;166(7):493-505.
7. Birkmeyer NJ, Weinstein JN, Tosteson AN, et al. Design of the Spine Patient Outcomes Research Trial (SPORT). *Spine (Phila Pa 1976)*. 2002;27(12):1361-1372.
8. Fritz JM, Lurie JD, Zhao W, et al. Associations between physical therapy and long-term outcomes for individuals with lumbar spinal stenosis in the SPORT study. *The spine journal : official journal of the North American Spine Society*. 2014;14(8):1611-1621.
9. Delitto A, Piva SR, Moore CG, et al. Surgery versus nonsurgical treatment of lumbar spinal stenosis: a randomized trial. *Ann Intern Med*. 2015;162(7):465-473.

10. Katz JN. Surgery for lumbar spinal stenosis: informed patient preferences should weigh heavily. *Ann Intern Med.* 2015;162(7):518-519.
11. Pearson A, Blood E, Lurie J, et al. Predominant leg pain is associated with better surgical outcomes in degenerative spondylolisthesis and spinal stenosis: results from the Spine Patient Outcomes Research Trial (SPORT). *Spine (Phila Pa 1976).* 2011;36(3):219-229.
12. Pearson A, Blood E, Lurie J, et al. Degenerative spondylolisthesis versus spinal stenosis: does a slip matter? Comparison of baseline characteristics and outcomes (SPORT). *Spine (Phila Pa 1976).* 2010;35(3):298-305.
13. Abdu WA, Lurie JD, Spratt KF, et al. Degenerative spondylolisthesis: does fusion method influence outcome? Four-year results of the spine patient outcomes research trial. *Spine (Phila Pa 1976).* 2009;34(21):2351-2360.
14. Deyo RA, Mirza SK, Martin BI, Kreuter W, Goodman DC, Jarvik JG. Trends, major medical complications, and charges associated with surgery for lumbar spinal stenosis in older adults. *Jama.* 2010;303(13):1259-1265.
15. Dartmouth Institute. Variation in the care of surgical conditions: spinal stenosis. 2014.
16. Yoshihara H, Yoneoka D. National trends in the surgical treatment for lumbar degenerative disc disease: United States, 2000 to 2009. *The spine journal : official journal of the North American Spine Society.* 2015;15(2):265-271.
17. Forsth P, Olafsson G, Carlsson T, et al. A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *The New England journal of medicine.* 2016;374(15):1413-1423.
18. Ghogawala Z, Dziura J, Butler WE, et al. Laminectomy plus fusion versus laminectomy alone for lumbar spondylolisthesis. *The New England journal of medicine.* 2016;374(15):1424-1434.
19. Peul WC, Moojen WA. Fusion surgery for lumbar spinal stenosis [letter]. *The New England journal of medicine.* 2016;375(6):601.
20. El Teclé NE, Dahdaleh NS. Fusion surgery for lumbar spinal stenosis [letter]. *The New England journal of medicine.* 2016;375(6):597.
21. Forsth P, Michaelsson K, Sanden B. Fusion surgery for lumbar spinal stenosis [letter]. *The New England journal of medicine.* 2016;375(6):599-600.
22. Su BW, Vaccaro AR. Fusion surgery for lumbar spinal stenosis [letter]. *The New England journal of medicine.* 2016;375(6):597-598.
23. Vasudeva VS, Chi JH. Fusion surgery for lumbar spinal stenosis [letter]. *The New England journal of medicine.* 2016;375(6):598.
24. Dijkerman ML, Overvest GM, Moojen WA, Vleggeert-Lankamp CLA. Decompression with or without concomitant fusion in lumbar stenosis due to degenerative spondylolisthesis: a systematic review. *Eur Spine J.* 2018;27(7):1629-1643.
25. Pearson AM. Fusion in degenerative spondylolisthesis: how to reconcile conflicting evidence. *J Spine Surg.* 2016;2(2):143-145.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

