Medical Coverage Policy | Low-Level Laser Therapy



EFFECTIVE DATE: 11 | 01 | 2022

POLICY LAST UPDATED: 07 | 20 | 2022

OVERVIEW

Low-level laser therapy (LLLT), also called photobiomodulation, is being evaluated to treat various conditions including oral mucositis, myofascial pain, joint pain, lymphedema, and chronic wounds.

MEDICAL CRITERIA

Not applicable.

PRIOR AUTHORIZATION

Not applicable.

POLICY STATEMENT

Medicare Advantage Plans

Low-level laser therapy is covered for prevention of oral mucositis in patients undergoing cancer treatment associated with increased risk of oral mucositis, including chemotherapy and/or radiotherapy, and/or hematopoietic stem cell transplantation.

Low-level laser therapy is not covered for all other indications, as the evidence is insufficient to determine the effects of the technology on health outcomes, including but not limited to:

- Carpal tunnel syndrome
- Neck pain
- Subacromial impingement
- Adhesive capsulitis
- Temporomandibular joint pain
- Low back pain
- Osteoarthritic knee pain
- Heel pain (ie, Achilles tendinopathy, plantar fasciitis)
- Rheumatoid arthritis
- Bell palsy
- Fibromyalgia
- Wound healing
- Lymphedema

Commercial Products

Low-level laser therapy is covered for prevention of oral mucositis in patients undergoing cancer treatment associated with increased risk of oral mucositis, including chemotherapy and/or radiotherapy, and/or hematopoietic stem cell transplantation.

Low-level laser therapy is not medically necessary for all other indications, as the evidence is insufficient to determine the effects of the technology on health outcomes, including but not limited to:

- Carpal tunnel syndrome
- Neck pain
- Subacromial impingement

- Adhesive capsulitis
- Temporomandibular joint pain
- Low back pain
- Osteoarthritic knee pain
- Heel pain (ie, Achilles tendinopathy, plantar fasciitis)
- Rheumatoid arthritis
- Bell palsy
- Fibromyalgia
- Wound healing
- Lymphedema

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Low-level laser therapy (LLLT) refers to the use of red-beam or near-infrared lasers with a wavelength between 600 and 1000 nm and power between 5 and 500 MW. In contrast, lasers used in surgery typically use 300 Watts. When applied to the skin, LLLT produces no sensation and does not burn the skin. Because of the low absorption by human skin, it is hypothesized that the laser light can penetrate deeply into the tissues where it has a photobiostimulative effect. The exact mechanism of its effect on tissue healing is unknown; hypotheses have included improved cellular repair and stimulation of the immune, lymphatic, and vascular systems. LLLT is being evaluated to treat a wide variety of conditions, including soft tissue injuries, myofascial pain, tendinopathies, nerve injuries, and joint pain. LLLT has also been evaluated for lymphedema.

ORAL MUCOSITIS

Oral mucositis describes inflammation of the oral mucosa and typically manifests as erythema or ulcerations that appear 7 to 10 days after initiation of high-dose cancer therapy. Oral mucositis can cause significant pain and increased risk of systemic infection, dependency on total parenteral nutrition, and use of narcotic analgesics.

Treatment

Treatment planning may also need to be modified due to dose-limiting toxicity. There are a number of interventions for oral mucositis that may partially control symptoms, but none is considered a criterion standard treatment. When uncomplicated by infection, oral mucositis is self-limited and usually heals within 2 to 4 weeks after cessation of cytotoxic chemotherapy. Low-level laser therapy (LLLT) has been used in cancer therapy—induced oral mucositis in patients treated with radiotherapy and/or chemotherapy and hematopoietic cell transplantation.

MUSCULOSKELETAL AND NEUROLOGIC DISORDERS

Carpal tunnel syndrome (CTS) is the most common entrapment neuropathy and the most commonly performed surgery of the hand. The syndrome is related to the bony anatomy of the wrist. The carpal tunnel is bound dorsally and laterally by the carpal bones and ventrally by the transverse carpal ligament. Through this contained space run the 9 flexor tendons and the median nerve. Therefore, any space-occupying lesion can compress the median nerve and produce the typical symptoms of CTS - pain, numbness, and tingling in the distribution of the median nerve. Symptoms of more severe cases include hypesthesia, clumsiness, loss of dexterity, and weakness of pinch. In the most severe cases, patients experience marked sensory loss and significant functional impairment with thenar atrophy.

Treatment

Mild-to-moderate cases of CTS are usually first treated conservatively with splinting and cessation of aggravating activities. Other conservative therapies include oral steroids, diuretics, nonsteroidal anti-inflammatory drugs, and steroid injections into the carpal tunnel itself. Patients who do not respond to

conservative therapy or who present with severe CTS with thenar atrophy may be considered candidates for surgical release of the carpal ligament, using either an open or endoscopic approach. LLLT is also used to treat CTS.

For individuals who have increased risk of oral mucositis due to some cancer treatments (eg, chemotherapy, radiotherapy) and/or hematopoietic cell transplantation who receive LLLT, the evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

The evidence for LLLT is insufficient to determine the effects of the technology on health outcomes in individuals who have the following conditions:

- orthopedic pain (ie, neck pain, osteoarthritic knee pain, low back pain, carpal tunnel syndrome)
- shoulder conditions (eg, subacromial impingement syndrome, adhesive capsulitis), heel pain, or temporomandibular joint pain
- bone, ligament, and joint conditions (eg, rheumatoid arthritis, fibromyalgia)
- Bell palsy
- lymphedema
- chronic non-healing wounds

CODING

Medicare Advantage Plans and Commercial Products

The following code(s) are covered when filed with the ICD-10 CM Diagnosis* codes below:

0552T Low-level laser therapy, dynamic photonic and dynamic thermokinetic energies, provided by a physician or other qualified health care professional

S8948 Application of a modality (requiring constant provider attendance) to one or more areas; low-level laser; each 15 minutes

*ICD-10 CM Diagnosis:

C00 - D49 K12.30-K12.39

RELATED POLICIES

Not applicable

PUBLISHED

Provider Update, September 2022 Provider Update, October 2021 Provider Update, December 2020 Provider Update, January 2020 Provider Update, September 2018

REFERENCES

- 1. Lalla RV, Bowen J, Barasch A, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. Cancer. May 15 2014; 120(10): 1453-61. PMID 24615748
- 2. Schubert MM, Eduardo FP, Guthrie KA, et al. A phase III randomized double-blind placebo-controlled clinical trial to determine the efficacy of low level laser therapy for the prevention of oral mucositis in patients undergoing hematopoietic cell transplantation. Support Care Cancer. Oct 2007; 15(10): 1145-54. PMID 17393191
- 3. Figueiredo AL, Lins L, Cattony AC, et al. Laser therapy in the control of oral mucositis: a meta-analysis. Rev Assoc Med Bras (1992). Sep-Oct 2013; 59(5): 467-74. PMID 24119379
- 4. Doeuk C, Hersant B, Bosc R, et al. Current indications for low level laser treatment in maxillofacial surgery: a review. Br J Oral Maxillofac Surg. Apr 2015; 53(4): 309-15. PMID 25740083
- 5. Peng J, Shi Y, Wang J, et al. Low-level laser therapy in the prevention and treatment of oral mucositis: a systematic review and meta-analysis. Oral Surg Oral Med Oral Pathol Oral Radiol. Oct 2020; 130(4): 387-397.e9. PMID 32624448

- 6. Oberoi S, Zamperlini-Netto G, Beyene J, et al. Effect of prophylactic low level laser therapy on oral mucositis: a systematic review and meta-analysis. PLoS One. 2014; 9(9): e107418. PMID 25198431
- 7. Rankin IA, Sargeant H, Rehman H, et al. Low-level laser therapy for carpal tunnel syndrome. Cochrane. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD012765/full Published 22 August 2017. Accessed April 25, 2022.
- 8. Li ZJ, Wang Y, Zhang HF, et al. Effectiveness of low-level laser on carpal tunnel syndrome: A metaanalysis of previously reported randomized trials. Medicine (Baltimore). Aug 2016; 95(31): e4424. PMID 27495063
- 9. Fusakul Y, Aranyavalai T, Saensri P, et al. Low-level laser therapy with a wrist splint to treat carpal tunnel syndrome: a double-blinded randomized controlled trial. Lasers Med Sci. May 2014; 29(3): 1279-87. PMID 24477392
- 10. Low-level laser therapy for carpal tunnel syndrome and chronic neck pain. Technol Eval Cent Assess Program Exec Summ. Nov 2010; 25(4): 1-2. PMID 21638940
- 11. Chow RT, Heller GZ, Barnsley L. The effect of 300 mW, 830 nm laser on chronic neck pain: a double-blind, randomized, placebo-controlled study. Pain. Sep 2006; 124(1-2): 201-10. PMID 16806710
- 12. Gross AR, Dziengo S, Boers O, et al. Low Level Laser Therapy (LLLT) for Neck Pain: A Systematic Review and Meta-Regression. Open Orthop J. 2013; 7: 396-419. PMID 24155802
- 13. Yeldan I, Cetin E, Ozdincler AR. The effectiveness of low-level laser therapy on shoulder function in subacromial impingement syndrome. Disabil Rehabil. 2009; 31(11): 935-40. PMID 19031167
- 14. Dogan SK, Ay S, Evcik D. The effectiveness of low laser therapy in subacromial impingement syndrome: a randomized placebo controlled double-blind prospective study. Clinics (Sao Paulo). 2010; 65(10): 1019-22. PMID 21120304
- 15. Abrisham SM, Kermani-Alghoraishi M, Ghahramani R, et al. Additive effects of low-level laser therapy with exercise on subacromial syndrome: a randomised, double-blind, controlled trial. Clin Rheumatol. Oct 2011; 30(10): 1341-6. PMID 21538218
- 16. Bal A, Eksioglu E, Gurcay E, et al. Low-level laser therapy in subacromial impingement syndrome. Photomed Laser Surg. Feb 2009; 27(1): 31-6. PMID 19250050
- 17. Calis HT, Berberoglu N, Calis M. Are ultrasound, laser and exercise superior to each other in the treatment of subacromial impingement syndrome? A randomized clinical trial. Eur J Phys Rehabil Med. Mar 2 2011;47(3):375-380. PMID 21364511
- 18. Alfredo PP, Bjordal JM, Junior WS, et al. Efficacy of low-level laser therapy combined with exercise for subacromial impingement syndrome: A randomised controlled trial. Clin Rehabil. Jun 2021; 35(6): 851-860. PMID 33307783
- 19. Badil Guloglu S. Comparison of low-level laser treatment and extracorporeal shock wave therapy in subacromial impingement syndrome: a randomized, prospective clinical study. Lasers Med Sci. Jun 2021; 36(4): 773-781. PMID 32638239
- 20. Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). Cochrane Database Syst Rev. Oct 01 2014; (10): CD011324. PMID 25271097
- 21. Stergioulas A, Stergioula M, Aarskog R, et al. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. Am J Sports Med. May 2008; 36(5): 881-7. PMID 18272794
- 22. Chen J, Huang Z, Ge M, et al. Efficacy of low-level laser therapy in the treatment of TMDs: a meta-analysis of 14 randomised controlled trials. J Oral Rehabil. Apr 2015; 42(4): 291-9. PMID 25491183
- 23. Chang WD, Lee CL, Lin HY, et al. A Meta-analysis of Clinical Effects of Low-level Laser Therapy on Temporomandibular Joint Pain. J Phys Ther Sci. Aug 2014; 26(8): 1297-300. PMID 25202201
- 24. Hanna R, Dalvi S, Bensadoun RJ, et al. Role of Photobiomodulation Therapy in Modulating Oxidative Stress in Temporomandibular Disorders. A Systematic Review and Meta-Analysis of Human Randomised Controlled Trials. Antioxidants (Basel). Jun 25 2021; 10(7). PMID 34202292
- 25. Conti PC. Low level laser therapy in the treatment of temporomandibular disorders (TMD): a double-blind pilot study. Cranio. Apr 1997; 15(2): 144-9. PMID 9586517
- 26. Kulekcioglu S, Sivrioglu K, Ozcan O, et al. Effectiveness of low-level laser therapy in temporomandibular disorder. Scand J Rheumatol. 2003; 32(2): 114-8. PMID 12737331

- 27. Venancio Rde A, Camparis CM, Lizarelli Rde F. Low intensity laser therapy in the treatment of temporomandibular disorders: a double-blind study. J Oral Rehabil. Nov 2005; 32(11): 800-7. PMID 16202043
- 28. Cetiner S, Kahraman SA, Yucetas S. Evaluation of low-level laser therapy in the treatment of temporomandibular disorders. Photomed Laser Surg. Oct 2006; 24(5): 637-41. PMID 17069496
- 29. Fikackova H, Dostalova T, Navratil L, et al. Effectiveness of low-level laser therapy in temporomandibular joint disorders: a placebo-controlled study. Photomed Laser Surg. Aug 2007; 25(4): 297-303. PMID 17803388
- 30. Mazzetto MO, Carrasco TG, Bidinelo EF, et al. Low intensity laser application in temporomandibular disorders: a phase I double-blind study. Cranio. Jul 2007; 25(3): 186-92. PMID 17696035
- 31. Frare J.C., Nicolau R.A. Clinical analysis of the effect of laser photobiomodulation (GaAs904 nm) on temporomandibular joint dysfunction. Rev. Bras. Fisioter. 2008;12:3742. doi: 10.1590/S1413-3555200800100008.
- 32.da Cunha LA, Firoozmand LM, da Silva AP, et al. Efficacy of low-level laser therapy in the treatment of temporomandibular disorder. Int Dent J. Aug 2008; 58(4): 213-7. PMID 18783114
- 33. Lassemi E., Jafari S.M., Motamedi M.H.K., Navi F., Lasemi R. Low- level laser therapy in the management of temporamandibular joint disorder. J. Oral Laser Appl. 2008;8:8386
- 34. Carrasco TG, Mazzetto MO, Mazzetto RG, et al. Low intensity laser therapy in temporomandibular disorder: a phase II double-blind study. Cranio. Oct 2008; 26(4): 274-81. PMID 19004308
- 35. Emshoff R, Bosch R, Pumpel E, et al. Low-level laser therapy for treatment of temporomandibular joint pain: a double-blind and placebo-controlled trial. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. Apr 2008; 105(4): 452-6. PMID 18329580
- 36. Carrasco TG, Guerisoli LD, Guerisoli DM, et al. Evaluation of low intensity laser therapy in myofascial pain syndrome. Cranio. Oct 2009; 27(4): 243-7. PMID 19891258
- 37. Shirani AM, Gutknecht N, Taghizadeh M, et al. Low-level laser therapy and myofacial pain dysfunction syndrome: a randomized controlled clinical trial. Lasers Med Sci. Sep 2009; 24(5): 715-20. PMID 19002646
- 38. Venezian GC, da Silva MA, Mazzetto RG, et al. Low level laser effects on pain to palpation and electromyographic activity in TMD patients: a double-blind, randomized, placebo-controlled study. Cranio. Apr 2010; 28(2): 84-91. PMID 20491229
- 39. Oz S, Gokcen-Rohlig B, Saruhanoglu A, et al. Management of myofascial pain: low-level laser therapy versus occlusal splints. J Craniofac Surg. Nov 2010; 21(6): 1722-8. PMID 21119408
- 40. Marini I, Gatto MR, Bonetti GA. Effects of superpulsed low-level laser therapy on temporomandibular joint pain. Clin J Pain. Sep 2010; 26(7): 611-6. PMID 20664343
- 41. Rohlig B.G., Kipirdi S., Meric U., Capan N., Keskin H. Masticatory muscle pain and low-level laser therapy: A double-blind and placebo-controlled study. Turk. J. Phys. Med. Rehabil. Turk. Fiz. Tip Rehabil. Derg. 2011;57:3137. doi: 10.4274/tftr.57.06.
- 42. Sattayut S, Bradley P. A study of the influence of low intensity laser therapy on painful temporomandibular disorder patients. Laser Ther. Sep 30 2012; 21(3): 183-92. PMID 24511188
- 43.de Carli ML, Guerra MB, Nunes TB, et al. Piroxicam and laser phototherapy in the treatment of TMJ arthralgia: a double-blind randomised controlled trial. J Oral Rehabil. Mar 2013; 40(3): 171-8. PMID 23252583
- 44.da Silva MA, Botelho AL, Turim CV, et al. Low level laser therapy as an adjunctive technique in the management of temporomandibular disorders. Cranio. Oct 2012; 30(4): 264-71. PMID 23156967
- 45. Panhoca VH, Lizarelli Rde F, Nunez SC, et al. Comparative clinical study of light analgesic effect on temporomandibular disorder (TMD) using red and infrared led therapy. Lasers Med Sci. Feb 2015; 30(2): 815-22. PMID 24197518
- 46. Uemoto L, Garcia MA, Gouvea CV, et al. Laser therapy and needling in myofascial trigger point deactivation. J Oral Sci. 2013; 55(2): 175-81. PMID 23748458
- 47. Ferreira LA, de Oliveira RG, Guimaraes JP, et al. Laser acupuncture in patients with temporomandibular dysfunction: a randomized controlled trial. Lasers Med Sci. Nov 2013; 28(6): 1549-58. PMID 23380907
- 48. Demirkol N, Sari F, Bulbul M, et al. Effectiveness of occlusal splints and low-level laser therapy on myofascial pain. Lasers Med Sci. Apr 2015; 30(3): 1007-12. PMID 24504660
- 49. Ahrari F, Madani AS, Ghafouri ZS, et al. The efficacy of low-level laser therapy for the treatment of myogenous temporomandibular joint disorder. Lasers Med Sci. Mar 2014; 29(2): 551-7. PMID 23318917

- 50. Pereira TS, Flecha OD, Guimaraes RC, et al. Efficacy of red and infrared lasers in treatment of temporomandibular disorders--a double-blind, randomized, parallel clinical trial. Cranio. Jan 2014; 32(1): 51-6. PMID 24660647
- 51.de Moraes Maia ML, Ribeiro MA, Maia LG, et al. Evaluation of low-level laser therapy effectiveness on the pain and masticatory performance of patients with myofascial pain. Lasers Med Sci. Jan 2014; 29(1): 29-35. PMID 23143142
- 52. Sancakli E, Gokcen-Rohlig B, Balik A, et al. Early results of low-level laser application for masticatory muscle pain: a double-blind randomized clinical study. BMC Oral Health. Oct 23 2015; 15(1): 131. PMID 26496720
- 53. Douglas De Oliveira DW, Lages FS, Guimaraes RC, et al. Do TMJ symptoms improve and last across time after treatment with red (660 nm) and infrared (790 nm) low level laser treatment (LLLT)? A survival analysis. Cranio. Nov 2017; 35(6): 372-378. PMID 28218006
- 54. Costa SAP, Florezi GP, Artes GE, et al. The analgesic effect of photobiomodulation therapy (830 nm) on the masticatory muscles: a randomized, double-blind study. Braz Oral Res. Dec 18 2017; 31: e107. PMID 29267668
- 55. Seifi M, Ebadifar A, Kabiri S, et al. Comparative effectiveness of Low Level Laser therapy and Transcutaneous Electric Nerve Stimulation on Temporomandibular Joint Disorders. J Lasers Med Sci. 2017; 8(Suppl 1): S27-S31. PMID 29071032
- 56. Shobha R, Narayanan VS, Jagadish Pai BS, et al. Low-level laser therapy: A novel therapeutic approach to temporomandibular disorder A randomized, double-blinded, placebo-controlled trial. Indian J Dent Res. Jul-Aug 2017; 28(4): 380-387. PMID 28836528
- 57. Rezazadeh F, Hajian K, Shahidi S, et al. Comparison of the Effects of Transcutaneous Electrical Nerve Stimulation and Low-Level Laser Therapy on Drug-Resistant Temporomandibular Disorders. J Dent (Shiraz). Sep 2017; 18(3): 187-192. PMID 29034273
- 58. Varma S.R., al Shayeb M., el Kaseh A., Kuduruthullah S., Ashekhi A., al Khader E. Effectiveness of low-level laser therapy in the Management of the Temporomandibular Joint Disorders: A Placebo-controlled Trial. World J. Dent. 2018;9:316320. doi: 10.5005/jp-journals-10015-1555.
- 59. Borges RMM, Cardoso DS, Flores BC, et al. Effects of different photobiomodulation dosimetries on temporomandibular dysfunction: a randomized, double-blind, placebo-controlled clinical trial. Lasers Med Sci. Dec 2018; 33(9): 1859-1866. PMID 29850961
- 60. Brochado FT, Jesus LH, Carrard VC, et al. Comparative effectiveness of photobiomodulation and manual therapy alone or combined in TMD patients: a randomized clinical trial. Braz Oral Res. Jul 10 2018; 32: e50. PMID 29995062
- 61. Rodrigues CA, Melchior MO, Valencise Magri L, et al. Can the severity of orofacial myofunctional conditions interfere with the response of analgesia promoted by active or placebo low-level laser therapy?. Cranio. Jul 2020; 38(4): 240-247. PMID 30244669
- 62. Peimani A., Keshavarz S., Fathollahi M.S. Comparison of Low-Level Laser Therapy and Drug Therapy in Patients with Temporomandibular Disorders: A Randomized Clinical Trial. J. Oral Health Dent. 2020;38:240247. doi: 10.1080/08869634.2018.1520950.
- 63. Nadershah M, Abdel-Alim HM, Bayoumi AM, et al. Photobiomodulation Therapy for Myofascial Pain in Temporomandibular Joint Dysfunction: A Double-Blinded Randomized Clinical Trial. J Maxillofac Oral Surg. Mar 2020; 19(1): 93-97. PMID 31988570
- 64. Magri LV, Bataglion C, Leite-Panissi CRA. Follow-up results of a randomized clinical trial for low-level laser therapy in painful TMD of muscular origins. Cranio. Nov 2021; 39(6): 502-509. PMID 31585522
- 65. Al-Quisi AF, Al-Anee AM, Al-Jumaily HA, et al. Efficacy of the LED Red Light Therapy in the Treatment of Temporomandibular Disorders: Double Blind Randomized Controlled Trial. Pain Res Treat. 2019; 2019: 8578703. PMID 31205787
- 66. Herpich CM, Leal-Junior ECP, Politti F, et al. Intraoral photobiomodulation diminishes pain and improves functioning in women with temporomandibular disorder: a randomized, sham-controlled, double-blind clinical trial: Intraoral photobiomodulation diminishes pain in women with temporomandibular disorder. Lasers Med Sci. Mar 2020; 35(2): 439-445. PMID 31325122
- 67. Khairnar S, Bhate K, S N SK, et al. Comparative evaluation of low-level laser therapy and ultrasound heat therapy in reducing temporomandibular joint disorder pain. J Dent Anesth Pain Med. Oct 2019; 19(5): 289-294. PMID 31723669

- 68. Sobral APT, Godoy CLH, Fernandes KPS, et al. Photomodulation in the treatment of chronic pain in patients with temporomandibular disorder: protocol for cost-effectiveness analysis. BMJ Open. May 05 2018; 8(5): e018326. PMID 29730613
- 69. Maracci LM, Stasiak G, de Oliveira Chami V, et al. Treatment of myofascial pain with a rapid laser therapy protocol compared to occlusal splint: A double-blind, randomized clinical trial. Cranio. Jun 03 2020: 1-7. PMID 32491964
- 70. Chellappa D, Thirupathy M. Comparative efficacy of low-Level laser and TENS in the symptomatic relief of temporomandibular joint disorders: A randomized clinical trial. Indian J Dent Res. Jan-Feb 2020; 31(1): 42-47. PMID 32246680
- 71. Monteiro L, Ferreira R, Resende T, et al. Effectiveness of Photobiomodulation in Temporomandibular Disorder-Related Pain Using a 635 nm Diode Laser: A Randomized, Blinded, and Placebo-Controlled Clinical Trial. Photobiomodul Photomed Laser Surg. May 2020; 38(5): 280-288. PMID 32427553
- 72. Del Vecchio A, Floravanti M, Boccassini A, et al. Evaluation of the efficacy of a new low-level laser therapy home protocol in the treatment of temporomandibular joint disorder-related pain: A randomized, double-blind, placebo-controlled clinical trial. Cranio. Mar 2021; 39(2): 141-150. PMID 30999823
- 73. Aisaiti A, Zhou Y, Wen Y, et al. Effect of photobiomodulation therapy on painful temporomandibular disorders. Sci Rep. Apr 27 2021; 11(1): 9049. PMID 33907210
- 74. Glazov G, Yelland M, Emery J. Low-level laser therapy for chronic non-specific low back pain: a meta-analysis of randomised controlled trials. Acupunct Med. Oct 2016; 34(5): 328-341. PMID 27207675
- 75. Huang Z, Ma J, Chen J, et al. The effectiveness of low-level laser therapy for nonspecific chronic low back pain: a systematic review and meta-analysis. Arthritis Res Ther. Dec 15 2015; 17: 360. PMID 26667480
- 76. Chen YJ, Liao CD, Hong JP, et al. Effects of laser therapy on chronic low back pain: A systematic review and meta-analysis of randomized controlled trials. Clin Rehabil. Mar 2022; 36(3): 289-302. PMID 34757882
- 77. Alayat MS, Atya AM, Ali MM, et al. Long-term effect of high-intensity laser therapy in the treatment of patients with chronic low back pain: a randomized blinded placebo-controlled trial. Lasers Med Sci. May 2014; 29(3): 1065-73. PMID 24178907
- 78. Ay S, Dogan SK, Evcik D. Is low-level laser therapy effective in acute or chronic low back pain? Clin Rheumatol. Aug 2010; 29(8): 905-10. PMID 20414695
- 79. Basford JR, Sheffield CG, Harmsen WS. Laser therapy: a randomized, controlled trial of the effects of low-intensity Nd:YAG laser irradiation on musculoskeletal back pain. Arch Phys Med Rehabil. Jun 1999; 80(6): 647-52. PMID 10378490
- 80. Djavid GE, Mehrdad R, Ghasemi M, et al. In chronic low back pain, low level laser therapy combined with exercise is more beneficial than exercise alone in the long term: a randomised trial. Aust J Physiother. 2007; 53(3): 155-60. PMID 17725472
- 81. Glazov G, Schattner P, Lopez D, et al. Laser acupuncture for chronic non-specific low back pain: a controlled clinical trial. Acupunct Med. Sep 2009; 27(3): 94-100. PMID 19734378
- 82. Glazov G, Yelland M, Emery J. Low-dose laser acupuncture for non-specific chronic low back pain: a double-blind randomised controlled trial. Acupunct Med. Apr 2014; 32(2): 116-23. PMID 24280948
- 83. Klein RG, Eek BC. Low-energy laser treatment and exercise for chronic low back pain: double-blind controlled trial. Arch Phys Med Rehabil. Jan 1990; 71(1): 34-7. PMID 2136991
- 84. Konstantinovic L, Lazovic M, Milovanovic N, et al.. Low level laser therapy in geriatric patients with low back pain. Eur J Pain Suppl (Poster Sessions) 2011;5:61 10.1016/S1754-3207(11)70205-X
- 85.Lin ML, Wu HC, Hsieh YH, et al. Evaluation of the effect of laser acupuncture and cupping with ryodoraku and visual analog scale on low back pain. Evid Based Complement Alternat Med. 2012; 2012: 521612. PMID 23118792
- 86.Okamoto H. Therapeutic effect of semiconductor laser irradiation on low-back pain. J Jpn A Phys Med Balneology Climatology 1989;52:13145.
- 87. Ruth M, Weber M, Zenz M. [Laser acupuncture for chronic back pain. A double-blind clinical study]. Schmerz. Sep 2010; 24(5): 485-93. PMID 20872127
- 88. Soriano F, Rios R. Gallium arsenide laser treatment of chronic low back pain: a prospective, randomized and double blind study. Laser Ther 1998;10:17580. 10.5978/islsm.10.175

- 89. Umegaki S. Effectiveness of low-power laser therapy on low-back pain: double blind comparative study to evaluate the analgesic effect of low-power laser therapy. Kiso to Rinsho (The Clinical Report) 1989;23:283946.
- 90. Vallone F, Benedicenti S, Sorrenti E, et al. Effect of diode laser in the treatment of patients with nonspecific chronic low back pain: a randomized controlled trial. Photomed Laser Surg. Sep 2014; 32(9): 490-4. PMID 25141218
- 91. Wallace G. The effects of laser acupuncture on chronic low back pain [Thesis]. Melbourne Monash University, 1996.
- 92. Gur A, Karakoc M, Cevik R, et al. Efficacy of low power laser therapy and exercise on pain and functions in chronic low back pain. Lasers Surg Med. 2003; 32(3): 233-8. PMID 12605431
- 93. Hsieh RL, Lee WC. Short-term therapeutic effects of 890-nanometer light therapy for chronic low back pain: a double-blind randomized placebo-controlled study. Lasers Med Sci. Mar 2014; 29(2): 671-9. PMID 23820974
- 94.de Carvalho ME, de Carvalho RM, Marques AP, et al. Low intensity laser and LED therapies associated with lateral decubitus position and flexion exercises of the lower limbs in patients with lumbar disk herniation: clinical randomized trial. Lasers Med Sci. Sep 2016; 31(7): 1455-63. PMID 27379776
- 95. Tantawy SA, Abdelbasset WK, Kamel DM, et al. Laser photobiomodulation is more effective than ultrasound therapy in patients with chronic nonspecific low back pain: a comparative study. Lasers Med Sci. Jun 2019; 34(4): 793-800. PMID 30334124
- 96. Nambi G, Kamal W, Es S, et al. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: a randomized controlled study. Eur J Phys Rehabil Med. Dec 2018; 54(6): 880-889. PMID 29687966
- 97. Shin JY, Ku B, Kim JU, et al. Short-Term Effect of Laser Acupuncture on Lower Back Pain: A Randomized, Placebo-Controlled, Double-Blind Trial. Evid Based Complement Alternat Med. 2015; 2015: 808425. PMID 26516333
- 98. Koldas Dogan S, Ay S, Evcik D. The effects of two different low level laser therapies in the treatment of patients with chronic low back pain: A double-blinded randomized clinical trial. J Back Musculoskelet Rehabil. 2017; 30(2): 235-240. PMID 27472858
- 99. Huang Z, Chen J, Ma J, et al. Effectiveness of low-level laser therapy in patients with knee osteoarthritis: a systematic review and meta-analysis. Osteoarthritis Cartilage. Sep 2015; 23(9): 1437-1444. PMID 25914044
- 100.Bjordal JM, Johnson MI, Lopes-Martins RA, et al. Short-term efficacy of physical interventions in osteoarthritic knee pain. A systematic review and meta-analysis of randomised placebo-controlled trials. BMC Musculoskelet Disord. Jun 22 2007; 8: 51. PMID 17587446
- 101. Stausholm MB, Naterstad IF, Joensen J, et al. Efficacy of low-level laser therapy on pain and disability in knee osteoarthritis: systematic review and meta-analysis of randomised placebo-controlled trials. BMJ Open. Oct 28 2019; 9(10): e031142. PMID 31662383
- 102.Al Rashoud AS, Abboud RJ, Wang W, et al. Efficacy of low-level laser therapy applied at acupuncture points in knee osteoarthritis: a randomised double-blind comparative trial. Physiotherapy. Sep 2014; 100(3): 242-8. PMID 24418801
- 103.Alfredo PP, Bjordal JM, Dreyer SH, et al. Efficacy of low level laser therapy associated with exercises in knee osteoarthritis: a randomized double-blind study. Clin Rehabil. Jun 2012; 26(6): 523-33. PMID 22169831
- 104. Alfredo PP, Bjordal JM, Junior WS, et al. Long-term results of a randomized, controlled, double-blind study of low-level laser therapy before exercises in knee osteoarthritis: laser and exercises in knee osteoarthritis. Clin Rehabil. Feb 2018; 32(2): 173-178. PMID 28776408
- 105.Alghadir A, Omar MT, Al-Askar AB, et al. Effect of low-level laser therapy in patients with chronic knee osteoarthritis: a single-blinded randomized clinical study. Lasers Med Sci. Mar 2014; 29(2): 749-55. PMID 23912778
- 106.Bagheri SR, Fatemi E, Fazeli SH, et al.. Efficacy of low level laser on knee osteoarthritis treatment [Persian]. Koomesh 2011;12:28592.
- 107.Bulow PM, Jensen H, Danneskiold-Samsoe B. Low power Ga-Al-As laser treatment of painful osteoarthritis of the knee. A double-blind placebo-controlled study. Scand J Rehabil Med. Sep 1994; 26(3): 155-9. PMID 7801065

- 108.Delkhosh CT, Fatemy E, Ghorbani R, et al. Comparing the immediate and long-term effects of low and high power laser on the symptoms of knee osteoarthritis [Persian]. Journal of mazandaran university of medical sciences 2018;28:6977.
- 109.Fukuda VO, Fukuda TY, Guimaraes M, et al. SHORT-TERM EFFICACY OF LOW-LEVEL LASER THERAPY IN PATIENTS WITH KNEE OSTEOARTHRITIS: A RANDOMIZED PLACEBO-CONTROLLED, DOUBLE-BLIND CLINICAL TRIAL. Rev Bras Ortop. Sep-Oct 2011; 46(5): 526-33. PMID 27027049
- 110.Gur A, Cosut A, Sarac AJ, et al. Efficacy of different therapy regimes of low-power laser in painful osteoarthritis of the knee: a double-blind and randomized-controlled trial. Lasers Surg Med. 2003; 33(5): 330-8. PMID 14677160
- 111.Gworys K, Gasztych J, Puzder A, et al. Influence of various laser therapy methods on knee joint pain and function in patients with knee osteoarthritis. Ortop Traumatol Rehabil. May-Jun 2012; 14(3): 269-77. PMID 22764339
- 112.Hegedus B, Viharos L, Gervain M, et al. The effect of low-level laser in knee osteoarthritis: a double-blind, randomized, placebo-controlled trial. Photomed Laser Surg. Aug 2009; 27(4): 577-84. PMID 19530911
- 113.Helianthi DR, Simadibrata C, Srilestari A, et al. Pain Reduction After Laser Acupuncture Treatment in Geriatric Patients with Knee Osteoarthritis: a Randomized Controlled Trial. Acta Med Indones. Apr 2016; 48(2): 114-21. PMID 27550880
- 114.Hinman RS, McCrory P, Pirotta M, et al. Acupuncture for chronic knee pain: a randomized clinical trial. JAMA. Oct 01 2014; 312(13): 1313-22. PMID 25268438
- 115.Jensen H, Harreby M, Kjer J. [Infrared laser--effect in painful arthrosis of the knee?]. Ugeskr Laeger. Nov 09 1987; 149(46): 3104-6. PMID 3445368
- 116.Kheshie AR, Alayat MS, Ali MM. High-intensity versus low-level laser therapy in the treatment of patients with knee osteoarthritis: a randomized controlled trial. Lasers Med Sci. Jul 2014; 29(4): 1371-6. PMID 24487957
- 117. Koutenaei FR, Mosallanezhad Z, Naghikhani M, et al.. The effect of low level laser therapy on pain and range of motion of patients with knee osteoarthritis. Physical Treatments Specific Physical Therapy 2017;7:1318.
- 118.Mohammed N, Allam H, Elghoroury E, et al. Evaluation of serum beta-endorphin and substance P in knee osteoarthritis patients treated by laser acupuncture. J Complement Integr Med. Jan 05 2018; 15(2). PMID 29303777
- 119.S GN, Kamal W, George J, et al. Radiological and biochemical effects (CTX-II, MMP-3, 8, and 13) of low-level laser therapy (LLLT) in chronic osteoarthritis in Al-Kharj, Saudi Arabia. Lasers Med Sci. Feb 2017; 32(2): 297-303. PMID 27913970
- 120.Nivbrant B, Friberg S. [Laser treatment of knee joint arthrosis seems to be effective but scientific evidence is lacking]. Lakartidningen. Mar 11 1992; 89(11): 859-61. PMID 1545640
- 121.Rayegani SM, Bahrami MH, Elyaspour D, et al.. Therapeutic effects of low level laser therapy (LLLT) in knee osteoarthritis, compared to therapeutic ultrasound. J Lasers Med Sci 2012;3:7174.
- 122. Tascioglu F, Armagan O, Tabak Y, et al. Low power laser treatment in patients with knee osteoarthritis. Swiss Med Wkly. May 01 2004; 134(17-18): 254-8. PMID 15243853
- 123. Youssef EF, Muaidi QI, Shanb AA. Effect of Laser Therapy on Chronic Osteoarthritis of the Knee in Older Subjects. J Lasers Med Sci. 2016; 7(2): 112-9. PMID 27330707
- 124. Tumilty S, McDonough S, Hurley DA, et al. Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of Achilles' tendinopathy: a randomized controlled trial. Arch Phys Med Rehabil. May 2012; 93(5): 733-9. PMID 22541305
- 125. Wang W, Jiang W, Tang C, et al. Clinical efficacy of low-level laser therapy in plantar fasciitis: A systematic review and meta-analysis. Medicine (Baltimore). Jan 2019; 98(3): e14088. PMID 30653125
- 126. Guimaraes JS, Arcanjo FL, Leporace G, et al. Effect of low-level laser therapy on pain and disability in patients with plantar fasciitis: A systematic review and meta-analysis. Musculoskelet Sci Pract. Feb 2022; 57: 102478. PMID 34847470
- 127.Macias DM, Coughlin MJ, Zang K, et al. Low-Level Laser Therapy at 635 nm for Treatment of Chronic Plantar Fasciitis: A Placebo-Controlled, Randomized Study. J Foot Ankle Surg. Sep-Oct 2015; 54(5): 768-72. PMID 25769363

- 128.Kiritsi O, Tsitas K, Malliaropoulos N, et al. Ultrasonographic evaluation of plantar fasciitis after low-level laser therapy: results of a double-blind, randomized, placebo-controlled trial. Lasers Med Sci. Mar 2010; 25(2): 275-81. PMID 19841862
- 129. Cinar E, Saxena S, Uygur F. Low-level laser therapy in the management of plantar fasciitis: a randomized controlled trial. Lasers Med Sci. Jul 2018; 33(5): 949-958. PMID 29273892
- 130.Brosseau L, Robinson V, Wells G, et al. Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis. Cochrane Database Syst Rev. Oct 19 2005; (4): CD002049. PMID 16235295
- 131.Meireles SM, Jones A, Jennings F, et al. Assessment of the effectiveness of low-level laser therapy on the hands of patients with rheumatoid arthritis: a randomized double-blind controlled trial. Clin Rheumatol. May 2010; 29(5): 501-9. PMID 20082104
- 132. Alayat MS, Elsodany AM, El Fiky AA. Efficacy of high and low level laser therapy in the treatment of Bell's palsy: a randomized double blind placebo-controlled trial. Lasers Med Sci. Jan 2014; 29(1): 335-42. PMID 23709010
- 133.Ordahan B, Karahan AY. Role of low-level laser therapy added to facial expression exercises in patients with idiopathic facial (Bell's) palsy. Lasers Med Sci. May 2017; 32(4): 931-936. PMID 28337563
- 134.Honda Y, Sakamoto J, Hamaue Y, et al. Effects of Physical-Agent Pain Relief Modalities for Fibromyalgia Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. Pain Res Manag. 2018; 2018: 2930632. PMID 30402199
- 135.Ruaro JA, Frez AR, Ruaro MB, et al. Low-level laser therapy to treat fibromyalgia. Lasers Med Sci. Nov 2014; 29(6): 1815-9. PMID 24801056
- 136.Matsutani LA, Marques AP, Ferreira EA, et al. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. Clin Exp Rheumatol. May-Jun 2007; 25(3): 410-5. PMID 17631737
- 137.Samson D, Lefevre F, Aronson N. Wound-healing technologies: low-level laser and vacuum-assisted closure. Evid Rep Technol Assess (Summ). Dec 2004; (111): 1-6. PMID 15663354
- 138.Chen C, Hou WH, Chan ES, et al. Phototherapy for treating pressure ulcers. Cochrane Database Syst Rev. Jul 11 2014; (7): CD009224. PMID 25019295
- 139.Machado RS, Viana S, Sbruzzi G. Low-level laser therapy in the treatment of pressure ulcers: systematic review. Lasers Med Sci. May 2017; 32(4): 937-944. PMID 28116536
- 140. Taradaj J, Halski T, Kucharzewski M, et al. Effect of laser irradiation at different wavelengths (940, 808, and 658 nm) on pressure ulcer healing: results from a clinical study. Evid Based Complement Alternat Med 2013;2013:960240. PMID
- 141.Lucas C, van Gemert MJ, de Haan RJ. Efficacy of low-level laser therapy in the management of stage III decubitus ulcers: a prospective, observer-blinded multicentre randomised clinical trial. Lasers Med Sci. 2003; 18(2): 72-7. PMID 12928815
- 142.Nussbaum EL, Biemann I, Mustard B. Comparison of ultrasound/ultraviolet-C and laser for treatment of pressure ulcers in patients with spinal cord injury. Phys Ther. Sep 1994; 74(9): 812-23; discussion 824-5. PMID 8066108
- 143. Taly AB, Sivaraman Nair KP, Murali T, et al. Efficacy of multiwavelength light therapy in the treatment of pressure ulcers in subjects with disorders of the spinal cord: A randomized double-blind controlled trial. Arch Phys Med Rehabil. Oct 2004; 85(10): 1657-61. PMID 15468027
- 144.Li S, Wang C, Wang B, et al. Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. Diabetes Res Clin Pract. Sep 2018; 143: 215-224. PMID 30009935
- 145.Smoot B, Chiavola-Larson L, Lee J, et al. Effect of low-level laser therapy on pain and swelling in women with breast cancer-related lymphedema: a systematic review and meta-analysis. J Cancer Surviv. Jun 2015; 9(2): 287-304. PMID 25432632
- 146.Omar MT, Shaheen AA, Zafar H. A systematic review of the effect of low-level laser therapy in the management of breast cancer-related lymphedema. Support Care Cancer. Nov 2012; 20(11): 2977-84. PMID 22875413
- 147.Sung L, Robinson P, Treister N, et al. Guideline for the prevention of oral and oropharyngeal mucositis in children receiving treatment for cancer or undergoing haematopoietic stem cell transplantation. BMJ Support Palliat Care. Mar 2017; 7(1): 7-16. PMID 25818385

- 148. Elad S, Cheng KKF, Lalla RV, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. Cancer. Oct 01 2020; 126(19): 4423-4431. PMID 32786044
- 149. Martin RL, Chimenti R, Cuddeford T, et al. Achilles Pain, Stiffness, and Muscle Power Deficits: Midportion Achilles Tendinopathy Revision 2018. J Orthop Sports Phys Ther. May 2018; 48(5): A1-A38. PMID 29712543
- 150. Carcia CR, Martin RL, Houck J, et al. Achilles pain, stiffness, and muscle power deficits: achilles tendinitis. J Orthop Sports Phys Ther. Sep 2010; 40(9): A1-26. PMID 20805627
- 151. National Institute for Health and Care Excellence (NICE). Low back pain in adults: early management [CG88]. 2009; https://www.nice.org.uk/guidance/CG88. Accessed April 26, 2022.
- 152. North American Spine Society. Evidence-based clinical guidelines for multidisciplinary spine care. 2020. https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.p df. Accessed April 25, 2022.
- 153. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. Ann Intern Med. Apr 04 2017; 166(7): 514-530. PMID 28192789
- 154.Qaseem A, McLean RM, O'Gurek D, et al. Nonpharmacologic and Pharmacologic Management of Acute Pain From Non-Low Back, Musculoskeletal Injuries in Adults: A Clinical Guideline From the American College of Physicians and American Academy of Family Physicians. Ann Intern Med. Nov 03 2020; 173(9): 739-748. PMID 32805126
- 155. American Academy of Orthopaedic Surgeons. Management of Carpal Tunnel Syndrome: Evidence-Based Clinical Guideline. 2016; https://www.aaos.org/quality/quality-programs/upper-extremityprograms/carpal-tunnel-syndrome/. Accessed April 25, 2022.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

