

EFFECTIVE DATE: 10|01|2023

POLICY LAST UPDATED: 09|06|2023

OVERVIEW

Mitochondrial diseases are multisystem diseases that arise from dysfunction in the mitochondrial protein complexes involved in oxidative metabolism. There are many related but distinct syndromes and some patients have overlapping syndromes. As a result, these disorders can be difficult to diagnose. Genetic testing has the potential to improve the accuracy of diagnosis for mitochondrial diseases. Genetic testing also has the potential to determine future risk of disease in individuals who have a close relative with a pathogenic variant.

The following test(s) is addressed in this policy:

- Genomic Unity® Comprehensive Mitochondrial Disorders Analysis (Variantyx, Inc.) CPT code 0417U

MEDICAL CRITERIA

Medicare Advantage Plans and Commercial Products

Genetic testing to establish a genetic diagnosis of a mitochondrial disorder may be considered **medically necessary** when signs and symptoms of a mitochondrial disorder are present and genetic testing may eliminate the need for muscle biopsy.

Targeted genetic testing for a known familial variant in at-risk relatives may be considered **medically necessary** as preconceptional carrier testing under the following conditions:

- There is a defined mitochondrial disorder in the family of sufficient severity to cause impairment of quality of life or functional status; AND
- A variant that is known to be pathogenic for that specific mitochondrial disorder has been identified in the index case.

PRIOR AUTHORIZATION

Medicare Advantage Plans and Commercial Products

Prior authorization is required for Medicare Advantage Plans and recommended for Commercial Products for the following Proprietary Laboratory Analyses (PLA) tests:

- Genomic Unity® Comprehensive Mitochondrial Disorders Analysis – CPT code 0417U

Note: Laboratories are not allowed to obtain clinical authorization or participate in the authorization process on behalf of the ordering physician. Only the ordering physician shall be involved in the authorization, appeal or other administrative processes related to prior authorization/medical necessity.

In no circumstance shall a laboratory or a physician/provider use a representative of a laboratory or anyone with a relationship to a laboratory and/or a third party to obtain authorization on behalf of the ordering physician, to facilitate any portion of the authorization process or any subsequent appeal of a claim where the authorization process was not followed and/or a denial for clinical appropriateness was issued, including any element of the preparation of necessary documentation of clinical appropriateness. If a laboratory or a third party is found to be supporting any portion of the authorization process, BCBSRI will deem the action a violation of this policy and severe action will be taken up to and including termination from the BCBSRI provider network. If a laboratory provides a laboratory service that has not been authorized, the service will be denied as the financial liability of the participating laboratory and may not be billed to the member.

POLICY STATEMENT

Medicare Advantage Plans and Commercial Products

The following Proprietary Laboratory Analyses (PLA) tests may be considered medically necessary when the medical criteria above have been met:

- Genomic Unity® Comprehensive Mitochondrial Disorders Analysis – CPT code 0417U

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable benefits/coverage.

BACKGROUND

Mitochondrial DNA

Mitochondria are organelles within each cell that contain their own set of DNA, distinct from the nuclear DNA (nDNA) that makes up most of the human genome. Human mitochondrial DNA (mtDNA) consists of 37 genes. Thirteen genes code for protein subunits of the mitochondrial oxidative phosphorylation complex and the remaining 24 genes are responsible for proteins involved in the translation and/or assembly of the mitochondrial complex. Additionally, there are over 1000 nuclear genes coding for proteins that support mitochondrial function. The protein products from these genes are produced in the nucleus and later migrate to the mitochondria.

Mitochondrial DNA differs from nDNA in several important ways. Inheritance of mtDNA does not follow traditional Mendelian patterns. Rather, mtDNA is inherited only from maternal DNA so disorders that result from variants in mtDNA can only be passed on by the mother. Also, there are thousands of copies of each mtDNA gene in each cell, as opposed to nDNA, which contains only 1 copy per cell. Because there are many copies of each gene, variants may be present in some copies of the gene but not others. This phenomenon is called heteroplasmy. Heteroplasmy can be expressed as a percentage of genes that have the variant ranging from 0% to 100%. Clinical expression of the variant will generally depend on a threshold effect (ie, clinical symptoms will begin to appear when the percentage of mutated genes exceeds a threshold amount).

Mitochondrial Diseases

Primary mitochondrial diseases arise from dysfunction of the mitochondrial respiratory chain. The mitochondrial respiratory chain is responsible for aerobic metabolism, and dysfunction, therefore, affects a wide variety of physiologic pathways dependent on aerobic metabolism. Organs with a high-energy requirement, such as the central nervous system, cardiovascular system, and skeletal muscle, are preferentially affected by mitochondrial dysfunction.

The prevalence of these disorders has risen over the last 2 decades as the pathophysiology and clinical manifestations have been better characterized. It is currently estimated that the minimum prevalence of primary mitochondrial diseases is at least 1 in 5000.

Some specific mitochondrial diseases are listed next:

- Mitochondrial encephalopathy with lactic acidosis and stroke-like symptoms (MELAS) syndrome;
- Myoclonus epilepsy with ragged red fibers syndrome (MERRF);
- Kearns-Sayre syndrome;
- Leigh syndrome;
- Chronic progressive external ophthalmoplegia (CPEO);
- Leber hereditary optic neuropathy (LHON);
- Neuropathy, ataxia, and retinitis pigmentosa (NARP).

Most of these disorders are characterized by multisystem dysfunction, which generally includes myopathies and neurologic dysfunction and may involve multiple other organs. Each defined mitochondrial disease has a characteristic set of signs or symptoms. The severity of illness is heterogeneous and can vary markedly. Some patients will have only mild symptoms for which they never require medical care, while other patients have severe symptoms, a large burden of morbidity, and a shortened life expectancy.

Diagnosis

The diagnosis of mitochondrial diseases can be difficult. The individual symptoms are nonspecific, and symptom patterns can overlap considerably. As a result, a patient often cannot be easily classified into a particular syndrome. Biochemical testing is indicated for patients who do not have a clear clinical picture of a specific disorder. Measurement of serum lactic acid is often used as a screening test but the test is neither sensitive nor specific for mitochondrial diseases.

A muscle biopsy can be performed if the diagnosis is uncertain after biochemical workup. However, this invasive test is not definitive in all cases. The presence of "ragged red fibers" on histologic analysis is consistent with a mitochondrial disease. Ragged red fibers represent a proliferation of defective mitochondria. This characteristic finding may not be present in all types of mitochondrial diseases and also may be absent early in the course of disease.

Treatment

Treatment of mitochondrial disease is largely supportive because there are no specific therapies that impact the natural history of the disorder. Identification of complications such as diabetes and cardiac dysfunction is important for early treatment of these conditions. A number of vitamins and cofactors (eg, coenzyme Q, riboflavin) have been used but empirical evidence of benefit is lacking.⁶ Exercise therapy for myopathy is often prescribed but the effect on clinical outcomes is uncertain.⁵ The possibility of gene transfer therapy is under consideration but is at an early stage of development and untested in clinical trials.

Genetic Testing

Mitochondrial diseases can be caused by pathogenic variants in the maternally inherited mtDNA or one of many nDNA genes. Genetic testing for mitochondrial diseases may involve testing for point mutations, deletion and duplication analysis, and/or whole exome sequencing of nuclear or mtDNA. The type of testing done depends on the specific disorder being considered. For some primary mitochondrial diseases such as MELAS and MERFF, most variants are point mutations, and there is a finite number of variants associated with the disorder. When testing for one of these disorders, known pathogenic variants can be tested for with polymerase chain reaction, or sequence analysis can be performed on the particular gene. For other mitochondrial diseases, such as CPEO and Kearns-Sayre syndrome, the most common variants are deletions, and therefore duplication and deletion analysis would be the first test when these disorders are suspected. Table 1 provides examples of clinical symptoms and particular genetic variants in mtDNA or nDNA associated with particular mitochondrial syndromes. A repository of published and unpublished data on variants in human mtDNA is available in the MITOMAP database. Lists of mtDNA and nDNA genes that may lead to mitochondrial diseases and testing laboratories in the U.S. are provided at Genetic Testing Registry of the National Center for Biotechnology Information website.

For individuals with signs and/or symptoms of a mitochondrial disease who receive genetic testing, the evidence includes case series and cohort studies. Relevant outcomes are test validity, other test performance measures, symptoms, functional outcomes, health status measures, and quality of life. There is some evidence on clinical validity that varies by the patient population and testing strategy. Studies reporting diagnostic yield for known pathogenic variants using next-generation sequencing (NGS) panels tend to report rates ranging from 15% to 25%. Clinical specificity is unknown, but population-based studies have indicated that the prevalence of certain variants exceeds the prevalence of clinical disease, suggesting that the variant will be found in some people without the clinical disease (false-positives). Clinical utility is relatively high for confirming the diagnosis of mitochondrial diseases in people who have signs and symptoms of the disease. In these patients, a positive result in genetic testing can avoid a muscle biopsy and eliminate the need for further clinical workup. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close relative with a mitochondrial disease and a known pathogenic variant and who receive targeted familial variant testing, the evidence includes case series and cohort studies. Relevant outcomes are test validity, other test performance measures, changes in reproductive decision making, symptoms, functional outcomes, health status measures, and quality of life. Clinical validity is expected to be high for targeted testing of a known familial variant, assuming sufficient analytic validity. Clinical utility can be demonstrated by testing at-risk family members who have a close relative with a pathogenic variant. When a specific mitochondrial disease is present in the family that is severe enough to cause impairment and/or

disability, genetic testing may impact reproductive decision making. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

CODING

Medicare Advantage Plans and Commercial Products

The following CPT code(s) are covered for Medicare Advantage Plans and Commercial Products when medical criteria above are met:

This code can be used for Genomic Unity® Comprehensive Mitochondrial Disorders Analysis:

0417U Rare diseases (constitutional/heritable disorders), whole mitochondrial genome sequence with heteroplasmy detection and deletion analysis, nuclear-encoded mitochondrial gene analysis of 335 nuclear genes, including sequence changes (New Code Effective 10/01/2023)

RELATED POLICIES

Proprietary Laboratory Analyses (PLA)

PUBLISHED

Provider Update, October 2023

REFERENCES

1. Schon EA, DiMauro S, Hirano M. Human mitochondrial DNA: roles of inherited and somatic mutations. *Nat Rev Genet.* Dec 2012; 13(12): 878-90. PMID 23154810
2. Wong LJ. Diagnostic challenges of mitochondrial DNA disorders. *Mitochondrion.* Feb-Apr 2007; 7(1-2): 45-52. PMID 17276740
3. DiMauro S, Schon EA. Mitochondrial DNA mutations in human disease. *Am J Med Genet.* 2001; 106(1): 18-26. PMID 11579421
4. Falk MJ, Sondheimer N. Mitochondrial genetic diseases. *Curr Opin Pediatr.* Dec 2010; 22(6): 711-6. PMID 21045694
5. Chinnery PF. Mitochondrial Disorders Overview. In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews.* Seattle, WA: University of Washington; 2014.
6. Chinnery P, Majamaa K, Turnbull D, et al. Treatment for mitochondrial disorders. *Cochrane Database Syst Rev.* Jan 25 2006; (1): CD004426. PMID 16437486
7. Angelini C, Bello L, Spinazzi M, et al. Mitochondrial disorders of the nuclear genome. *Acta Myol.* Jul 2009; 28(1): 16-23. PMID 19772191
8. FOSWIKI. MITOMAP: a human mitochondrial genome database. 2018; <https://www.mitomap.org//MITOMAP>. Accessed August 5, 2022.
9. National Center for Biotechnology Information. GTR: Genetic Testing Registry. n.d.; <https://www.ncbi.nlm.nih.gov/gtr/>. Accessed August 5, 2022.
10. Riley LG, Cowley MJ, Gayevskiy V, et al. The diagnostic utility of genome sequencing in a pediatric cohort with suspected mitochondrial disease. *Genet Med.* Jul 2020; 22(7): 1254-1261. PMID 32313153
11. Nogueira C, Silva L, Pereira C, et al. Targeted next generation sequencing identifies novel pathogenic variants and provides molecular diagnoses in a cohort of pediatric and adult patients with unexplained mitochondrial dysfunction. *Mitochondrion.* Jul 2019; 47: 309-317. PMID 30831263
12. Fang F, Liu Z, Fang H, et al. The clinical and genetic characteristics in children with mitochondrial disease in China. *Sci China Life Sci.* Jul 2017; 60(7): 746-757. PMID 28639102
13. Legati A, Reyes A, Nasca A, et al. New genes and pathomechanisms in mitochondrial disorders unraveled by NGS technologies. *Biochim Biophys Acta.* Aug 2016; 1857(8): 1326-1335. PMID 26968897
14. Pronicka E, Piekutowska-Abramczuk D, Ciara E, et al. New perspective in diagnostics of mitochondrial disorders: two years' experience with whole-exome sequencing at a national paediatric centre. *J Transl Med.* Jun 12 2016; 14(1): 174. PMID 27290639
15. Kohda M, Tokuzawa Y, Kishita Y, et al. A Comprehensive Genomic Analysis Reveals the Genetic Landscape of Mitochondrial Respiratory Chain Complex Deficiencies. *PLoS Genet.* Jan 2016; 12(1): e1005679. PMID 26741492
16. Wortmann SB, Koolen DA, Smeitink JA, et al. Whole exome sequencing of suspected mitochondrial patients in clinical practice. *J Inher Metab Dis.* May 2015; 38(3): 437-43. PMID 25735936

17. Ohtake A, Murayama K, Mori M, et al. Diagnosis and molecular basis of mitochondrial respiratory chain disorders: exome sequencing for disease gene identification. *Biochim Biophys Acta*. Apr 2014; 1840(4): 1355-9. PMID 24462578
18. Taylor RW, Pyle A, Griffin H, et al. Use of whole-exome sequencing to determine the genetic basis of multiple mitochondrial respiratory chain complex deficiencies. *JAMA*. Jul 02 2014; 312(1): 68-77. PMID 25058219
19. Lieber DS, Calvo SE, Shanahan K, et al. Targeted exome sequencing of suspected mitochondrial disorders. *Neurology*. May 07 2013; 80(19): 1762-70. PMID 23596069
20. DaRe JT, Vasta V, Penn J, et al. Targeted exome sequencing for mitochondrial disorders reveals high genetic heterogeneity. *BMC Med Genet*. Nov 11 2013; 14: 118. PMID 24215330
21. McCormick E, Place E, Falk MJ. Molecular genetic testing for mitochondrial disease: from one generation to the next. *Neurotherapeutics*. Apr 2013; 10(2): 251-61. PMID 23269497
22. Calvo SE, Compton AG, Hershman SG, et al. Molecular diagnosis of infantile mitochondrial disease with targeted next-generation sequencing. *Sci Transl Med*. Jan 25 2012; 4(118): 118ra10. PMID 22277967
23. Qi Y, Zhang Y, Wang Z, et al. Screening of common mitochondrial mutations in Chinese patients with mitochondrial encephalomyopathies. *Mitochondrion*. Feb-Apr 2007; 7(1-2): 147-50. PMID 17276742
24. Deschauer M, Krasnianski A, Zierz S, et al. False-positive diagnosis of a single, large-scale mitochondrial DNA deletion by Southern blot analysis: the role of neutral polymorphisms. *Genet Test*. 2004; 8(4): 395-9. PMID 15684869
25. Elliott HR, Samuels DC, Eden JA, et al. Pathogenic mitochondrial DNA mutations are common in the general population. *Am J Hum Genet*. Aug 2008; 83(2): 254-60. PMID 18674747
26. Majamaa K, Moilanen JS, Uimonen S, et al. Epidemiology of A3243G, the mutation for mitochondrial encephalomyopathy, lactic acidosis, and strokelike episodes: prevalence of the mutation in an adult population. *Am J Hum Genet*. Aug 1998; 63(2): 447-54. PMID 9683591
27. DiMauro S, Hirano M. Melas. In: Pagon RA, Adam MP, Ardinger HH, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2013.
28. Jean-Francois MJ, Lertrit P, Berkovic SF, et al. Heterogeneity in the phenotypic expression of the mutation in the mitochondrial tRNA(Leu) (UUR) gene generally associated with the MELAS subset of mitochondrial encephalomyopathies. *Aust N Z J Med*. Apr 1994; 24(2): 188-93. PMID 8042948
29. Pfeiffer G, Majamaa K, Turnbull DM, et al. Treatment for mitochondrial disorders. *Cochrane Database Syst Rev*. Apr 18 2012; (4): CD004426. PMID 22513923
30. Parikh S, Goldstein A, Koenig MK, et al. Diagnosis and management of mitochondrial disease: a consensus statement from the Mitochondrial Medicine Society. *Genet Med*. Sep 2015; 17(9): 689-701. PMID 25503498

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

