

Medical Coverage Policy | Disposable Negative Pressure Wound Therapy



EFFECTIVE DATE: 07|01|2017

POLICY LAST REVIEWED: 02|07|2024

OVERVIEW

Negative pressure wound therapy (NPWT) involves the use of negative pressure or suction device to aspirate and remove fluids, debris, and infectious materials from the wound bed to promote the formation of granulation tissue and wound healing.

This policy is applicable only to disposable negative pressure wound therapy.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

The use of (powered or nonpowered) disposable single-use negative pressure wound therapy (NPWT) system devices for the treatment of acute or chronic wounds including but not limited to diabetic, venous, surgical, and traumatic wounds, is not covered, as they do not meet the durable medical equipment (DME) benefit durability requirement.

Commercial Products

The use of (powered or nonpowered) disposable single-use negative pressure wound therapy (NPWT) system devices for the treatment of acute or chronic wounds including but not limited to diabetic, venous, surgical, and traumatic wounds, is considered not medically necessary, as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

The management and treatment of chronic wounds, including decubitus ulcers, is challenging. Most chronic wounds will heal only if the underlying cause (ie, venous stasis, pressure, infection) is addressed. Also, cleaning the wound to remove nonviable tissue, microorganisms, and foreign bodies is essential to create optimal conditions for either re-epithelialization (ie, healing by secondary intention) or preparation for wound closure with skin grafts or flaps (ie, healing by primary intention). Therefore, debridement, irrigation, whirlpool treatments, and wet-to-dry dressings are common components of chronic wound care.

Negative pressure wound therapy (NPWT) involves the use of a negative pressure therapy or suction device to aspirate and remove fluids, debris, and infectious materials from the wound bed to promote the formation of granulation tissue. The devices may also be used as an adjunct to surgical therapy or as an alternative to surgery in a debilitated patient. Although the exact mechanism has not been elucidated, it is hypothesized that negative pressure contributes to wound healing by removing excess interstitial fluid, increasing the vascularity of the wound, reducing edema, and/or creating beneficial mechanical forces that lead to cell growth and expansion.

Disposable negative pressure therapy or suction devices cleared by the U.S. Food and Drug Administration (FDA) for treating chronic wounds include but are not limited to: Smart Negative Pressure Wound Care System, PICO Single Use Negative Pressure Wound Therapy System (Smith & Nephew), and the Prevena™ Incision Management System (KCI) is designed specifically for closed surgical incisions.

A nonpowered (mechanical) NPWT system has also been developed; the Smart Negative Pressure Wound Care System is portable and lightweight (3 oz) and can be worn underneath clothing. This system consists of a cartridge, dressing, and strap; the cartridge acts as the negative pressure source. The system is reported to generate negative pressure levels similar to other NPWT systems. This system is fully disposable.

For individuals who have diabetic lower-extremity ulcers or amputation wounds who receive portable, single-use outpatient NPWT, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, quality of life (QOL), and treatment-related morbidity. A 2019 RCT compared the PICO device with standard NPWT. In this study, the PICO device demonstrated noninferiority for wound area reduction. A statistically significant benefit in complete wound closure was noted for patients with diabetic foot ulcers (DFUs), but was not duplicated in the per protocol population due to a high number of exclusions. One study of the Smart Negative Pressure nonpowered Wound Care System (SNaP) showed noninferiority to a Vacuum-Assisted Closure Therapy (V.A.C.) device for wound size reduction. No significant difference in complete wound closure was reported. Interpretation of this study is limited by a high loss to follow-up. Well-designed comparative studies with larger numbers of patients powered to detect differences in complete wound closure are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have lower-extremity ulcers due to venous insufficiency who receive portable, single-use outpatient NPWT, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity. A 2019 RCT compared the PICO device with standard NPWT. In this study, the PICO device demonstrated noninferiority for wound area reduction. No significant benefit in complete wound closure was found in patients with venous ulcers. One study of the SNaP System showed noninferiority to a V.A.C. device for wound size reduction. A subgroup analysis of this study found a significant difference in complete wound closure for patients with venous ulcers. However, interpretation of this study is limited by a high loss to follow-up and a lack of a control group treated with standard dressings. Well-designed comparative studies with larger numbers of patients powered to detect differences in complete wound closure are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have traumatic or surgical wounds who receive portable, single-use outpatient NPWT, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity. The PICO device was studied in an adequately powered but unblinded RCT of combined in- and outpatient use after total joint arthroplasty. The evidence base for the Prevena System is not sufficiently robust for conclusions on efficacy to be drawn. Well-designed comparative studies with larger numbers of patients treated in an outpatient setting are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

CODING

Medicare Advantage Plans and Commercial Products

The following HCPCS code(s) is not covered for Medicare Advantage Plans (as it does not meet the DME benefit durability requirement) and is not medically necessary for Commercial Products:

A9272 Wound suction, disposable, includes dressing, all accessories and components, any type, each

The following CPT code(s) are not covered for Medicare Advantage Plans and not medically necessary for Commercial Products:

97607 Negative pressure wound therapy, (eg, vacuum assisted drainage collection), utilizing disposable, non-durable medical equipment including provision of exudate management collection system, topical

application(s), wound assessment, and instructions for ongoing care, per session; total wound(s) surface area less than or equal to 50 square centimeters

- 97608** Negative pressure wound therapy, (eg, vacuum assisted drainage collection), utilizing disposable, non-durable medical equipment including provision of exudate management collection system, topical application(s), wound assessment, and instructions for ongoing care, per session; total wound(s) surface area greater than 50 square centimeters

RELATED POLICIES

Durable Medical Equipment

PUBLISHED

Provider Update, April 2024
Provider Update, April 2023
Provider Update, June 2022
Provider Update, May 2021
Provider Update, April 2020

REFERENCES

- 1.U.S. Food and Drug Administration. UPDATE on Serious Complications Associated with Negative Pressure WoundTherapy Systems: FDA Safety Communication. 2011 Feb; <http://wayback.archive-it.org/7993/20170722215801/https://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm244211.htm>. Accessed November 11, 2022.
- 2.Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Vacuum-assisted closure in the treatment ofchronic wounds. TEC Assessments. 2000;Volume 15:Tab 23.
- 3.Food and Drug Administration (FDA). Guidance for Industry. Chronic Cutaneous Ulcer and Burn Wounds - DevelopingProducts for Treatment. June 2006; <https://www.fda.gov/media/71278/download>. Accessed November 15, 2023.
- 4.Rhee SM, Valle MF, Wilson LM, et al. Negative Pressure Wound Therapy Technologies For Chronic Wound Care in theHome Setting. Evidence Report/Technology Assessment (Contract No. 290-201-200007-I) Rockville, MD: Agency forHealthcare Research and Quality; 2014.
- 5.Rhee SM, Valle MF, Wilson LM, et al. Negative pressure wound therapy technologies for chronic wound care in the homesetting: A systematic review. *Wound Repair Regen*. 2015; 23(4): 506-17. PMID 25845268
- 6.Sullivan N, Snyder DL, Tipton K, et al. Technology assessment: Negative pressure wound therapy devices (Contract No.290-2007-10063). Rockville, MD: Agency for Healthcare Research and Quality; 2009.
- 7.Dumville JC, Hinchliffe RJ, Cullum N, et al. Negative pressure wound therapy for treating foot wounds in people withdiabetes mellitus. *Cochrane Database Syst Rev*. Oct 17 2013; (10): CD010318. PMID 24132761
- 8.Liu Z, Dumville JC, Hinchliffe RJ, et al. Negative pressure wound therapy for treating foot wounds in people with diabetesmellitus. *Cochrane Database Syst Rev*. Oct 17 2018; 10(10): CD010318. PMID 30328611
- 9.Wynn M, Freeman S. The efficacy of negative pressure wound therapy for diabetic foot ulcers: A systematised review. *JTissue Viability*. Aug 2019; 28(3): 152-160. PMID 31056407
- 10.Chen L, Zhang S, Da J, et al. A systematic review and meta-analysis of efficacy and safety of negative pressure woundtherapy in the treatment of diabetic foot ulcer. *Ann Palliat Med*. Oct 2021; 10(10): 10830-10839. PMID 34763444
- 11.Kirsner R, Dove C, Reyzelman A, et al. A prospective, randomized, controlled clinical trial on the efficacy of a single-usenegative pressure wound therapy system, compared to traditional negative pressure wound therapy in the treatment ofchronic ulcers of the lower extremities. *Wound Repair Regen*. Sep 2019; 27(5): 519-529. PMID 31087729
- 12.Kirsner RS, Zimnitsky D, Robinson M. A prospective, randomized, controlled clinical study on the effectiveness of a single-use negative pressure wound therapy system, compared to traditional negative pressure wound therapy in the treatmentof diabetic ulcers of the lower extremities. *Wound Repair Regen*. Nov 2021; 29(6): 908-911. PMID 34525239
- 13.Armstrong DG, Marston WA, Reyzelman AM, et al. Comparison of negative pressure wound therapy with an ultraportablemechanically powered device vs. traditional electrically powered device for the treatment of

- chronic lower extremity ulcers:a multicenter randomized-controlled trial. *Wound Repair Regen.* 2011; 19(2): 173-80. PMID 21362084
- 14.Armstrong DG, Marston WA, Reyzelman AM, et al. Comparative effectiveness of mechanically and electrically powerednegative pressure wound therapy devices: a multicenter randomized controlled trial. *Wound Repair Regen.* 2012; 20(3):332-41. PMID 22564228
- 15.Lerman B, Oldenbrook L, Eichstadt SL, et al. Evaluation of chronic wound treatment with the SNaP wound care systemversus modern dressing protocols. *Plast Reconstr Surg.* Oct 2010; 126(4): 1253-1261. PMID 20885246
- 16.Shi J, Gao Y, Tian J, et al. Negative pressure wound therapy for treating pressure ulcers. *Cochrane Database Syst Rev.* May 26 2023; 5(5): CD011334. PMID 37232410
- 17.Wanner MB, Schwarzl F, Strub B, et al. Vacuum-assisted wound closure for cheaper and more comfortable healing ofpressure sores: a prospective study. *Scand J Plast Reconstr Surg Hand Surg.* 2003; 37(1): 28-33. PMID 12625392
- 18.Dumville JC, Land L, Evans D, et al. Negative pressure wound therapy for treating leg ulcers. *Cochrane Database SystRev.* Jul 14 2015; 2015(7): CD011354. PMID 26171910
- 19.Vuerstaek JD, Vainas T, Wuite J, et al. State-of-the-art treatment of chronic leg ulcers: A randomized controlled trialcomparing vacuum-assisted closure (V.A.C.) with modern wound dressings. *J Vasc Surg.* Nov 2006; 44(5): 1029-37;discussion 1038. PMID 17000077
- 20.Marston WA, Armstrong DG, Reyzelman AM, et al. A Multicenter Randomized Controlled Trial Comparing Treatment ofVenous Leg Ulcers Using Mechanically Versus Electrically Powered Negative Pressure Wound Therapy. *Adv Wound Care(New Rochelle).* Feb 01 2015; 4(2): 75-82. PMID 25713749
- 21.Dumville JC, Munson C, Christie J. Negative pressure wound therapy for partial-thickness burns. *Cochrane DatabaseSyst Rev.* Dec 15 2014; 2014(12): CD006215. PMID 25500895
- 22.Bloemen MC, van der Wal MB, Verhaegen PD, et al. Clinical effectiveness of dermal substitution in burns by topicalnegative pressure: a multicenter randomized controlled trial. *Wound Repair Regen.* 2012; 20(6): 797-805. PMID 23110478
- 23.Krug E, Berg L, Lee C, et al. Evidence-based recommendations for the use of Negative Pressure Wound Therapy intraumatic wounds and reconstructive surgery: steps towards an international consensus. *Injury.* Feb 2011; 42 Suppl 1: S1-12. PMID 21316515
- 24.Ehrl D, Heidekrueger PI, Broer PN, et al. Topical Negative Pressure Wound Therapy of Burned Hands: FunctionalOutcomes. *J Burn Care Res.* Jan 01 2018; 39(1): 121-128. PMID 28368916
- 25.Norman G, Shi C, Goh EL, et al. Negative pressure wound therapy for surgical wounds healing by primary closure.*Cochrane Database Syst Rev.* Apr 26 2022; 4(4): CD009261. PMID 35471497
- 26.Li HZ, Xu XH, Wang DW, et al. Negative pressure wound therapy for surgical site infections: a systematic review andmeta-analysis of randomized controlled trials. *Clin Microbiol Infect.* Nov 2019; 25(11): 1328-1338. PMID 31220604
- 27.De Vries FEE, Wallert ED, Solomkin JS, et al. A systematic review and meta-analysis including GRADE qualification of therisk of surgical site infections after prophylactic negative pressure wound therapy compared with conventional dressings inclean and contaminated surgery. *Medicine (Baltimore).* Sep 2016; 95(36): e4673. PMID 27603360
- 28.Iheozor-Ejiofor Z, Newton K, Dumville JC, et al. Negative pressure wound therapy for open traumatic wounds. *CochraneDatabase Syst Rev.* Jul 03 2018; 7(7): CD012522. PMID 29969521
- 29.Stannard JP, Volgas DA, McGwin G, et al. Incisional negative pressure wound therapy after high-risk lower extremityfractures. *J Orthop Trauma.* Jan 2012; 26(1): 37-42. PMID 21804414
- 30.Costa ML, Achten J, Bruce J, et al. Effect of Negative Pressure Wound Therapy vs Standard Wound Management on 12-Month Disability Among Adults With Severe Open Fracture of the Lower Limb: The WOLFF Randomized Clinical Trial.*JAMA.* Jun 12 2018; 319(22): 2280-2288. PMID 29896626
- 31.Seidel D, Diedrich S, Herrle F, et al. Negative Pressure Wound Therapy vs Conventional Wound Treatment inSubcutaneous Abdominal Wound Healing Impairment: The SAWHI Randomized Clinical Trial. *JAMA Surg.* Jun 01 2020;155(6): 469-478. PMID 32293657
- 32.Stannard JP, Robinson JT, Anderson ER, et al. Negative pressure wound therapy to treat hematomas and surgicalincisions following high-energy trauma. *J Trauma.* Jun 2006; 60(6): 1301-6. PMID 16766975

33. Monsen C, Acosta S, Mani K, et al. A randomised study of NPWT closure versus alginate dressings in peri-vascular groin infections: quality of life, pain and cost. *J Wound Care*. Jun 2015; 24(6): 252, 254-6, 258-0. PMID 26075373
34. Costa ML, Achten J, Parsons NR. Five-year outcomes for patients sustaining severe fractures of the lower limb : mid-term results from the Wound management for Open Lower Limb Fracture (WOLLF) trial. *Bone Joint J*. May 2022; 104-B(5):633-639. PMID 35491582
35. Seidel D, Lefering R. NPWT Resource Use Compared With Conventional Wound Treatment in Subcutaneous Abdominal Wounds With Healing Impairment After Surgery: SAWHI Randomized Clinical Trial Results. *Ann Surg*. Feb 01 2022; 275(2): e290-e298. PMID 34117147
36. Karlakki SL, Hamad AK, Whittall C, et al. Incisional negative pressure wound therapy dressings (iNPWTd) in routine primary hip and knee arthroplasties: A randomised controlled trial. *Bone Joint Res*. Aug 2016; 5(8): 328-37. PMID 27496913
37. Peterson AT, Bakaysa SL, Driscoll JM, et al. Randomized controlled trial of single-use negative-pressure wound therapy dressings in morbidly obese patients undergoing cesarean delivery. *Am J Obstet Gynecol MFM*. Sep 2021; 3(5): 100410. PMID 34058423
38. Gonzalez MG, Barske ME, Kjellsson KB, et al. Topical negative pressure wound therapy to prevent wound complications following caesarean delivery in high-risk obstetric patients: A randomised controlled trial. *Aust N Z J Obstet Gynaecol*. Aug 2023; 63(4): 516-520. PMID 37140175
39. Pauser J, Nordmeyer M, Biber R, et al. Incisional negative pressure wound therapy after hemiarthroplasty for femoral neck fractures - reduction of wound complications. *Int Wound J*. Oct 2016; 13(5): 663-7. PMID 25125244
40. Murphy PB, Knowles S, Chadi SA, et al. Negative Pressure Wound Therapy Use to Decrease Surgical Nosocomial Events in Colorectal Resections (NEPTUNE): A Randomized Controlled Trial. *Ann Surg*. Jul 2019; 270(1): 38-42. PMID 30499799
41. Hussamy DJ, Wortman AC, McIntire DD, et al. Closed Incision Negative Pressure Therapy in Morbidly Obese Women Undergoing Cesarean Delivery: A Randomized Controlled Trial. *Obstet Gynecol*. Oct 2019; 134(4): 781-789. PMID 31503147
42. Tuuli MG, Liu J, Tita ATN, et al. Effect of Prophylactic Negative Pressure Wound Therapy vs Standard Wound Dressing on Surgical-Site Infection in Obese Women After Cesarean Delivery: A Randomized Clinical Trial. *JAMA*. Sep 22 2020; 324(12): 1180-1189. PMID 32960242
43. Bertges DJ, Smith L, Scully RE, et al. A multicenter, prospective randomized trial of negative pressure wound therapy for infrainguinal revascularization with a groin incision. *J Vasc Surg*. Jul 2021; 74(1): 257-267.e1. PMID 33548422
44. Ceppa EP, Kim RC, Niedzwiecki D, et al. Closed Incision Negative Pressure Therapy to Reduce Surgical Site Infection in High-Risk Gastrointestinal Surgery: A Randomized Controlled Trial. *J Am Coll Surg*. Apr 01 2023; 236(4): 698-708. PMID 36728375
45. American Academy of Orthopaedic Surgeons. Prevention of Surgical Site Infections After Major Extremity Trauma Evidence-Based Clinical Practice Guideline. www.aaos.org/SSItraumacpg. Published 03/21/22. Accessed November 15, 2023.
46. Qaseem A, Humphrey LL, Forcica MA, et al. Treatment of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. Mar 03 2015; 162(5): 370-9. PMID 25732279
47. Association for the Advancement of Wound Care (AAWC). International Consolidated Venous Ulcer Guideline (ICVUG). Update of AAWC Venous Ulcer Guideline, 2005 and 2010. 2015; <https://aawconline.memberclicks.net/assets/appendix%20c%20guideline%20icvug-textformatrecommendations-final%20v42%20changessaved18aug17.pdf>. Accessed November 16, 2023.
48. Willy C, Agarwal A, Andersen CA, et al. Closed incision negative pressure therapy: international multidisciplinary consensus recommendations. *Int Wound J*. Apr 2017; 14(2): 385-398. PMID 27170231
49. Senneville É, Albalawi Z, van Asten SA, et al. IWGDF/IDSA guidelines on the diagnosis and treatment of diabetes-related foot infections (IWGDF/IDSA 2023). *Diabetes Metab Res Rev*. Oct 01 2023; e3687. PMID 37779323
50. National Institute for Health and Care Excellence (NICE). Negative Pressure Wound Therapy for the Open Abdomen [IPG467]. 2013; <https://www.nice.org.uk/guidance/ipg467>. Accessed November 11, 2023.

51. National Institute for Health and Care Excellence (NICE). Diabetic Foot Problems: Prevention and Management [NG19]. Published August 2015; Updated October 2019; <https://www.nice.org.uk/guidance/ng19/evidence>. Accessed November 14, 2023.
52. National Institute for Health and Care Excellence (NICE). Pressure ulcers: prevention and management [CG179]. 2014; <https://www.nice.org.uk/guidance/cg179>. Accessed November 10, 2023.
53. National Institute for Health and Care Excellence (NICE). PICO negative pressure wound dressings for closed surgical incisions [MTG43]. 2019; <https://www.nice.org.uk/guidance/mtg43>. Accessed November 16, 2023.
54. National Institute for Health and Care Excellence (NICE). Cesarean birth [NG192]. Published: March 2021. Updated: September 2023; <https://www.nice.org.uk/guidance/ng192>. Accessed November 15, 2023.
55. National Institute for Health and Care Excellence (NICE). The VAC Veraflo Therapy system for acute infected or chronic wounds that are failing to heal [MTG54]. 2021; <https://www.nice.org.uk/guidance/mtg54>. Accessed November 9, 2023.
56. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): Negative Pressure Wound Therapy Pumps (L33821)
57. Centers for Medicare and Medicaid Services (CMS). Local Coverage Article: Billing and Coding: Article - Negative Pressure Wound Therapy Pumps - Policy Article (A52511)

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

