

Medical Coverage Policy | Pelvic Floor Stimulation as a Treatment of Urinary and Fecal Incontinence



EFFECTIVE DATE: 01|01|2026

POLICY LAST REVIEWED: 10|01|2025

OVERVIEW

Pelvic floor stimulation is proposed as a nonsurgical treatment option for women and men with urinary or fecal incontinence. This approach involves either electrical stimulation of pelvic floor musculature or extracorporeal pulsed magnetic stimulation.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

Pelvic floor electrical stimulation with a non-implantable stimulator is covered for Medicare Advantage Plans members only for the treatment of stress and/or urge urinary incontinence.

Pelvic floor magnetic stimulation with a non-implantable stimulator for the treatment of stress and/or urge urinary incontinence is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

Pelvic floor electrical or magnetic stimulation with a non-implantable stimulator for fecal incontinence is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

Note: Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all Medicare Advantage Plans policies. Therefore, Medicare Advantage Plans policies may differ from Commercial products. In some instances, benefits for Medicare Advantage Plans may be greater than what is allowed by the CMS.

Commercial Products

Pelvic floor electrical or magnetic stimulation with a non-implantable stimulator for urinary and fecal incontinence is not covered and considered contract exclusion.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable "Therapies, Acupuncture and Acupuncturist Services, and Biofeedback" benefits/coverage.

BACKGROUND

Commercial Products

Pelvic floor stimulation (PFS) involves electrical stimulation of pelvic floor muscles using either a probe wired to a device for controlling the electrical stimulation or, more recently, extracorporeal electromagnetic (also called magnetic) pulses. Stimulation of the pudendal nerve to activate the pelvic floor musculature may improve urethral closure. In addition, PFS is thought to improve partially denervated urethral and pelvic floor musculature by enhancing the process of reinnervation. Methods of electrical PFS have varied in location (eg, vaginal, rectal), stimulus frequency, stimulus intensity or amplitude, pulse duration, pulse to rest ratio,

treatments per day, number of treatment days per week, length of time for each treatment session, and overall time period for device use between clinical and home settings. Variations in the amplitude and frequency of the electrical pulse are used to mimic and stimulate the different physiologic mechanisms of the voiding response, depending on the etiology of the incontinence (ie, either detrusor instability, stress incontinence, or a mixed pattern). Magnetic PFS does not require an internal electrode; instead, patients sit fully clothed on a specialized chair with an embedded magnet. Patients receiving electrical PFS may undergo treatment in a physician's office or physical therapy facility, or patients may undergo initial training in a physician's office followed by home treatment with a rented or purchased pelvic floor stimulator. Magnetic PFS may be administered in the physician's office.

Urinary Incontinence

For individuals who have urinary incontinence who receive electrical PFS, the evidence includes systematic reviews of randomized controlled trials (RCTs). Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Findings from systematic reviews have not found that electrical PFS used to treat urinary incontinence in women consistently improves the net health outcome compared with placebo or other conservative treatments. Moreover, meta-analyses of RCTs have not found a significant benefit of electrical PFS in men with postprostatectomy incontinence compared with a control intervention. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have urinary incontinence who receive magnetic PFS, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. A systematic review of RCTs on magnetic PFS for urinary incontinence in women concluded that the evidence was insufficient due to the following factors: a low number of trials with short-term follow-up, methodologic limitations, as well as heterogeneity in patient populations, interventions, and outcomes reported. One RCT evaluating magnetic stimulation for treating men with postprostatectomy urinary incontinence reported short-term results favoring magnetic PFS; however, the trial was small and lacked a sham comparator. The evidence is insufficient to determine the effects of the technology on health outcomes.

Fecal Incontinence

For individuals who have fecal incontinence who receive electrical PFS, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Among the RCTs that have evaluated electrical PFS as a treatment for fecal incontinence only one trial was sham-controlled, and it did not find that electrical stimulation improved the net health outcome. Systematic reviews of RCTs have not found that electrical stimulation is superior to control interventions for treating fecal incontinence. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have fecal incontinence who receive magnetic PFS, the evidence includes no RCTs or non-RCTs. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. The evidence is insufficient to determine the effects of the technology on health outcomes.

Medicare Advantage Plans

According to Medicare National Coverage Determination, non-implantable pelvic floor electrical stimulators provide neuromuscular electrical stimulation through the pelvic floor with the intent of strengthening and exercising pelvic floor musculature. Stimulation is generally delivered by vaginal or anal probes connected to an external pulse generator. The methods of pelvic floor electrical stimulation vary in location, stimulus frequency (Hz), stimulus intensity or amplitude (mA), pulse duration (duty cycle), treatments per day, number of treatment days per week, length of time for each treatment session, overall time period for device use and between clinic and home settings. In general, the stimulus frequency and other parameters are chosen based on the patient's clinical diagnosis. Pelvic floor electrical stimulation with a non-implantable stimulator is covered for the treatment of stress and/or urge urinary incontinence in cognitively intact patients who have failed a documented trial of pelvic muscle exercise (PME) training. A failed trial of PME training is defined as

no clinically significant improvement in urinary continence after completing 4 weeks of an ordered plan of pelvic muscle exercises designed to increase periurethral muscle strength.

Several electrical stimulators have been cleared by the U.S. Food and Drug Administration (FDA). In 2006, the MyoTrac Infiniti™ (Thought Technology) and in 2015, the ApexM (InControl Medical), nonimplanted electrical stimulators for treating urinary incontinence, were cleared for marketing by the FDA through the 510(k) process. Predicate devices also used to treat urinary incontinence, including the Pathway™ CTS 2000 (Prometheus Group) and the InCare® PRS (Hollister). In 2011, the itouch Sure Pelvic Floor Exerciser (TensCare) was cleared for marketing.

In 2000, the NeoControl® Pelvic Floor Therapy System (Neotonus) cleared through the FDA 510(k) process for treating urinary incontinence in women. This device, formerly known as the Neotonus Model 1000 Magnetic Stimulator, provides noninvasive electromagnetic stimulation of pelvic floor musculature. The magnetic system is embedded in a chair seat; patients sit on the chair fully clothed and receive the treatment. The magnetic fields are controlled by a separate power unit.

In 2014, the InTone® MV (InControl Medical), a nonimplantable device that provides electrical stimulation and/or biofeedback via manometry, was cleared by the FDA. The device is intended to treat male and female urinary and fecal incontinence.

CODING

Medicare Advantage Plans for Urinary Incontinence

There are no specific CPT code(s) for this service and therefore the unlisted code should be used:

53899 Unlisted procedure, urinary system (to be used for pulsed magnetic stimulation for the treatment of incontinence)

The following HCPCS code(s) is covered for Medicare Advantage Plans only:

E0740 Nonimplanted pelvic floor electrical stimulator, complete system

Commercial Products for Urinary Incontinence

There are no specific CPT codes for this service and therefore the unlisted code should be used:

53899 Unlisted procedure, urinary system (to be used for pulsed magnetic stimulation for the treatment of incontinence)

The following HCPCS code(s) is not covered for Commercial Products.

E0740 Nonimplanted pelvic floor electrical stimulator, complete system

Medicare Advantage Plans and Commercial Products for Fecal Incontinence

There are no specific CPT code(s) for this service. Therefore, an unlisted code should be used.

RELATED POLICIES

Biofeedback

Unlisted Procedures

PUBLISHED

Provider Update, November/December 2025

Provider Update, November 2024

Provider Update, October 2023

Provider Update, January 2023

Provider Update, October 2021

REFERENCES

1. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD): Non-Implantable Pelvic Floor Electrical Stimulator (230.8); <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCDId=231>. Accessed September 23, 2025
2. Centers for Medicare & Medicaid Services (CMS). CMS Manual System: Pub 100-03 Medicare National Coverage Determinations; Transmittal 48. 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=231>. Accessed July 8, 2025.
3. Gorina Y, Schappert S, Bercovitz A, et al. Prevalence of incontinence among older americans. *Vital Health Stat* 3. Jun 2014; (36): 1-33. PMID 24964267
4. Markland AD, Goode PS, Redden DT, et al. Prevalence of urinary incontinence in men: results from the national health and nutrition examination survey. *J Urol*. Sep 2010; 184(3): 1022-7. PMID 20643440
5. Abdelbary AM, El-Dessoukey AA, Massoud AM, et al. Combined Vaginal Pelvic Floor Electrical Stimulation (PFS) and Local Vaginal Estrogen for Treatment of Overactive Bladder (OAB) in Perimenopausal Females. Randomized Controlled Trial (RCT). *Urology*. Sep 2015; 86(3): 482-6. PMID 26135813
6. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Pelvic floor electrical stimulation in the treatment of urinary incontinence in adults. *TEC Assessments*. 2000;Volume 15:Tab 2.
7. Leonardo K, Seno DH, Mirza H, et al. Biofeedback-assisted pelvic floor muscle training and pelvic electrical stimulation in women with overactive bladder: A systematic review and meta-analysis of randomized controlled trials. *Neurourol Urodyn*. Aug 2022; 41(6): 1258-1269. PMID 35686543
8. Stewart F, Berghmans B, Bø K, et al. Electrical stimulation with non-implanted devices for stress urinary incontinence in women. *Cochrane Database Syst Rev*. Dec 22 2017; 12(12): CD012390. PMID 29271482
9. Shamliyan T, Wyman J, Kane R. Nonsurgical Treatments for Urinary Incontinence in Adult Women: Diagnosis and Comparative Effectiveness (Comparative Effectiveness Review 36). Rockville, MD: Agency for Healthcare Research and Quality; 2012.
10. Moroni RM, Magnani PS, Haddad JM, et al. Conservative Treatment of Stress Urinary Incontinence: A Systematic Review with Meta-analysis of Randomized Controlled Trials. *Rev Bras Ginecol Obstet*. Feb 2016; 38(2): 97-111. PMID 26883864
11. Tang G, Liu M, Chen X, et al. Effectiveness of electrical stimulation for treating male urinary incontinence after prostatectomy: a meta-analysis and systematic review. *Int J Surg*. Jun 27 2025. PMID 40576184
12. Sciarra A, Viscuso P, Arditi A, et al. A biofeedback-guided programme or pelvic floor muscle electric stimulation can improve early recovery of urinary continence after radical prostatectomy: A meta-analysis and systematic review. *Int J Clin Pract*. Oct 2021; 75(10): e14208. PMID 33811418
13. Berghmans B, Hendriks E, Bernards A, et al. Electrical stimulation with non-implanted electrodes for urinary incontinence in men. *Cochrane Database Syst Rev*. Jun 06 2013; (6): CD001202. PMID 23740763
14. Johnson EE, Mamoulakis C, Stoniute A, et al. Conservative interventions for managing urinary incontinence after prostate surgery. *Cochrane Database Syst Rev*. Apr 18 2023; 4(4): CD014799. PMID 37070660
15. Zhu YP, Yao XD, Zhang SL, et al. Pelvic floor electrical stimulation for postprostatectomy urinary incontinence: a meta-analysis. *Urology*. Mar 2012; 79(3): 552-5. PMID 22386394
16. Cohen-Zubary N, Gingold-Belfer R, Lambort I, et al. Home electrical stimulation for women with fecal incontinence: a preliminary randomized controlled trial. *Int J Colorectal Dis*. Apr 2015; 30(4): 521-8. PMID 25619464
17. Norton C, Gibbs A, Kamm MA. Randomized, controlled trial of anal electrical stimulation for fecal incontinence. *Dis Colon Rectum*. Feb 2006; 49(2): 190-6. PMID 16362803
18. Vonthein R, Heimerl T, Schwandner T, et al. Electrical stimulation and biofeedback for the treatment of fecal incontinence: a systematic review. *Int J Colorectal Dis*. Nov 2013; 28(11): 1567-77. PMID 23900652
19. Schwandner T, König IR, Heimerl T, et al. Triple target treatment (3T) is more effective than biofeedback alone for anal incontinence: the 3T-AI study. *Dis Colon Rectum*. Jul 2010; 53(7): 1007-16. PMID 20551752
20. Schwandner T, Hemmelmann C, Heimerl T, et al. Triple-target treatment versus low-frequency electrostimulation for anal incontinence: a randomized, controlled trial. *Dtsch Arztebl Int*. Sep 2011; 108(39): 653-60. PMID 22013492

21. Hosker G, Cody JD, Norton CC. Electrical stimulation for faecal incontinence in adults. *Cochrane Database Syst Rev*. Jul 18 2007; 2007(3): CD001310. PMID 17636665
22. Lim R, Lee SW, Tan PY, et al. Efficacy of electromagnetic therapy for urinary incontinence: A systematic review. *Neurourol Urodyn*. Nov 2015; 34(8): 713-22. PMID 25251335
23. Lim R, Liong ML, Leong WS, et al. Pulsed Magnetic Stimulation for Stress Urinary Incontinence: 1-Year Followup Results. *J Urol*. May 2017; 197(5): 1302-1308. PMID 27871927
24. Unal B, Sarsan A, Yildiz N, et al. Efficacy of Magnetic Stimulation in Men With Urinary Incontinence After Radical Prostatectomy: A Randomized, Quadruple-Blind, Sham-Controlled Clinical Trial. *Neurourol Urodyn*. Jun 2025; 44(5): 1140-1148. PMID 40223765
25. Wald A, Bharucha AE, Limketkai B, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. *Am J Gastroenterol*. Oct 01 2021; 116(10): 1987-2008. PMID 34618700
26. Bordeianou LG, Thorsen AJ, Keller DS, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Fecal Incontinence. *Dis Colon Rectum*. May 01 2023; 66(5): 647-661. PMID 36799739
27. Cameron AP, Chung DE, Dielubanza EJ, et al. The AUA/SUFU Guideline on the Diagnosis and Treatment of Idiopathic Overactive Bladder. *J Urol*. Jul 2024; 212(1): 11-20. PMID 38651651
28. Sandhu JS, Breyer B, Comiter C, et al. Incontinence after Prostate Treatment: AUA/SUFU Guideline. *J Urol*. Aug 2019; 202(2): 369-378. PMID 31059663
29. Breyer BN, Kim SK, Kirkby E, et al. Updates to Incontinence After Prostate Treatment: AUA/GURS/SUFU Guideline (2024). *J Urol*. Oct 2024; 212(4): 531-538. PMID 38934789
30. National Institute for Health and Care Excellence (NICE) Guideline. Urinary Incontinence and Pelvic Organ Prolapse in Women: Management. NICE Guideline. 2019. <https://www.nice.org.uk/guidance/ng123>. Accessed July 7, 2025.
31. National Institute for Health and Care Excellence (NICE). Faecal incontinence in adults: management [CG49]. 2007; <https://www.nice.org.uk/guidance/cg49>. Accessed July 8, 2025.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

