

## Medical Coverage Policy | Lutathera (lutetium Lu 177 dotatate)



**EFFECTIVE DATE:** 07|01|2018

**POLICY LAST UPDATED:** 06|5|2018

### OVERVIEW

Lutathera (lutetium Lu 177 dotatate) is used for the treatment of somatostatin receptor-positive gastroenteropancreatic neuroendocrine tumors (GEP-NETs), including foregut, midgut, and hindgut neuroendocrine tumors in adults.

This policy is applicable to BlueCHiP for Medicare products only. For Commercial Products, see related policy section

### PRIOR AUTHORIZATION

Prior authorization is required for BlueCHiP for Medicare

### POLICY STATEMENT

#### Blue CHiP for Medicare

Lutathera is medically necessary when the criteria is met.

### MEDICAL CRITERIA

#### Initial Evaluation

Lutathera will be approved when ALL of the following are met:

1. ONE of the following:
  - A. There is documentation that the patient is currently being treated with the requested agent  
**OR**
  - B. The prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed  
**OR**
  - C. The patient has a diagnosis of somatostatin-positive, gastroenteropancreatic neuroendocrine tumor (GEP-NETS) AND BOTH of the following:
    - i. The patient has locally advanced, inoperable, or metastatic carcinoid tumor **AND**
    - ii. The patient will discontinue long-acting somatostatin analog (e.g. octreotide LAR) for at least 4 weeks prior to initiating the requested agent  
**OR**
  - D. The patient has another FDA approved indication for the requested agent  
**AND**
2. The prescriber is a specialist (e.g., oncologist) or the prescriber has consulted with a specialist  
**AND**
3. The patient does NOT have any FDA labeled contraindications to the requested agent  
**AND**
4. The requested dose is within FDA labeled dosing for the requested indication  
**AND**
5. The patient has NOT exceeded 4 treatment doses in lifetime

**Length of Approval:** GEP-NETs – 12 months for maximum 4 doses per lifetime; All other FDA approved diagnosis – 12 months.

### **Renewal Evaluation**

**Lutathera** will be approved when ALL of the following are met:

1. The patient has been previously approved for the requested agent through the Medical Drug Review process  
**AND**
2. Treatment-related toxicities (e.g., anemia, hepatotoxicity, neutropenia, renal toxicity, thrombocytopenia) are resolved prior to re-starting the requested agent
3. The patient has NOT exceeded 4 treatment doses in lifetime

**Length of Approval:** GEP-NETs – 12 months for maximum 4 doses per lifetime; All other FDA approved diagnosis – 12 months.

The requested agent will also be approved when the following are met:

1. The patient has been previously approved  
**AND**
2. Treatment-related toxicities (e.g., anemia, hepatotoxicity, neutropenia, renal toxicity, thrombocytopenia) are resolved prior to re-starting the requested agent

**Length of Approval:** 12 months

## **BACKGROUND**

### **Carcinoid Syndrome**

Carcinoid syndrome is a constellation of symptoms resulting from hormones secreted by carcinoid tumors. These tumors can either be active or inactive. Active tumors secrete various hormones and vasoactive peptide substances such as adrenocorticotropic hormone (ACTH), serotonin, histamine, and tachykinins. The most common symptoms of carcinoid syndrome are flushing and diarrhea.

Somatostatin analogs (octreotide or lanreotide) are recommended as initial therapy in patients with carcinoid syndrome. Addition of telotristat to somatostatin analog therapy is recommended for patients who are refractory to somatostatin analog therapy.

### **Neuroendocrine Tumors**

Neuroendocrine tumors are thought to arise from cells throughout the diffuse endocrine system. They comprise a broad family of tumors, the most common of which are carcinoid tumors (most commonly arising in the lungs and bronchi, small intestine, appendix, rectum, and thymus) and pancreatic neuroendocrine tumors. Other less common neuroendocrine tumors include those arising in the parathyroid, thyroid, adrenal, and pituitary glands.

Patients with neuroendocrine tumors may or may not have symptoms attributable to hormonal hypersecretion. These symptoms include intermittent flushing and diarrhea in patients with carcinoid syndrome, hypertension in patients with pheochromocytoma, and symptoms attributable to secretion of insulin, glucagon, gastrin, and other peptides in patients with pancreatic neuroendocrine tumors. Patients with hormonal symptoms are considered to have functional tumors and those without symptoms are considered to have nonfunctional tumors.

Two-thirds of carcinoid tumors arise in the GI tract, including the stomach, small intestine, appendix, and rectum. Neuroendocrine tumors of the GI tract, lung, or thymus may secrete various hormones and vasoactive peptides. Tumors arising in the small intestine or appendix are more commonly associated with

carcinoid syndrome. Approximately 50 to 66% of patients with carcinoid syndrome develop valvular cardiac complications consisting of tricuspid regurgitation and/or pulmonary stenosis.

Somatostatin receptor-based imaging is recommended to assess the somatostatin receptor status of locoregional unresectable and/or metastatic neuroendocrine tumors of the GI tract, lung, or thymus, if treatment with octreotide or lanreotide is being considered. Somatostatin analogs (octreotide or lanreotide) are recommended for control of symptoms and tumor growth in patients who have metastatic neuroendocrine tumors and carcinoid syndrome. The long-acting release (LAR) formulation of octreotide is commonly used for the chronic management of symptoms in patients with carcinoid syndrome. Telotristat in combination with octreotide or lanreotide should be considered for persistent diarrhea.

Approximately 1% of pancreatic cancers are malignant pancreatic neuroendocrine tumors. Majority of pancreatic neuroendocrine tumors are nonfunction (40 to 91%). The remaining manifest with clinically evident hormonal symptoms. VIPoma is the rarest form of pancreatic neuroendocrine tumors, also described as cholecystikininoma (CCKoma). Primary treatment approach for localized pancreatic neuroendocrine tumors is resection, when possible. For patients with unresectable disease, those who initially present with clinically significant tumor burden, or those with clinically significant disease progression, octreotide or lanreotide should be considered. Patients with pancreatic neuroendocrine tumors and symptoms of hormone secretion should, in most cases, receive treatment with either lanreotide or octreotide and/or other medication to manage their symptoms.

### **Efficacy**

The efficacy of Lutathera in patients with progressive, well-differentiated, locally advanced/inoperable or metastatic somatostatin receptor positive midgut carcinoid tumors was established in NETTER-1 (NCT01578239), a randomized, multicenter, open-label, active-controlled trial. Key eligibility criteria included Ki67 index  $\leq$  20%, Karnofsky performance status  $\geq$  60, confirmed presence of somatostatin receptors on all lesions (OctreoScan uptake  $\geq$  normal liver), creatinine clearance  $\geq$  50 mL/min, no prior treatment with peptide receptor radionuclide therapy (PRRT), and no prior external radiation therapy to more than 25% of the bone marrow.

Two hundred twenty-nine (229) patients were randomized (1:1) to receive either Lutathera 7.4 GBq (200 mCi) every 8 weeks for up to 4 administrations (maximum cumulative dose of 29.6 GBq) or high-dose long-acting octreotide (defined as 60 mg by intramuscular injection every 4 weeks). Patients in the Lutathera arm also received long-acting octreotide 30 mg as an intramuscular injection 4 to 24 hours after each Lutathera dose and every 4 weeks after completion of Lutathera treatment until disease progression or until week 76 of the study. Patients in both arms could receive short-acting octreotide for symptom management; however, short-acting octreotide was withheld for at least 24 hours before each Lutathera dose.

Randomization was stratified by OctreoScan tumor uptake score (Grade 2, 3 or 4) and the length of time that patients had been on the most recent constant dose of octreotide prior to randomization ( $\leq$  6 or  $>$  6 months). The major efficacy outcome measure was progression free survival (PFS) as determined by a blinded independent radiology committee (IRC) per RECIST v1.1. Additional efficacy outcome measures were overall response rate (ORR) by IRC, duration of response, and overall survival (OS).

### **COVERAGE**

Benefits may vary between groups and contracts. Please refer to the appropriate Evidence of Coverage for applicable physician administered drug services benefits/coverage.

### **CODING**

#### **Blue CHiP for Medicare and Commercial**

Prior to 10/1/18 there is no specific HCPCS code, claims must be filed with an unlisted code such as J3490 and the NDC number.

For claims between 10/1/18 to 12/31/18 use the following HCPCS code:  
C9031 Lutetium Lu 177, dotatate, therapeutic, 1 mCi

For claims after 12/31/18 use the following HCPCS code:  
A9513 Lutetium Lu 177, dotatate, therapeutic, 1 mCi

### RELATED POLICIES

Prior Authorization of Drugs

### PUBLISHED

Provider Update July 2018

### REFERENCES

1. Lutathera prescribing information. Advanced Accelerator Applications S.A. January 2018.
2. Jonathan R Strosberg. Treatment of carcinoid syndrome. UpToDate. Accessed September 2017.
3. NCCN Clinical Practice Guidelines. Neuroendocrine Tumors. Version 3.2017 – June 13,2017. Available at: [https://www.nccn.org/professionals/physician\\_gls/PDF/neuroendocrine.pdf](https://www.nccn.org/professionals/physician_gls/PDF/neuroendocrine.pdf). Accessed February 2018.

**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

