



EFFECTIVE DATE: 12|01|2014
POLICY LAST UPDATED: 05|21|2019

OVERVIEW

This policy documents coverage guidelines for BlueCHiP for Medicare and Commercial Products for enhanced external counterpulsation (EECP) used in outpatient treatment. EECP is a noninvasive treatment used to augment diastolic pressure, decrease left ventricular afterload, and increase venous return. It has been studied primarily as a treatment for patients with refractory angina and heart failure.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare

EECP is covered for BlueCHiP for Medicare members.

Note: Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all BlueCHiP for Medicare policies. Therefore, BlueCHiP for Medicare policies may differ from Commercial products. In some instances, benefits for BlueCHiP for Medicare may be greater than what is allowed by the CMS.

Commercial Products

EECP used in outpatient treatment is not medically necessary for all indications, including but not limited to, treatment of chronic stable angina pectoris, heart failure, erectile dysfunction, or ischemic stroke as there is insufficient peer-reviewed scientific literature to demonstrate that the procedure is effective.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate section of the Benefit Booklet, Evidence of Coverage or Subscriber Agreement for services not medically necessary.

BACKGROUND

Enhanced external counterpulsation (EECP) uses timed, sequential inflation of pressure cuffs on the calves, thighs, and buttocks to augment diastolic pressure, decrease left ventricular afterload, and increase venous return. The proposed mechanism of action is the augmentation of diastolic pressure by displacement of a volume of blood backward into the coronary arteries during diastole when the heart is in a state of relaxation and resistance in the coronary arteries is at a minimum. The resulting increase in coronary artery perfusion pressure may enhance coronary collateral development or increase flow through existing collaterals. Also, when the left ventricular contracts, it faces reduced aortic counterpressure, because the counterpulsation has somewhat emptied the aorta. EECP has been primarily investigated as a treatment for chronic stable angina.

Intra-aortic balloon counterpulsation is a more familiar, invasive form of counterpulsation that is used as a method of temporary circulatory assistance for the ischemic heart, often after acute myocardial infarction. In contrast, EECP is thought to provide a permanent effect on the heart by enhancing the coronary collateral development. A full course of therapy usually consists of 35, one-hour treatments, which may be offered once or twice daily, usually 5 days a week. The multiple components of the procedure include the use of the device itself, finger plethysmography to follow the blood flow, continuous electrocardiograms to trigger inflation and deflation, and optional use of pulse oximetry to measure oxygen saturation before and after treatment.

Regulatory Status

A variety of enhanced external counterpulsation (EECP) devices have been cleared for marketing by the Food and Drug Administration (FDA) through the 510(k) process. Examples of EECP devices with FDA clearance are the Renew® NCP-5 External Counterpulsation System, ECP Health System Model, CardiAssist™ Counter Pulsation System, ACS Model NCP-2 External Counterpulsation Device, and the EECP® Therapy System.

For individuals who have chronic stable angina who receive enhanced external counterpulsation (EECP), the evidence includes randomized controlled trials (RCTs), observational studies, and systematic reviews. Relevant outcomes are overall survival, symptoms, morbid events, and functional outcomes. There is a single blinded RCT that includes clinical outcomes, and it reported benefit on only 1 of 4 main angina outcomes. Additional small RCTs have reported changes in physiologic measures associated with EECP but did not provide relevant evidence on clinical efficacy. Because of the variable natural history of angina, the multiple confounding variables for cardiac outcomes, and the potential for a placebo effect, RCT evidence is needed. Therefore, Observational studies, including registry studies with large numbers of patients, add little to determinations of efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have heart failure who receive EECP, the evidence includes RCTs, observational studies, and systematic reviews. Relevant outcomes are overall survival, symptoms, morbid events, and functional outcomes. One RCT that reported on clinical outcomes found a modest benefit with EECP on some outcomes and not others. A second RCT reported improvements on the 6-minute walk test with EECP, but had methodologic limitations: RCT findings ultimately proved inconclusive. The observational studies on EECP in heart failure have limited ability to inform the evidence on EECP due to the multiple confounding variables for cardiac outcomes and the potential for a placebo effect. The evidence is insufficient to determine the effects of the technology on health outcomes. Therefore this service is considered not medically necessary for Commercial products.

BlueCHiP for Medicare

Medicare has published a national coverage decision on EECP that allows coverage for the following indications:

“Coverage is provided for the use of EECP for patients who have been diagnosed with disabling angina who, in the opinion of a cardiologist or cardiothoracic surgeon, are not readily amenable to surgical intervention, such as percutaneous transluminal coronary angioplasty or cardiac bypass because: 1) Their condition is inoperable, or at high risk of operative complications or post-operative failure; 2) Their coronary anatomy is not readily amendable to such procedures; or 3) They have co-morbid states which create excessive risk.”

CODING

BlueCHiP for Medicare and Commercial Products

The following HCPCS code is covered for BlueCHiP for Medicare and not medically necessary for Commercial products:

G0166 External counterpulsation, per treatment session

RELATED POLICIES

None

PUBLISHED

Provider Update, July 2019
Provider Update, July 2018
Provider Update, May 2017
Provider Update, August 2016
Provider Update, August 2015
Provider Update, January 2015
Provider Update, November 2013

REFERENCES

1. Center for Medicare & Medicaid Services (CMS). National Coverage Determination for external counterpulsation (ECP) therapy for severe angina (20.20). 2006; [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?CALId=185&CalName=PSA+\(Addition+of+ICD-9-CM+600.10%2C+Nodular+prostate+without+urinary+obstruction+and+600.11%2C+with+urinary+obstruction%2C+as+covered+indications\)&ExpandComments=n&CommentPeriod=0&NCDId=97&ncdver=2&CoverageSelection=National&ncd_id=20.20&ncd_version=2&basket=ncd%2525253A20%2525252E20%2525253A2%2525253AExternal+Counterpulsation+%25252528ECP%25252529+for+Severe+Angin&bc=gAAAABAAQEAAA%3D%3D&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?CALId=185&CalName=PSA+(Addition+of+ICD-9-CM+600.10%2C+Nodular+prostate+without+urinary+obstruction+and+600.11%2C+with+urinary+obstruction%2C+as+covered+indications)&ExpandComments=n&CommentPeriod=0&NCDId=97&ncdver=2&CoverageSelection=National&ncd_id=20.20&ncd_version=2&basket=ncd%2525253A20%2525252E20%2525253A2%2525253AExternal+Counterpulsation+%25252528ECP%25252529+for+Severe+Angin&bc=gAAAABAAQEAAA%3D%3D&). Accessed April 6, 2018.
2. Arora RR, Chou TM, Jain D, et al. The multicenter study of enhanced external counterpulsation (MUST-EECP): effect of EECP on exercise-induced myocardial ischemia and anginal episodes. *J Am Coll Cardiol*. Jun 1999;33(7):1833-1840. PMID 10362181
3. Arora RR, Chou TM, Jain D, et al. Effects of enhanced external counterpulsation on health-related quality of life continue 12 months after treatment: a substudy of the Multicenter Study of Enhanced External Counterpulsation. *J Investig Med*. Jan 2002;50(1):25-32. PMID 11813825
4. Gloekler S, Meier P, de Marchi SF, et al. Coronary collateral growth by external counterpulsation: a randomised controlled trial. *Heart*. Feb 2010;96(3):202-207. PMID 19897461
5. Buschmann EE, Utz W, Pagonas N, et al. Improvement of fractional flow reserve and collateral flow by treatment with external counterpulsation (Art.Net.-2 Trial). *Eur J Clin Invest*. Oct 2009;39(10):866-875. PMID 19572918
6. Braith RW, Conti CR, Nichols WW, et al. Enhanced external counterpulsation improves peripheral artery flow-mediated dilation in patients with chronic angina: a randomized sham-controlled study. *Circulation*. Oct 19 2010;122(16):1612-1620. PMID 20921442
7. Casey DP, Beck DT, Nichols WW, et al. Effects of enhanced external counterpulsation on arterial stiffness and myocardial oxygen demand in patients with chronic angina pectoris. *Am J Cardiol*. May 15 2011;107(10):1466-1472. PMID 21420062
8. Shakouri SK, Razavi Z, Eslamian F, et al. Effect of enhanced external counterpulsation and cardiac rehabilitation on quality of life, plasma nitric oxide, endothelin 1 and high sensitive CRP in patients with coronary artery disease: a pilot study. *Ann Rehabil Med*. Apr 2015;39(2):191-198. PMID 25932415
9. Holubkov R, Kennard ED, Foris JM, et al. Comparison of patients undergoing enhanced external counterpulsation and percutaneous coronary intervention for stable angina pectoris. *Am J Cardiol*. May 15 2002;89(10):1182-1186. PMID 12008172
10. Shechter M, Matetzky S, Feinberg MS, et al. External counterpulsation therapy improves endothelial function in patients with refractory angina pectoris. *J Am Coll Cardiol*. Dec 17 2003;42(12):2090-2095. PMID 14680732
11. Amin F, Al Hajeri A, Civelek B, et al. Enhanced external counterpulsation for chronic angina pectoris. *Cochrane Database Syst Rev*. Feb 17 2010(2):CD007219. PMID 20166092

12. Shah SA, Shapiro RJ, Mehta R, et al. Impact of enhanced external counterpulsation on Canadian Cardiovascular Society angina class in patients with chronic stable angina: a meta-analysis. *Pharmacotherapy*. Jul 2010;30(7):639-645. PMID 20575628
13. McKenna C, McDaid C, Suekarran S, et al. Enhanced external counterpulsation for the treatment of stable angina and heart failure: a systematic review and economic analysis. *Health Technol Assess*. Apr 2009;13(24):iii-iv, ix-xi, 1-90. PMID 19409154
14. Barsheshet A, Hod H, Shechter M, et al. The effects of external counter pulsation therapy on circulating endothelial progenitor cells in patients with angina pectoris. *Cardiology*. Dec 2008;110(3):160-166. PMID 18057883
15. Qin X, Deng Y, Wu D, et al. Does enhanced external counterpulsation (EECP) Significantly affect myocardial perfusion?: a systematic review & meta-analysis. *PLoS One*. Apr 2016;11(4):e0151822. PMID 27045935
16. Soran O, Kennard ED, Bart BA, et al. Impact of external counterpulsation treatment on emergency department visits and hospitalizations in refractory angina patients with left ventricular dysfunction. *Congest Heart Fail*. Jan-Feb 2007;13(1):36-40. PMID 17268208
17. Loh PH, Cleland JG, Louis AA, et al. Enhanced external counterpulsation in the treatment of chronic refractory angina: a long-term follow-up outcome from the International Enhanced External Counterpulsation Patient Registry. *Clin Cardiol*. Apr 2008;31(4):159-164. PMID 18404725
18. Thakkar BV, Hirsch AT, Satran D, et al. The efficacy and safety of enhanced external counterpulsation in patients with peripheral arterial disease. *Vasc Med*. Feb 2010;15(1):15-20. PMID 19841026
19. Kumar A, Aronow WS, Vadnerkar A, et al. Effect of enhanced external counterpulsation on clinical symptoms, quality of life, 6-minute walking distance, and echocardiographic measurements of left ventricular systolic and diastolic function after 35 days of treatment and at 1-year follow up in 47 patients with chronic refractory angina pectoris. *Am J Ther*. Mar-Apr 2009;16(2):116-118. PMID 19300038
20. Pettersson T, Bondesson S, Cojocaru D, et al. One year follow-up of patients with refractory angina pectoris treated with enhanced external counterpulsation. *BMC Cardiovasc Disord*. Jun 15 2006;6:28. PMID 16776842
21. Loh PH, Louis AA, Windram J, et al. The immediate and long-term outcome of enhanced external counterpulsation in treatment of chronic stable refractory angina. *J Intern Med*. Mar 2006;259(3):276-284. PMID 16476105
22. Feldman AM, Silver MA, Francis GS, et al. Treating heart failure with enhanced external counterpulsation (EECP): design of the Prospective Evaluation of EECP in Heart Failure (PEECH) trial. *J Card Fail*. Apr 2005;11(3):240-245. PMID 15812754
23. Feldman AM, Silver MA, Francis GS, et al. Enhanced external counterpulsation improves exercise tolerance in patients with chronic heart failure. *J Am Coll Cardiol*. Sep 19 2006;48(6):1198-1205. PMID 16979005
24. Feldman AM, Silver MA, Francis GS, et al. Enhanced external counterpulsation improves exercise tolerance in patients with chronic heart failure. *J Am Coll Cardiol*. Sep 19 2006;48(6):1198-1205. PMID 16979005
25. Abbottsmith CW, Chung ES, Varricchione T, et al. Enhanced external counterpulsation improves exercise duration and peak oxygen consumption in older patients with heart failure: a subgroup analysis of the PEECH trial. *Congest Heart Fail*. Nov-Dec 2006;12(6):307-311. PMID 17170583

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

