Medical Coverage Policy | Electrical Bone Growth Stimulation of the Appendicular Skeleton-Implantable and Semi-Implantable

Blue Cross Blue Shield of Rhode Island

EFFECTIVE DATE:10|01|2017 **POLICY LAST UPDATED:** 09|03|2019

OVERVIEW

In the appendicular skeleton, electrical stimulation (with either implantable electrodes or noninvasive surface stimulators) has been investigated for the treatment of delayed union, nonunion, and fresh fractures. This policy addresses only implantable and semi-invasive devices.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION Not applicable

POLICY STATEMENT

BlueCHiP for Medicare

The implantable electric bone growth stimulator is covered as the Centers for Medicare and Medicaid Services (CMS) has determined that it is medically necessary.

Note: Medicare policy is developed separately from BCBSRI policy. Medicare policy incorporates consideration of governmental regulations from CMS, such as national coverage determinations or local coverage determinations. In addition to benefit differences, CMS may reach different conclusions regarding the scientific evidence than does BCBSRI. Medicare and BCBSRI policies may differ. However, BlueCHiP for Medicare members must be offered, at least, the same services that Medicare offers.

Commercial Products

Implantable electrical bone growth stimulator is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes

BlueCHiP for Medicare and Commercial Products

The semi-invasive electrical bone growth stimulator is not covered as there are devices with U.S. Food and Drug Administration (FDA) approval or clearance.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Evidence of Coverage, Subscriber Agreement for applicable not medically necessary benefits/coverage.

BACKGROUND

In the appendicular skeleton, electrical stimulation has been primarily used to treat tibial fractures, and thus this technique has often been thought of as a treatment of the long bones. According to orthopedic anatomy, the skeleton consists of long bones, short bones, flat bones, and irregular bones. Long bones act as levels to facilitate motion, while short bones function to dissipate concussive forces. Short bones include those composing the carpus and tarsus. Flat bones, such as the scapula or pelvis, provide a broad surface area for attachment of muscles. Despite their anatomic classification, all bones are composed of a combination of cortical and trabecular (also called cancellous) bone. Each bone, depending on its physiologic function, has a different proportion of cancellous to trabecular bone. At a cellular level, however, both bone types are composed of lamellar bone and cannot be distinguished microscopically.

Electrical and electromagnetic fields can be generated and applied to bones through the following methods:

• Surgical implantation of a cathode at the fracture site with the production of direct current electrical stimulation. Invasive devices require surgical implantation of a current generator in an intramuscular or subcutaneous space, while an electrode is implanted within the fragments of bone graft at the fusion site. The implantable device typically remains functional for 6 to 9 months after implantation, and, although the current generator is removed in a second surgical procedure when stimulation is completed, the electrode may or may not be removed. Implantable electrodes provide constant stimulation at the nonunion or fracture site but carry increased risks associated with implantable leads.

• Semi-invasive (semi-implantable) stimulators use percutaneous electrodes and an external power supply obviating the need for a surgical procedure to remove the generator when treatment is finished. No semi-invasive electrical bone growth stimulator devices with FDA approval or clearance were identified.

For individuals who have fracture, pseudoarthroses, or who have had surgery of the appendicular skeleton who receive implantable and semi-invasive electrical bone growth stimulation, the evidence includes a small number of case series. Relevant outcomes are symptoms, change in disease status, and functional outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes

BlueCHiP for Medicare

CMS has a National Coverage Determination (NCD) which states that the invasive stimulator device is covered only for the following indications:

- Nonunion of long bone fractures;
- Effective July 1, 1996, as an adjunct to spinal fusion surgery for patients at high risk of pseudarthrosis due to previously failed spinal fusion at the same site or for those undergoing multiple level fusion. A multiple level fusion involves 3 or more vertebrae (e.g., L3-L5, L4-S1, etc).
- Effective September 15, 1980, nonunion of long bone fractures is considered to exist only after 6 or more months have elapsed without healing of the fracture.
- Effective April 1, 2000, nonunion of long bone fractures is considered to exist only when starting treatment with the electrical osteogenic stimulator. Serial radiographs must include a minimum of 2 sets of radiographs, each including multiple views of the fracture site, separated by a minimum of 90 days.

Based on this coverage determination, the invasive(implantable) stimulator will be covered.

CODING

The following code is covered for BlueCHiP for Medicare and not medically necessary for Commercial products:

20975 Electrical stimulation to aid bone healing; invasive (operative)

E0749 Osteogenesis stimulator, electrical, surgically implanted

Note - There is no code for the semi-invasive stimulator as there is not a FDA approved device.

RELATED POLICIES

Preauthorization via Web-Based Tool for Durable Medical Equipment (DME)

PUBLISHED

Provider Update, November 2019 Provider Update, September 2018 Provider Update, December 2017 Provider Update, January 2017 Provider Update, August 2015

REFERENCES

1.Buza JA, 3rd, Einhorn T. Bone healing in 2016. Clin Cases Miner Bone Metab. May-Aug 2016;13(2):101-105. PMID 27920804

2. Bhandari M, Fong K, Sprague S, et al. Variability in the definition and perceived causes of delayed unions and nonunions: a cross-sectional, multinational survey of orthopaedic surgeons. J Bone Joint Surg Am. Aug 12012;94(15):e1091-1096. PMID 22854998

3. Ahl T, Andersson G, Herberts P, et al. Electrical treatment of non-united fractures. Acta Orthop Scand. Dec 1984;55(6):585-588. PMID 6335345

4. Connolly JF. Selection, evaluation and indications for electrical stimulation of ununited fractures. Clin Orthop Relat Res. Nov-Dec 1981(161):39-53. PMID 6975690

5. Connolly JF. Electrical treatment of nonunions. Its use and abuse in 100 consecutive fractures. Orthop Clin North Am. Jan 1984;15(1):89-106. PMID 6607443

6. de Haas WG, Beaupre A, Cameron H, et al. The Canadian experience with pulsed magnetic fields in the treatment of ununited tibial fractures. Clin Orthop Relat Res. Jul 1986(208):55-58. PMID 3720140

7. Sharrard WJ, Sutcliffe ML, Robson MJ, et al. The treatment of fibrous non-union of fractures by pulsing electromagnetic stimulation. J Bone Joint Surg Br. 1982;64(2):189-193. PMID 6978339

8. Aleem IS, Aleem I, Evaniew N, et al. Efficacy of electrical stimulators for bone healing: a meta-analysis of randomized sham-controlled trials. Sci Rep. Aug 19 2016;6:31724. PMID 27539550

9. Simonis RB, Parnell EJ, Ray PS, et al. Electrical treatment of tibial non-union: a prospective, randomised, double-blind trial. Injury. May 2003;34(5):357-362. PMID 12719164

10. Barker AT, Dixon RA, Sharrard WJ, et al. Pulsed magnetic field therapy for tibial non-union. Interim results of a double-blind trial. Lancet. May 5 1984;1(8384):994-996. PMID 6143970

11. Scott G, King JB. A prospective, double-blind trial of electrical capacitive coupling in the treatment of non-union of long bones. J Bone Joint Surg Am. Jun 1994;76(6):820-826. PMID 8200888

12. Shi HF, Xiong J, Chen YX, et al. Early application of pulsed electromagnetic field in the treatment of postoperative delayed union of long-bone fractures: a prospective randomized controlled study. BMC Musculoskelet Disord. 2013;14:35. PMID 23331333

13. Sharrard WJ. A double-blind trial of pulsed electromagnetic fields for delayed union of tibial fractures. J Bone Joint Surg Br. May 1990;72(3):347-355. PMID 2187877

Griffin XL, Warner F, Costa M. The role of electromagnetic stimulation in the management of established nonunion of long bone fractures: what is the evidence? Injury. Apr 2008;39(4):419-429. PMID 18321512 15. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Electrical bone growth stimulation for delayed union or nonunion of fractures. TEC Assessment. 1992:Volume 7:332-351.

16. Griffin XL, Costa ML, Parsons N, et al. Electromagnetic field stimulation for treating delayed union or non-union of long bone fractures in adults. Cochrane Database Syst Rev. 2011(4):CD008471. PMID 21491410

17. Adie S, Harris IA, Naylor JM, et al. Pulsed electromagnetic field stimulation for acute tibial shaft fractures: a multicenter, double-blind, randomized trial. J Bone Joint Surg Am. Sep 7 2011;93(17):1569-1576. PMID 21915570

18. Faldini C, Cadossi M, Luciani D, et al. Electromagnetic bone growth stimulation in patients with femoral neck fractures treated with screws: prospective randomized double-blind study. Curr Orthop Pract. 2010;21(3):282-287.

19. Hannemann PF, Gottgens KW, van Wely BJ, et al. The clinical and radiological outcome of pulsed electromagnetic field treatment for acute scaphoid fractures: a randomised double-blind placebo-controlled multicentre trial. J Bone Joint Surg Br. Oct 2012;94(10):1403-1408. PMID 23015569 20. Hannemann PF, van Wezenbeek MR, Kolkman KA, et al. CT scan-evaluated outcome of pulsed electromagnetic fields in the treatment of acute scaphoid fractures: a randomised, multicentre, double-blind, placebo-controlled trial. Bone Joint J. Aug 2014;96-B(8):1070-1076. PMID 25086123

21. Martinez-Rondanelli A, Martinez JP, Moncada ME, et al. Electromagnetic stimulation as coadjuvant in the healing of diaphyseal femoral fractures: a randomized controlled trial. Colomb Med (Cali). Apr-Jun

2014;45(2):67-71. PMID 25100891 22. Hannemann PF, Essers BA, Schots JP, et al. Functional outcome and cost-effectiveness of pulsed electromagnetic fields in the treatment of acute scaphoid fractures: a cost-utility analysis. BMC Musculoskelet Disord. Apr 11 2015;16:84. PMID 25880388

23. Beck BR, Matheson GO, Bergman G, et al. Do capacitively coupled electric fields accelerate tibial stress fracture healing? A randomized controlled trial. Am J Sports Med. Mar 2008;36(3):545-553. PMID 18055921
24. Borsalino G, Bagnacani M, Bettati E, et al. Electrical stimulation of human femoral intertrochanteric osteotomies. Double-blind study. Clin Orthop Relat Res. Dec 1988(237):256-263. PMID 3191636
25. Dhawan SK, Conti SF, Towers J, et al. The effect of pulsed electromagnetic fields on hindfoot arthrodesis: a prospective study. J Foot Ankle Surg. Mar-Apr 2004;43(2):93-96. PMID 15057855
26. Petrisor B, Lau JT. Electrical bone stimulation: an overview and its use in high risk and Charcot foot and

ankle reconstructions. Foot Ankle Clin. Dec 2005;10(4):609-620, vii-viii. PMID 16297822

27. Lau JT, Stamatis ED, Myerson MS, et al. Implantable direct-current bone stimulators in high-risk and revision foot and ankle surgery: a retrospective analysis with outcome assessment. Am J Orthop (Belle Mead NJ). Jul 2007;36(7):354-357. PMID 17694182

28. Saxena A, DiDomenico LA, Widtfeldt A, et al. Implantable electrical bone stimulation for arthrodeses of the foot and ankle in high-risk patients: a multicenter study. J Foot Ankle Surg. Nov-Dec 2005;44(6):450-454. PMID 16257674

29. Centers for Medicare and Medicaid Services. National Coverage Determination for Osteogenic Stimulators (150.2). 2005; https://www.cms.gov/medicare-coverage-database/search/document-id-searchresults.

aspx?DocID=150.2&ncd_id=150.2&ncd_version=2&basket=ncd%25253A150%25252E2%25253A2%252 53AOsteogenic+Stimulators&bc=gAAAAAAAAAAAAA. Accessed August, 2011.

----- CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield Association.

