

EFFECTIVE DATE: 01|01|2017

POLICY LAST UPDATED: 09|17|2019

OVERVIEW

Chelation therapy, an established treatment for heavy metal toxicities and transfusional hemosiderosis, has been investigated for a variety of off-label applications, such as treatment of atherosclerosis, Alzheimer disease, and autism. This policy addresses the following off-label uses of chelation therapy:

- Alzheimer disease
- Atherosclerotic cardiovascular disease
- Arthritis, including rheumatoid arthritis
- Autism spectrum disorder
- Diabetes
- Multiple sclerosis

This policy does not address the following U.S Food and Drug Administration (FDA)-approved indications for which chelation therapy is considered standard of care treatment:

- Extreme conditions of metal toxicity
- Treatment of chronic iron overload due to blood transfusions (transfusional hemosiderosis) or due to non-transfusion-dependent thalassemia (NTDT)
- Wilson disease (hepatolenticular degeneration)
- Lead poisoning
- Control of ventricular arrhythmias or heart block associated with digitalis toxicity
- Emergency treatment of hypercalcemia

This policy is applicable to Commercial Products only. For BlueCHiP for Medicare, see related policy section.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Commercial Products

Off-label applications of chelation therapy (non-FDA-approved uses) are considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes, including, but no limited to:

- Alzheimer disease
- Arthritis (includes rheumatoid arthritis)
- Atherosclerosis (eg, coronary artery disease, secondary prevention in patients with myocardial infarction, or peripheral vascular disease)
- Autism
- Diabetes
- Multiple sclerosis

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable not medically necessary benefits/coverage.

BACKGROUND

Chelation therapy is an established treatment for the removal of metal toxins by converting them to a chemically inert form that can be excreted in the urine. Chelation therapy comprises intravenous or oral administration of chelating agents that remove metal ions such as lead, aluminum, mercury, arsenic, zinc, iron, copper, and calcium from the body. Specific chelating agents are used for particular heavy metal toxicities. For example, desferrioxamine (not FDA-approved) is used for patients with iron toxicity, and calcium-ethylenediaminetetraacetic acid (EDTA) is used for patients with lead poisoning. (Disodium-EDTA is not recommended for acute lead poisoning due to the increased risk of death from hypocalcemia.)

Another class of chelating agents, called metal protein attenuating compounds (MPACs), is under investigation for the treatment of Alzheimer disease, which is associated with the disequilibrium of cerebral metals. Unlike traditional systemic chelators that bind and remove metals from tissues systemically, MPACs have subtle effects on metal homeostasis and abnormal metal interactions. In animal models of Alzheimer disease, they promote the solubilization and clearance of beta amyloid by binding its metal-ion complex, and also inhibit redox reactions that generate neurotoxic free radicals. MPACs therefore interrupt 2 putative pathogenic processes of Alzheimer disease. However, no MPACs have received FDA approval for the treatment of Alzheimer disease.

Chelation therapy also has been discussed as a treatment for other indications including atherosclerosis and autism spectrum disorder. For example, EDTA chelation therapy has been proposed in patients with atherosclerosis as a method of decreasing obstruction in the arteries.

In 1953, EDTA (Versenate) was approved by the FDA for lowering blood lead levels among both pediatric and adult patients with lead poisoning. In 1991, succimer (Chemet) was approved by FDA for the treatment of lead poisoning in pediatric patients only. The FDA approved disodium-EDTA for use in selected patients with hypercalcemia and use in patients with heart rhythm problems due to intoxication with digitalis. In 2008, FDA withdrew approval of disodium-EDTA due to safety concerns and recommended that other forms of chelation therapy be used.

Several iron chelating agents are FDA-approved:

- In 1968, deferoxamine (Desferal®; Novartis) was approved by FDA for subcutaneous, intramuscular, or intravenous injections to treat acute iron intoxication and chronic iron overload due to transfusion-dependent anemia. Several generic forms of deferoxamine have been approved by FDA.
- In 2005, deferasirox (Exjade®; Novartis) was approved by FDA, is available as a tablet for oral suspension, and is indicated for the treatment of chronic iron overload due to blood transfusions in patients ages 2 years and older. Under the accelerated approval program, FDA expanded the indications for deferasirox in 2013 to include treatment of patients age 10 years and older with chronic iron overload due to non-transfusion-dependent thalassemia syndromes and specific liver iron concentration and serum ferritin levels. A generic version of deferasirox tablet for oral suspension has also been approved by FDA. In 2015, an oral tablet formulation for deferasirox (Jadenu™) was approved by FDA. All formulations of deferasirox carry a black box warning because it may cause serious and fatal renal toxicity and failure, hepatic toxicity and failure, and gastrointestinal hemorrhage. As a result, treatment with deferasirox requires close patient monitoring, including laboratory tests of renal and hepatic function.
- In 2011, the iron chelator deferiprone (Ferriprox®) was approved by FDA for treatment of patients with transfusional overload due to thalassemia syndromes when another chelation therapy is

inadequate. Deferiprone is available in tablet and oral solution. Ferriprox® carries a black box warning because it can cause agranulocytosis, which can lead to serious infections and death. As a result, absolute neutrophil count should be monitored before and during treatment.

In a June 2014 warning to consumers, FDA advised that FDA-approved chelating agents would be available by prescription only. There are no FDA-approved over-the-counter chelation products.

For individuals who have Alzheimer disease, cardiovascular disease, autism spectrum disorder, diabetes, multiple sclerosis, or arthritis who receive chelation therapy, the evidence is insufficient to determine the effect of the technology on health outcomes. Thus, chelation therapy for these off-label applications is considered not medically necessary.

CODING

Commercial Products

The following code represents the infusion service only and is not separately reimbursed:

S9355 Home Infusion Therapy, chelation therapy; administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem.

Chemical Endarterectomy

The following code and any of the medications utilized as part of the service are not medically necessary when filed with the ICD-10 diagnosis codes below:

M0300 IV chelation therapy (chemical endarterectomy)

E08.00-E13.9 Diabetes mellitus code range

F84.0-F84.9 Autism spectrum disorders

G30.0-G30.9 Alzheimer's disease code range

G35 Multiple sclerosis

I25.10-I25.9 Atherosclerosis code range

M05.00-M06.09 Rheumatoid arthritis code range

M15.0-M19.93 Osteoarthritis code range

Failure of participating providers to report Chemical Endarterectomy using M0300 will be considered improper coding by BCBSRI.

RELATED POLICIES

BlueCHiP for Medicare National and Local Coverage Determinations Policy

Non Reimbursable Health Service Codes

PUBLISHED

Provider Update, November 2019

Provider Update, November 2018

Provider Update, February 2018

Provider Update, January 2017

Provider Update, August 2015

REFERENCES

1. Centers for Disease Control and Prevention. Deaths associated with hypocalcemia from chelation therapy--Texas, Pennsylvania, and Oregon, 2003-2005. *MMWR Morb Mortal Wkly Rep*. Mar 03 2006;55(8):204-207. PMID 16511441
2. Food and Drug Administration. Hospira, Inc., et al.; Withdrawal of Approval of One New Drug Application and Two Abbreviated New Drug Application. *Federal Register*. 2008;73(113):33440-33441. PMID

3. U.S Food and Drug Administration. FDA warns consumers about potential health risks from using Thorne Research's Captomer products. 2014 June 12; http://www.fda.gov/Drugs/DrugSafety/ucm400977.htm?source=govdelivery&utm_medium=email&utm_source=govdelivery. Accessed January 23, 2018.
4. Sampson E, Jenagaratnam L, McShane R. Metal protein attenuating compounds for the treatment of Alzheimer's disease. *Cochrane Database Syst Rev*. Jan 23 2008(1):CD005380. PMID 18254079
5. Ritchie CW, Bush AI, Mackinnon A, et al. Metal-protein attenuation with iodochlorhydroxyquin (clioquinol) targeting Abeta amyloid deposition and toxicity in Alzheimer disease: a pilot phase 2 clinical trial. *Arch Neurol*. Dec 2003;60(12):1685-1691. PMID 14676042
6. Sampson EL, Jenagaratnam L, McShane R. Metal protein attenuating compounds for the treatment of Alzheimer's dementia. *Cochrane Database Syst Rev*. May 16 2012;5(5):CD005380. PMID 22592705
7. Lannfelt L, Blennow K, Zetterberg H, et al. Safety, efficacy, and biomarker findings of PBT2 in targeting Abeta as a modifying therapy for Alzheimer's disease: a phase IIa, double-blind, randomised, placebo-controlled trial. *Lancet Neurol*. Sep 2008;7(9):779-786. PMID 18672400
8. Villarruz MV, Dans A, Tan F. Chelation therapy for atherosclerotic cardiovascular disease. *Cochrane Database Syst Rev*. Jan 2002(4):CD002785. PMID 12519577
9. Knudtson ML, Wyse DG, Galbraith PD, et al. Chelation therapy for ischemic heart disease: a randomized controlled trial. *Jama*. Jan 23-30 2002;287(4):481-486. PMID 11798370
10. Anderson TJ, Hubacek J, Wyse DG, et al. Effect of chelation therapy on endothelial function in patients with coronary artery disease: PATCH substudy. *J Am Coll Cardiol*. Feb 5 2003;41(3):420-425. PMID 12575969
11. Guldager B, Jelnes R, Jorgensen SJ, et al. EDTA treatment of intermittent claudication--a double-blind, placebocontrolled study. *J Intern Med*. Mar 1992;231(3):261-267. PMID 1556523
12. van Rij AM, Solomon C, Packer SG, et al. Chelation therapy for intermittent claudication. A double-blind, randomized, controlled trial. *Circulation*. Sep 1994;90(3):1194-1199. PMID 8087928
13. Lamas GA, Goertz C, Boineau R, et al. Effect of disodium EDTA chelation regimen on cardiovascular events in patients with previous myocardial infarction: the TACT randomized trial. *Jama*. Mar 27 2013;309(12):1241-1250. PMID 23532240
14. Mark DB, Anstrom KJ, Clapp-Channing NE, et al. Quality-of-life outcomes with a disodium EDTA chelation regimen for coronary disease: results from the trial to assess chelation therapy randomized trial. *Circ Cardiovasc Qual Outcomes*. Jul 2014;7(4):508-516. PMID 24987051
15. Lamas GA, Boineau R, Goertz C, et al. EDTA chelation therapy alone and in combination with oral high-dose multivitamins and minerals for coronary disease: The factorial group results of the Trial to Assess Chelation Therapy. *Am Heart J*. Jul 2014;168(1):37-44 e35. PMID 24952858
16. Nissen SE. Concerns about reliability in the Trial to Assess Chelation Therapy (TACT) [editorial]. *Jama*. Mar 27 2013;309(12):1293-1294. PMID 23532246
17. Maron DJ, Hlatky MA. Trial to Assess Chelation Therapy (TACT) and equipoise: When evidence conflicts with beliefs [editorial]. *Am Heart J*. Jul 2014;168(1):4-5. PMID 24952853
18. Bernard S, Enayati A, Redwood L, et al. Autism: a novel form of mercury poisoning. *Med Hypotheses*. Apr 2001;56(4):462-471. PMID 11339848
19. Nelson KB, Bauman ML. Thimerosal and autism? *Pediatrics*. Mar 2003;111(3):674-679. PMID 12612255
20. Ng DK, Chan CH, Soo MT, et al. Low-level chronic mercury exposure in children and adolescents: metaanalysis. *Pediatr Int*. Feb 2007;49(1):80-87. PMID 17250511
21. Rossignol DA. Novel and emerging treatments for autism spectrum disorders: a systematic review. *Ann Clin Psychiatry*. Oct-Dec 2009;21(4):213-236. PMID 19917212
22. Cooper GJ, Young AA, Gamble GD, et al. A copper(II)-selective chelator ameliorates left-ventricular hypertrophy in type 2 diabetic patients: a randomised placebo-controlled study. *Diabetologia*. Apr 2009;52(4):715-722. PMID 19172243

23. Escolar E, Lamas GA, Mark DB, et al. The effect of an EDTA-based chelation regimen on patients with diabetes mellitus and prior myocardial infarction in the Trial to Assess Chelation Therapy (TACT). *Circ Cardiovasc Qual Outcomes*. Jan 2014;7(1):15-24. PMID 24254885
24. Chen KH, Lin JL, Lin-Tan DT, et al. Effect of chelation therapy on progressive diabetic nephropathy in patients with type 2 diabetes and high-normal body lead burdens. *Am J Kidney Dis*. Oct 2012;60(4):530-538. PMID 22721929
25. U.S. Department of Labor, Occupational Health and Safety Administration. Safety and Health Regulations for Construction: Substance Data Sheet for Occupational Exposure to Lead. 1993; http://www.osha.gov/pls/oshaweb/owadispl.show_document?p_table=STANDARDS&p_id=1064
2. Accessed January 23, 2018.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

