

**EFFECTIVE DATE:** 01|01|2016

**POLICY LAST UPDATED:** 02|06|2020

## OVERVIEW

Numerous lipid and nonlipid biomarkers have been proposed as potential risk markers for cardiovascular disease. Biomarkers assessed here are those that have the most evidence in support of their use in clinical care, including apolipoprotein B (apo B), apolipoprotein AI (apo AI), apolipoprotein E (apo E), high-density lipoprotein (HDL) subclass, low-density lipoprotein (LDL) subclass, lipoprotein (a), B-type natriuretic peptide, cystatin C, fibrinogen, and leptin. These biomarkers have been studied as alternatives or additions to standard lipid panels for risk stratification in cardiovascular disease or as treatment targets for lipid-lowering therapy.

For coverage of tests filed with PLA codes (0052U-VAP Cholesterol Test), please refer to the related policy “Proprietary Laboratory Analyses (PLA).”

## MEDICAL CRITERIA

Not applicable

## PRIOR AUTHORIZATION

Not applicable

## POLICY STATEMENT

### Commercial Products

Measurement of novel lipid and non-lipid risk factors (ie, apolipoprotein B, apolipoprotein AI, apolipoprotein E, cystatin C, fibrinogen, leptin, low-density lipoprotein (LDL) subclass, high-density lipoprotein (HDL) subclass, lipoprotein[a]) are not medically necessary for Commercial products as the evidence is insufficient to determine the effects of the technology on health outcomes. This policy is applicable to Commercial Products only. For BlueCHiP for Medicare, see the related policy for BlueCHiP for Medicare National and Local Coverage Determinations.

### BlueCHiP for Medicare and Commercial Products

B type natriuretic peptide testing is covered but not separately reimbursed when used in conjunction with standard diagnostic tests, medical history and clinical findings during an evaluation of heart failure in an acute care setting or other setting (i.e. emergency department) where test results are immediately determined.

## COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate section of the Benefit Booklet, Evidence of Coverage or Subscriber Agreement for services not medically necessary.

## BACKGROUND

### Commercial Products

Low-density lipoproteins (LDLs) have been identified as the major atherogenic lipoproteins and has long been identified by the National Cholesterol Education Project (NCEP) as the primary target of cholesterol-lowering therapy. LDL particles consist of a surface coat composed of phospholipids, free cholesterol, and apolipoproteins surrounding an inner lipid core composed of cholesterol ester and triglycerides. Traditional lipid risk factors such as LDL-cholesterol (LDL-C), while predictive on a population basis, are weaker markers of risk on an individual basis. Only a minority of subjects with elevated LDL and cholesterol levels will develop

clinical disease, and up to 50% of cases of coronary artery disease (CAD) occur in subjects with ‘normal’ levels of total and LDL-C. Thus, there is considerable potential to improve the accuracy of current cardiovascular risk prediction models. Other nonlipid markers have been identified as having an association with cardiovascular disease including B-type natriuretic peptide, cystatin C, fibrinogen, and leptin. These biomarkers may have a predictive role in identifying cardiovascular disease risk or in targeting for therapy.

## **Lipid Markers**

### **Apolipoprotein B**

Apolipoprotein B (apo B) is the major protein moiety of all lipoproteins except for HDL. The most abundant form of apo B, large B or B100, constitutes the apo B found in LDL and very-low-density lipoproteins (VLDL). Because both LDL and VLDL each contain 1 molecule of apo B, measurement of apo B reflects the total number of these atherogenic particles, 90% of which are LDL. Because LDL particles can vary both in size and in cholesterol content, for a given concentration of LDL-C, there can be a wide variety of both size and numbers of LDL particles. Thus, it has been postulated that apo B is a better measure of the atherogenic potential of serum LDL than is LDL concentration.

### **Apolipoprotein AI**

HDL contains 2 associated apolipoproteins, (ie, AI and AII). HDL particles can also be classified by whether they contain apolipoprotein AI (apo AI) only or whether they contain both apo AI and apolipoprotein AII (apo AII). All lipoproteins contain apo AI, and some also contain apo AII. Because all HDL particles contain apo AI, this lipid marker can be used as an approximation for HDL number, similar to the way apo B has been proposed as an approximation of the LDL number. Direct measurement of apo AI has been proposed as more accurate than the traditional use of HDL level in evaluation of the cardioprotective, or “good,” cholesterol. In addition, the ratio of apo B/apo AI has been proposed as a superior measure of the ratio of proatherogenic (ie, “bad”) cholesterol to anti-atherogenic (ie, “good”) cholesterol.

### **Apolipoprotein E**

Apolipoprotein E (apo E) is the primary apolipoprotein found in VLDLs and chylomicrons. Apo E is the primary binding protein for LDL receptors in the liver and is thought to play an important role in lipid metabolism. The apolipoprotein E (*APOE*) gene is polymorphic, consisting of 3 epsilon alleles (e2, e3, e4) that code for 3 protein isoforms, known as E2, E3, and E4, which differ from one another by 1 amino acid. These molecules mediate lipid metabolism through their different interactions with the LDL receptors. The genotype of apo E alleles can be assessed by gene amplification techniques, or the *APOE* phenotype can be assessed by measuring plasma levels of *APOE*. It has been proposed that various *APOE* genotypes are more atherogenic than others and that *APOE* measurement may provide information on risk of CAD above traditional risk factor measurement. It has also been proposed that the *APOE* genotype may be useful in the selection of specific components of lipid-lowering therapy, such as drug selection. In the major lipid-lowering intervention trials, including trials of statin therapy, there is considerable variability in response to therapy that cannot be explained by factors such as compliance. *APOE* genotype may be a factor that determines an individual’s degree of response to interventions such as statin therapy.

### **HDL Subclass**

HDL particles exhibit considerable heterogeneity, and it has been proposed that various subclasses of HDL may have a greater role in protection from atherosclerosis. Particles of HDL can be characterized based on size or density and/or on apolipoprotein composition. Using size or density, HDL can be classified into HDL2, the larger, less dense particles that may have the greatest degree of cardioprotection, and HDL3, which are smaller, denser particles.

An alternative to measuring the concentration of subclasses of HDL (eg, HDL2, HDL3) is direct measurement of HDL particle size and/or number. Particle size can be measured by nuclear magnetic resonance spectroscopy or by gradient-gel electrophoresis. HDL particle numbers can be measured by nuclear magnetic resonance spectroscopy. Several commercial labs offer these measurements of HDL particle

size and number. Measurement of apo AI has used HDL particle number as a surrogate, based on the premise that each HDL particle contains a single apo AI molecule.

### **LDL Subclass**

Two main subclass patterns of LDL, called A and B, have been described. In subclass pattern A, particles have a diameter larger than 25 nm and are less dense, while in subclass pattern B, particles have a diameter less than 25 nm and a higher density. Subclass pattern B is a commonly inherited disorder associated with a more atherogenic lipoprotein profile, also termed “atherogenic dyslipidemia.” In addition to small, dense LDL, this pattern includes elevated levels of triglycerides, elevated levels of apo B, and low levels of HDL. This lipid profile is commonly seen in type 2 diabetes and is a component of the “metabolic syndrome,” defined by the Third Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III [ATP III]) to also include high normal blood pressure, insulin resistance, increased levels of inflammatory markers such as C-reactive protein, and a prothrombotic state. Presence of the metabolic syndrome is considered by ATP III to be a substantial risk-enhancing factor for CAD.

LDL size has also been proposed as a potentially useful measure of treatment response. Lipid-lowering treatment decreases total LDL and may also induce a shift in the type of LDL, from smaller, dense particles to larger particles. It has been proposed that this shift in lipid profile may be beneficial in reducing risk for CAD independent of the total LDL level. Also, some drugs may cause a greater shift in lipid profile than others. Niacin and/or fibrates may cause a greater shift from small to large LDL size than statins. Therefore, measurement of LDL size may potentially play a role in drug selection or may be useful in deciding whether to use a combination of drugs rather than a statin alone.

In addition to the size of LDL particles, interest has been shown in assessing the concentration of LDL particles as a distinct cardiac risk factor. For example, the commonly performed test for LDL-C is not a direct measure of LDL, but, chosen for its convenience, measures the amount of cholesterol incorporated into LDL particles. Because LDL particles carry much of the cholesterol in the bloodstream, the concentration of cholesterol in LDL correlates reasonably well with the number of LDL particles when examined in large populations. However, for an individual patient, the LDL-C level may not reflect the number of particles due to varying levels of cholesterol in different sized particles. It is proposed that the discrepancy between the number of LDL particles and the serum level of LDL-C represents a significant source of unrecognized atherogenic risk. The size and number of particles are interrelated. For example, all LDL particles can invade the arterial wall and initiate atherosclerosis. However, small, dense particles are thought to be more atherogenic than larger particles. Therefore, for patients with elevated numbers of LDL particles, cardiac risk may be further enhanced when the particles are smaller vs larger.

### **Lipoprotein (a)**

Lipoprotein (a) (Lp[a]) is a lipid-rich particle similar to LDL. Apo B is the major apolipoprotein associated with LDL; in Lp(a), however, there is an additional apo A covalently linked to the apo B. The apo A molecule is structurally similar to plasminogen, suggesting that Lp(a) may contribute to the thrombotic and atherogenic basis of CVD. Levels of Lp(a) are relatively stable in individuals over time but vary up to 1000-fold between individuals, presumably on a genetic basis. The similarity between Lp(a) and fibrinogen has stimulated intense interest in Lp(a) as a link between atherosclerosis and thrombosis. In addition, approximately 20% of patients with CAD have elevated Lp(a) levels. Therefore, it has been proposed that levels of Lp(a) may be an independent risk factor for CAD.

### **Non-Lipid Markers**

#### **B-Type or Brain Natriuretic Peptide**

Brain natriuretic peptide (BNP) is an amino acid polypeptide that is secreted primarily by the ventricles of the heart when pressure to the cardiac muscles increases or there is myocardial ischemia. Elevations in BNP levels reflect deterioration in cardiac loading levels and may predict

adverse events. BNP has been studied as a biomarker for managing heart failure and predicting cardiovascular and heart failure risk.

### **Cystatin C**

Cystatin C is a small serine protease inhibitor protein secreted from all functional cells in the body. It has primarily been used as a biomarker of kidney function. Cystatin C has also been studied to determine whether it may serve as a biomarker for predicting cardiovascular risk. Cystatin C is encoded by the *CST3* gene.

### **Fibrinogen**

Fibrinogen is a circulating clotting factor and precursor of fibrin. It is important in platelet aggregation and a determinant of blood viscosity. Fibrinogen levels have been shown to be associated with future risk of CVD and all-cause mortality.

### **Leptin**

Leptin is a protein secreted by fat cells that has been found to be elevated in heart disease. Leptin has been studied to determine if it has any relationship with the development of cardiovascular disease.

For individuals who are asymptomatic with risk of CVD who receive novel cardiac biomarker testing (eg, apo B, apo AI, apo E, HDL subclass, LDL subclass, Lp[a], BNP, cystatin C, fibrinogen, leptin), the evidence includes systematic reviews, meta-analyses, and large, prospective cohort studies. Relevant outcomes are overall survival, other test performance measures, change in disease status, morbid events, and medication use. The evidence from cohort studies and meta-analyses of these studies has suggested that some of these markers are associated with increased cardiovascular risk and may provide incremental accuracy in risk prediction. In particular, apo B and apo AI have been identified as adding some incremental predictive value. However, it has not been established whether the incremental accuracy provides clinically important information beyond that of traditional lipid measures. Furthermore, no study has provided high-quality evidence that measurement of markers leads to changes in management that improve health outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with hyperlipidemia managed with lipid-lowering therapy who receive novel cardiac biomarker testing (eg, apo B, apo AI, apo E, apo E, HDL subclass, LDL subclass, lipoprotein [a], BNP, cystatin C, fibrinogen, leptin), the evidence includes analyses of the intervention arm(s) of lipid-lowering medication trials. Relevant outcomes are overall survival, change in disease status, morbid events, and medication use. In particular, apo B, apo AI, and apo E have been evaluated as markers of lipid-lowering treatment success, and evidence from the intervention arms of several randomized controlled trials has suggested that these markers are associated with treatment success. However, there is no direct evidence that using markers other than LDL and HDL as a lipid-lowering treatment target leads to improved health outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes. Therefore, these services are considered not medically necessary for Commercial products.

There is a Local Coverage Determination (LCD) for B-type Natriuretic Peptide (BNP) Testing that indicates: BNP measurements may be considered reasonable and necessary when used in combination with other medical data such as medical history, physical examination, laboratory studies, chest x-ray, and electrocardiography:

- To distinguish cardiac cause of acute dyspnea from pulmonary or other non-cardiac causes. Plasma BNP levels are significantly increased in patients with CHF presenting with acute dyspnea compared with patients presenting with acute dyspnea due to other causes.
- To distinguish decompensated CHF from exacerbated chronic obstructive pulmonary disease (COPD) in a symptomatic patient with combined chronic CHF and COPD. Plasma BNP levels are significantly increased in patients with CHF with or without concurrent lung disease compared with patients who have primary lung disease.
- To establish prognosis or disease severity in chronic CHF when needed to guide therapy

- To achieve optimal dosing of guideline-directed medical therapy (GDMT) in select clinically euvolemic patients followed in a well-structured heart failure (HF) disease management program
- To guide therapeutic decision-making in individuals who have amyloidosis

BNP measurements must be analyzed in conjunction with standard diagnostic tests, medical history and clinical findings. The efficacy of BNP measurement as a stand-alone test has not yet been established. Clinicians should be aware that certain conditions such as ischemia, infarction and renal insufficiency, may cause elevation of circulating BNP concentration and require alterations of the interpretation of BNP results. Therefore, B type natriuretic peptide testing is covered but not separately reimbursed when used in conjunction with standard diagnostic tests, medical history and clinical findings during an evaluation of heart failure in an acute care setting or other setting (i.e. emergency department) where test results are immediately determined.

## **CODING**

### **Commercial Products**

The following CPT codes are not medically necessary:

- 82610** Cystatin C
- 83695** Lipoprotein (a)
- 83700** Lipoprotein, blood; electrophoretic separation and quantitation
- 83701** Lipoprotein, blood; high resolution fractionation and quantitation of lipoproteins including lipoprotein subclasses when performed (eg, electrophoresis, ultracentrifugation)
- 83704** Lipoprotein, blood; quantitation of lipoprotein particle number(s) (eg, by nuclear magnetic resonance spectroscopy), includes lipoprotein particle subclass(es), when performed
- 83722** Lipoprotein, Direct Measurement; Small Dense LDL Cholesterol (Effective 1/1/2019)
- 85384** Fibrinogen; activity
- 85385** Fibrinogen; antigen

### **BlueCHiP for Medicare and Commercial Products:**

The following CPT code is covered but not separately reimbursed for BlueCHiP for Medicare and Commercial Products:

- 83880** Natriuretic peptide

## **RELATED POLICIES**

BlueCHiP for Medicare National and Local Coverage Determinations Policy  
 Measurement of Small Low-Density Lipoprotein (LDL) Particles  
 Propriety Laboratory Analyses (PLA)

## **PUBLISHED**

Provider Update, April 2020  
 Provider Update, July 2019  
 Provider Update, October 2018  
 Provider Update, July 2017  
 Provider Update, September, 2016  
 Provider Update, November, 2015

## **REFERENCES:**

1. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*. May 16 2001;285(19):2486-2497. PMID 11368702
2. BlueCross and BlueShield Technology Evaluation Center. C-Reactive Protein as a Cardiac Risk Marker (Special Report). *TEC Assessment*. 2002;Volume 17:Tab 23.



3. Perera R, McFadden E, McLellan J, et al. Optimal strategies for monitoring lipid levels in patients at risk or with cardiovascular disease: a systematic review with statistical and cost-effectiveness modelling. *Health Technol Assess.* Dec 2015;19(100):1-401, vii-viii. PMID 26680162
4. Thanassoulis G, Williams K, Ye K, et al. Relations of change in plasma levels of LDL-C, non-HDL-C and apoB with risk reduction from statin therapy: a meta-analysis of randomized trials. *J Am Heart Assoc.* Apr 14 2014;3(2):e000759. PMID 24732920
5. van Holten TC, Waanders LF, de Groot PG, et al. Circulating biomarkers for predicting cardiovascular disease risk; a systematic review and comprehensive overview of meta-analyses. *PLoS One.* May 2013;8(4):e62080. PMID 23630624
6. Tzoulaki I, Siontis KC, Evangelou E, et al. Bias in associations of emerging biomarkers with cardiovascular disease. *JAMA Intern Med.* Apr 22 2013;173(8):664-671. PMID 23529078
7. Willis A, Davies M, Yates T, et al. Primary prevention of cardiovascular disease using validated risk scores: a systematic review. *J R Soc Med.* Aug 2012;105(8):348-356. PMID 22907552
8. Robinson JG, Wang S, Jacobson TA. Meta-analysis of comparison of effectiveness of lowering apolipoprotein B versus low-density lipoprotein cholesterol and nonhigh-density lipoprotein cholesterol for cardiovascular risk reduction in randomized trials. *Am J Cardiol.* Nov 15 2012;110(10):1468-1476. PMID 22906895
9. Sniderman AD, Islam S, Yusuf S, et al. Discordance analysis of apolipoprotein B and non-high density lipoprotein cholesterol as markers of cardiovascular risk in the INTERHEART study. *Atherosclerosis.* Dec 2012;225(2):444-449. PMID 23068583
10. Emerging Risk Factors Collaboration, Di Angelantonio E, Gao P, et al. Lipid-related markers and cardiovascular disease prediction. *JAMA.* Jun 20 2012;307(23):2499-2506. PMID 22797450
11. Lamarche B, Moorjani S, Lupien PJ, et al. Apolipoprotein A-I and B levels and the risk of ischemic heart disease during a five-year follow-up of men in the Quebec Cardiovascular Study. *Circulation.* Aug 1 1996;94(3):273-278. PMID 8759066
12. Walldius G, Jungner I, Holme I, et al. High apolipoprotein B, low apolipoprotein A-I, and improvement in the prediction of fatal myocardial infarction (AMORIS study): a prospective study. *Lancet.* Dec 15 2001;358(9298):2026-2033. PMID 11755609
13. Ridker PM, Rifai N, Cook NR, et al. Non-HDL cholesterol, apolipoproteins A-I and B100, standard lipid measures, lipid ratios, and CRP as risk factors for cardiovascular disease in women. *JAMA.* Jul 20 2005;294(3):326-333. PMID 16030277
14. Benn M, Nordestgaard BG, Jensen GB, et al. Improving prediction of ischemic cardiovascular disease in the general population using apolipoprotein B: the Copenhagen City Heart Study. *Arterioscler Thromb Vasc Biol.* Mar 2007;27(3):661-670. PMID 17170368
15. Sharrett AR, Ballantyne CM, Coady SA, et al. Coronary heart disease prediction from lipoprotein cholesterol levels, triglycerides, lipoprotein(a), apolipoproteins A-I and B, and HDL density subfractions: The Atherosclerosis Risk in Communities (ARIC) Study. *Circulation.* Sep 4 2001;104(10):1108-1113. PMID 11535564
16. Rasouli M, Kiasari AM, Mokhberi V. The ratio of apoB/apoAI, apoB and lipoprotein(a) are the best predictors of stable coronary artery disease. *Clin Chem Lab Med.* Aug 2006;44(8):1015-1021. PMID 16879071
17. Walldius G, Jungner I. Apolipoprotein B and apolipoprotein A-I: risk indicators of coronary heart disease and targets for lipid-modifying therapy. *J Intern Med.* Feb 2004;255(2):188-205. PMID 14746556
18. Kappelle PJ, Gansevoort RT, Hillege JL, et al. Apolipoprotein B/A-I and total cholesterol/high-density lipoprotein cholesterol ratios both predict cardiovascular events in the general population independently of nonlipid risk factors, albuminuria and C-reactive protein. *J Intern Med.* Feb 2011;269(2):232-242. PMID 21129046
19. Ridker PM, Buring JE, Rifai N, et al. Development and validation of improved algorithms for the assessment of global cardiovascular risk in women: the Reynolds Risk Score. *JAMA.* Feb 14 2007;297(6):611-619. PMID 17299196
20. Ingelsson E, Schaefer EJ, Contois JH, et al. Clinical utility of different lipid measures for prediction of coronary heart disease in men and women. *JAMA.* Aug 15 2007;298(7):776-785. PMID 17699011

21. Pencina MJ, D'Agostino RB, Zdrojewski T, et al. Apolipoprotein B improves risk assessment of future coronary heart disease in the Framingham Heart Study beyond LDL-C and non-HDL-C. *Eur J Prev Cardiol.* Oct 2015;22(10):1321-1327. PMID 25633587
22. Campos H, Moye LA, Glasser SP, et al. Low-density lipoprotein size, pravastatin treatment, and coronary events. *JAMA.* Sep 26 2001;286(12):1468-1474. PMID 11572739
23. Ridker PM, Hennekens CH, Stampfer MJ. A prospective study of lipoprotein(a) and the risk of myocardial infarction. *JAMA.* Nov 10 1993;270(18):2195-2199. PMID 8411602
24. Suk Danik J, Rifai N, Buring JE, et al. Lipoprotein(a), hormone replacement therapy, and risk of future cardiovascular events. *J Am Coll Cardiol.* Jul 8 2008;52(2):124-131. PMID 18598891
25. Bennet A, Di Angelantonio E, Erqou S, et al. Lipoprotein(a) levels and risk of future coronary heart disease: large-scale prospective data. *Arch Intern Med.* Mar 24 2008;168(6):598-608. PMID 18362252
26. Smolders B, Lemmens R, Thijs V. Lipoprotein (a) and stroke: a meta-analysis of observational studies. *Stroke.* Jun 2007;38(6):1959-1966. PMID 17478739
27. CMS.gov, Centers for Medicare and Medicaid Services Local Coverage Determination (LCD): B-type Natriuretic Peptide (BNP) Testing (L33573)<https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33573&ver=12&SearchType=Advanced&CoverageSelection=Local&PolicyType=Both&s=47&CntrctrType=13&DateTag=C&kq=true&bc=IAAAACAAAAAAA%3d%3d&>

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

