

## Medical Coverage Policy | Confocal Laser Endomicroscopy

**EFFECTIVE DATE:** 09|01|2018

**POLICY LAST UPDATED:** 03|19|2020



**Blue Cross  
Blue Shield**  
of Rhode Island

### OVERVIEW

Confocal laser endomicroscopy (CLE), also known as confocal fluorescent endomicroscopy and optical endomicroscopy, allows *in vivo* microscopic imaging of cells during endoscopy. CLE is proposed for a variety of purposes, especially as a real-time alternative to biopsy/polypectomy and histopathologic analysis during colonoscopy and for targeting areas to undergo biopsy in patients with inflammatory bowel disease or Barrett esophagus.

### MEDICAL CRITERIA

Not applicable

### PRIOR AUTHORIZATION

Not applicable

### POLICY STATEMENT

#### BlueCHiP for Medicare

Use of confocal laser endomicroscopy is not covered as the evidence is insufficient to determine the effects of the technology on health outcomes.

#### Commercial Products

Use of confocal laser endomicroscopy is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

### COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

### BACKGROUND

Confocal laser endomicroscopy (CLE), also known as confocal fluorescent endomicroscopy and optical endomicroscopy, allows *in vivo* microscopic imaging of the mucosal epithelium during endoscopy. The process uses light from a low-power laser to illuminate tissue and, subsequently, the same lens detects light reflected from the tissue through a pinhole. The term *confocal* refers to having both illumination and collection systems in the same focal plane. Light reflected and scattered at other geometric angles that is not reflected through the pinhole is excluded from detection, which dramatically increases the resolution of CLE images.

To date, 2 CLE systems have been cleared by the U.S. Food and Drug Administration (FDA). One is an endoscope-based system with a confocal probe incorporated onto the tip of a conventional endoscope. The other is a probe-based system; the probe is placed through the biopsy channel of a conventional endoscope. The depth of view is up to 250  $\mu\text{m}$  with the endoscopic system and about 120  $\mu\text{m}$  with the probe-based system. A limited area can be examined - no more than 700  $\mu\text{m}$  in the endoscopic-based system and less with the probe-based system. As pointed out in systematic reviews, the limited viewing area emphasizes the need for careful conventional endoscopy to target areas for evaluation. Both CLE systems are optimized using a contrast agent. The most widely used agent is intravenous fluorescein, which is FDA-approved for ophthalmologic imaging of blood vessels when used with a laser scanning ophthalmoscope.

Unlike techniques such as chromoendoscopy, which are primarily intended to improve the sensitivity of colonoscopy, CLE is unique in that it is designed to characterize the cellular structure of lesions immediately. CLE can thus potentially be used to make a diagnosis of polyp histology, particularly in association with screening or surveillance colonoscopy, which could allow for small hyperplastic lesions to be overlooked rather than removed and sent for histologic evaluation. Using CLE would reduce risks associated with biopsy and reduce the number of biopsies and histologic evaluations.

Another potential application of CLE technology is targeting areas for biopsy in patients with Barrett esophagus undergoing surveillance endoscopy. This alternative to the current standard approach, recommended by the American Gastroenterological Association, is that patients with Barrett esophagus who do not have dysplasia undergo endoscopic surveillance every 3 to 5 years. The American Gastroenterological Association has further recommended that random 4-quadrant biopsies every 2 cm be taken with white-light endoscopy in patients without known dysplasia.

Other potential uses of CLE under investigation include better diagnosis and differentiation of conditions such as gastric metaplasia, lung cancer, and bladder cancer.

As noted, limitations of CLE systems include a limited viewing area and depth of view. Another issue is standardization of systems for classifying lesions viewed with CLE devices. Although there is currently no internationally accepted classification system for colorectal lesions, two systems have been used in a number of studies conducted in different countries. They are the Mainz criteria for endoscopy-based CLE devices and the Miami classification system for probe-based CLE devices. Lesion classification systems are less developed for non-gastrointestinal lesions viewed by CLE devices (e.g., those in the lung or bladder). Another challenge is the learning curve for obtaining high-quality images and classifying lesions. Several recent studies, however, have found that the ability to acquire high-quality images and interpret them accurately can be learned relatively quickly; these studies were specific to colorectal applications of CLE.

#### REGULATORY STATUS

Two CLE devices have been cleared for marketing by FDA through the 510(k) process.

Cellvizio® (Mauna Kea Technologies, Paris, France) is a confocal microscopy with a fiber optic probe (ie, a probe-based CLE system). The device consists of a laser scanning unit, proprietary software, a flat-panel display, and miniaturized fiber optic probes. The F-600 system, cleared by the FDA in 2006, can be used with any standard endoscope with a working channel of at least 2.8 mm. According to FDA, the device is intended for imaging the internal microstructure of tissues in the anatomic tract (gastrointestinal or respiratory) that are accessed by an endoscope. The 100 series version of the system was cleared by FDA in 2015 for imaging the internal microstructure of tissues and for visualization of body cavities organs and canals during endoscopic and laparoscopic surgery. In 2018, the CranoFlex™ Confocal Miniprobe (Mauna Kea Technologies) was cleared to “provide visualization within central nervous system during cranial diagnostic and therapeutic procedures such as tumor biopsy and resection.”

Confocal Video Colonoscope (Pentax Medical) is an endoscopy-based CLE system. The EC-3S7OCILK system, cleared by FDA in 2004, is used with a Pentax Video Processor and with a Pentax Confocal Laser System. According to the FDA, the device is intended to provide optical and microscopic visualization of and therapeutic access to the lower gastrointestinal tract.

For individuals who have suspected or known colorectal lesions who receive CLE as an adjunct to colonoscopy, the evidence includes multiple diagnostic accuracy studies. The relevant outcomes are overall survival, disease-specific survival, test validity, and resource utilization. While the reported sensitivity and specificity in these studies are high, it is uncertain whether the accuracy is sufficiently high to replace biopsy/polypectomy and histopathologic analysis. Moreover, issues remain concerning the use

of this technology in clinical practice (eg, the learning curve, interpretation of lesions). The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have Barrett esophagus who are undergoing surveillance who receive CLE with targeted biopsy, the evidence includes several RCTs and a meta-analysis. The relevant outcomes are overall survival, disease-specific survival, test validity, and resource utilization. Evidence from RCTs has suggested CLE is more sensitive than standard endoscopy for identifying areas of dysplasia. However, a 2014 meta-analysis found that the pooled sensitivity, specificity, and negative predictive value of available studies were not sufficiently high to replace the standard surveillance protocol. National guidelines continue to recommend 4-quadrant random biopsies for patients with Barrett esophagus undergoing surveillance. The single RCT, which compared high-definition white-light endoscopy with high-definition white-light endoscopy plus CLE, was stopped early because an interim analysis did not find a between-group difference in outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have gastrointestinal lesions and have had endoscopic treatment who receive CLE to assess the adequacy of endoscopic treatment, the evidence includes a systematic review. The relevant outcomes are overall survival, disease-specific survival, test validity, and resource utilization. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have a suspicion of a condition diagnosed by identification and biopsy of lesions (e.g., lung, bladder, or gastric cancer) who receive CLE, the evidence includes a small number of diagnostic accuracy studies. The relevant outcomes are overall survival, disease-specific survival, test validity, and resource utilization. There is limited evidence on the diagnostic accuracy of CLE for these other indications. The evidence is insufficient to determine the effects of the technology on health outcomes.

## CODING

The following codes are not covered for BlueCHiP for Medicare and not medically necessary for Commercial Products:

**43206** Esophagoscopy, flexible, transoral; with optical endomicroscopy

**43252** Esophagogastroduodenoscopy, flexible, transoral; with optical endomicroscopy

**88375** Optical endomicroscopic image(s), interpretation and report, real-time or referred,  
Each endoscopic session

**0397T** Endoscopic retrograde cholangiopancreatography (ERCP), with optical endomicroscopy  
(List separately in addition to code for primary procedure)

## RELATED POLICIES

New Technology

## PUBLISHED

Provider Update, May 2020

Provider Update, July 2019

Provider Update, August 2018

## REFERENCES:

1. American Gastroenterological Association, Spechler SJ, Sharma P, et al. American Gastroenterological Association medical position statement on the management of Barrett's esophagus. *Gastroenterology*. Mar 2011;140(3):1084-1091. PMID 21376940
2. Salvatori F, Siciliano S, Maione F, et al. Confocal laser endomicroscopy in the study of colonic mucosa in IBD patients: a review. *Gastroenterol Res Pract*. Apr 2012;2012:525098. PMID 22474440
3. Neumann H, Vieth M, Atreya R, et al. Prospective evaluation of the learning curve of confocal laser endomicroscopy in patients with IBD. *Histol Histopathol*. Jul 2011;26(7):867-872. PMID 21630216

4. Buchner AM, Gomez V, Heckman MG, et al. The learning curve of in vivo probe-based confocal laser endomicroscopy for prediction of colorectal neoplasia. *Gastrointest Endosc.* Mar 2011;73(3):556-560. PMID 21353852
5. Su P, Liu Y, Lin S, et al. Efficacy of confocal laser endomicroscopy for discriminating colorectal neoplasms from non-neoplasms: a systematic review and meta-analysis. *Colorectal Dis.* Jan 2013;15(1):e1-12. PMID 23006609
6. Dong YY, Li YQ, Yu YB, et al. Meta-analysis of confocal laser endomicroscopy for the detection of colorectal neoplasia. *Colorectal Dis.* Sep 2013;15(9):e488-495. PMID 23810105
7. Wanders LK, East JE, Uitentuis SE, et al. Diagnostic performance of narrowed spectrum endoscopy, autofluorescence imaging, and confocal laser endomicroscopy for optical diagnosis of colonic polyps: a meta-analysis. *Lancet Oncol.* Dec 2013;14(13):1337-1347. PMID 24239209
8. Xie XJ, Li CQ, Zuo XL, et al. Differentiation of colonic polyps by confocal laser endomicroscopy. *Endoscopy.* Feb 2011;43(2):87-93. PMID 21038291
9. Buchner AM, Shahid MW, Heckman MG, et al. Comparison of probe-based confocal laser endomicroscopy with virtual chromoendoscopy for classification of colon polyps. *Gastroenterology.* Mar 2010;138(3):834-842. PMID 19909747
10. Shahid MW, Buchner AM, Raimondo M, et al. Accuracy of real-time vs. blinded offline diagnosis of neoplastic colorectal polyps using probe-based confocal laser endomicroscopy: a pilot study. *Endoscopy.* Apr 2012;44(4):343-348. PMID 22382851
11. Hlavaty T, Huorka M, Koller T, et al. Colorectal cancer screening in patients with ulcerative and Crohn's colitis with use of colonoscopy, chromoendoscopy and confocal endomicroscopy. *Eur J Gastroenterol Hepatol.* Aug 2011;23(8):680-689. PMID 21602687
12. Xiong YQ, Ma SJ, Zhou JH, et al. A meta-analysis of confocal laser endomicroscopy for the detection of neoplasia in patients with Barrett's esophagus. *J Gastroenterol Hepatol.* Jun 2016;31(6):1102-1110. PMID 26676646
13. Gupta A, Attar BM, Koduru P, et al. Utility of confocal laser endomicroscopy in identifying high-grade dysplasia and adenocarcinoma in Barrett's esophagus: a systematic review and meta-analysis. *Eur J Gastroenterol Hepatol.* Apr 2014;26(4):369-377. PMID 24535597
14. Canto MI, Anandasabapathy S, Brugge W, et al. In vivo endomicroscopy improves detection of Barrett's esophagus-related neoplasia: a multicenter international randomized controlled trial (with video). *Gastrointest Endosc.* Feb 2014;79(2):211-221. PMID 24219822
15. Sharma P, Meining AR, Coron E, et al. Real-time increased detection of neoplastic tissue in Barrett's esophagus with probe-based confocal laser endomicroscopy: final results of an international multicenter, prospective, randomized, controlled trial. *Gastrointest Endosc.* Sep 2011;74(3):465-472. PMID 21741642
16. Dunbar KB, Okolo P, 3rd, Montgomery E, et al. Confocal laser endomicroscopy in Barrett's esophagus and endoscopically inapparent Barrett's neoplasia: a prospective, randomized, double-blind, controlled, crossover trial. *Gastrointest Endosc.* Oct 2009;70(4):645-654. PMID 19559419
17. Ypsilantis E, Pissas D, Papagrigoriadis S, et al. Use of confocal laser endomicroscopy to assess the adequacy of endoscopic treatment of gastrointestinal neoplasia: a systematic review and meta-analysis. *Surg Laparosc Endosc Percutan Tech.* Feb 2015;25(1):1-5. PMID 24910941
18. Wallace MB, Crook JE, Saunders M, et al. Multicenter, randomized, controlled trial of confocal laser endomicroscopy assessment of residual metaplasia after mucosal ablation or resection of GI neoplasia in Barrett's esophagus. *Gastrointest Endosc.* Sep 2012;76(3):539-547 e531. PMID 22749368
19. Sorokina A, Danilevskaya O, Averyanov A, et al. Comparative study of ex vivo probe-based confocal laser endomicroscopy and light microscopy in lung cancer diagnostics. *Respirology.* Aug 2014;19(6):907-913. PMID 24909555
20. Wellikoff AS, Holladay RC, Downie GH, et al. Comparison of in vivo probe-based confocal laser endomicroscopy with histopathology in lung cancer: A move toward optical biopsy. *Respirology.* Aug 2015;20(6):967-974. PMID 26094505
21. Fuchs FS, Zirlik S, Hildner K, et al. Confocal laser endomicroscopy for diagnosing lung cancer in vivo. *Eur Respir J.* Jun 2013;41(6):1401-1408. PMID 22997220

22. Sonn GA, Jones SN, Tarin TV, et al. Optical biopsy of human bladder neoplasia with in vivo confocal laser endomicroscopy. *J Urol.* Oct 2009;182(4):1299-1305. PMID 19683270
23. Liu JJ, Droller MJ, Liao JC. New optical imaging technologies for bladder cancer: considerations and perspectives. *J Urol.* Aug 2012;188(2):361-368. PMID 22698620
24. ASGE Technology Committee. Confocal laser endomicroscopy. *Gastrointest Endosc.* Dec 2014;80(6):928-938. PMID 25442092
25. Hirota WK, Zuckerman MJ, Adler DG, et al. ASGE guideline: the role of endoscopy in the surveillance of premalignant conditions of the upper GI tract. *Gastrointest Endosc.* Apr 2006;63(4):570-580. PMID 165648

**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

