

Medical Coverage Policy | Carotid
Angioplasty/Stenting Without Embolic Protection



EFFECTIVE DATE: 04|07|2005
POLICY LAST UPDATED: 06|04|2021

OVERVIEW

Carotid artery angioplasty with stenting (CAS) is a treatment for carotid stenosis that is intended to prevent future stroke. It is an alternative to medical therapy and a less invasive alternative to carotid endarterectomy (CEA).

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

Carotid angioplasty without associated stenting and embolic protection is not covered as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Commercial Products

Carotid angioplasty without associated stenting and embolic protection is considered not medically necessary the evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Carotid artery angioplasty with stenting is a treatment for carotid stenosis that is intended to prevent future stroke. It is an alternative to medical therapy and a less-invasive alternative to carotid endarterectomy.

Combined with optimal medical management, carotid angioplasty with or without stenting has been evaluated as an alternative to carotid endarterectomy (CEA). Carotid artery stenting (CAS) involves the introduction of coaxial systems of catheters, microcatheters, balloons, and other devices. The procedure is most often performed through the femoral artery, but a transcervical approach can also be used to avoid traversing the aortic arch. The procedure typically takes 20–40 minutes. Interventionalists almost uniformly use an embolic protection device (EPD) designed to reduce the risk of stroke caused by thromboembolic material dislodged during CAS. Embolic protection devices can be deployed proximally (with flow reversal) or distally (using a filter). Carotid angioplasty is rarely performed without stent placement.

Proposed advantages of CAS over CEA include:

- General anesthesia is not used (although CEA can be performed under local/regional anesthesia)
- Cranial nerve palsies are infrequent sequelae (although almost all following CEA resolve over time)
- Simultaneous procedures may be performed on the coronary and carotid arteries

A number of carotid artery stents and embolic protection devices (EPDs) have been approved by the U.S. Food and Drug Administration (FDA) through the premarket approval or the 510(k) process.

Each FDA-approved carotid stent is indicated for combined use with an EPD to reduce risk of stroke in patients considered at increased risk for periprocedural complications from carotid endarterectomy (CEA) who are symptomatic with greater than 50% stenosis, or asymptomatic with greater than 80% stenosis with degree of stenosis assessed by ultrasound or angiogram, with computed tomography angiography also used. Patients are considered at increased risk for complications during CEA if affected by any item from a list of anatomic features and comorbid conditions included in each stent system's Information for Prescribers.

The RX Acculink™ Carotid Stent System is also approved for use in conventional risk patients (not considered at increased risk for complications during CEA) with symptoms and 70% or more stenosis by ultrasound or 50% or more stenosis by angiogram, and asymptomatic patients with 70% or more stenosis by ultrasound or 60% or more stenosis by angiogram.

The FDA-approved stents and EPDs differ in the deployment methods used once they reach the target lesion, with the rapid exchange (RX) devices designed for more rapid stent and filter expansion. FDA has mandated postmarketing studies for EPDs, including longer follow-up for patients already reported to FDA and additional registry studies, primarily to compare outcomes as a function of clinician training and facility experience. Each manufacturer's system is available in various configurations (e.g., straight or tapered) and sizes (diameters and lengths) to match the vessel lumen that will receive the stent.

In 2015, the ENROUTE™ Transcarotid NPS was cleared for marketing by FDA through the 510(k) process. ENROUTE™ is a flow-reversal device designed to be placed via direct carotid access.

In 2007, a decision memo reaffirmed the Centers for Medicare and Medicaid Services (CMS) previous decision following a request to expand coverage while clarifying that "CAS is only covered when used with an embolic protection device and is, therefore, not covered if deployment of the distal embolic protection device is not technically possible." In 2008 in the sixth reconsideration, and on, in 2009 in the seventh reconsideration, CMS reaffirmed their prior coverage decisions.

Therefore, Carotid angioplasty without associated stenting and embolic protection is considered not covered for Medicare Advantage Plans and not medically necessary for Commercial products as the evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

The following code is not covered for Medicare Advantage Plans and not medically necessary for Commercial Products:

37216 Transcatheter placement of intravascular stent(s), cervical carotid artery, percutaneous; without distal embolic protection

RELATED POLICIES

Preauthorization via Web-Based tool for Procedures

PUBLISHED

Provider Update, August 2021
Provider Update, July 2020
Provider Update, August 2019
Provider Update, Sep 2018
Provider Update, October 2017

REFERENCES

1. Centers for Medicare & Medicaid Services (CMS). Decision Memo for Carotid Artery Stenting (CAG-00085R). 2005; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=157> Accessed March 23, 2021
2. Jordan WD, Jr., Voellinger DC, Fisher WS, et al. A comparison of carotid angioplasty with stenting versus endarterectomy with regional anesthesia. *J Vasc Surg*. Sep 1998;28(3):397-402; discussion 402-393. PMID 9737448
3. Vincent S, Eberg M, Eisenberg MJ, et al. Meta-Analysis of Randomized Controlled Trials Comparing the Long-Term Outcomes of Carotid Artery Stenting Versus Endarterectomy. *Circ Cardiovasc Qual Outcomes*. Oct 2015; 8(6 Suppl 3): S99-108. PMID 26515216
4. Gurm HS, Yadav JS, Fayad P, et al. Long-Term Results of Carotid Stenting versus Endarterectomy in High-Risk Patients. *New England Journal of Medicine*; April 10, 2008;358;15;1572-1579.
5. Mas J, Chatellier G, Beyssen B, et al. Endarterectomy versus Stenting in Patients with Symptomatic Severe Carotid Stenosis. *NEJM*: 2007; 355:1660-1671
6. Yadav JS, Wholey MH, Kuntz RE, et al. Protected Carotid-Artery Stenting versus Endarterectomy in High-Risk Patients. *NEJM*: 2004; 351:1493-1501.
7. Arquizan C, Trinquart L, Touboul PJ, et al. Restenosis is more frequent after carotid stenting than after endarterectomy: the EVA-3S study. *Stroke*. Apr 2011; 42(4):1015-1020. PMID 21311065
8. Bonati LH, Dobson J, Featherstone RL, et al. Long-term outcomes after stenting versus endarterectomy for treatment of symptomatic carotid stenosis: the International Carotid Stenting Study (ICSS) randomised trial. *Lancet*. Oct 14 2014. PMID 25453443
9. Silver FL, Mackey A, Clark WM, et al. Safety of Stenting and Endarterectomy by Symptomatic Status in the Carotid Revascularization Endarterectomy Versus Stenting Trial (CREST). *Stroke*. Feb 9 2011. PMID 21307169
32. Lal BK, Beach KW, Roubin GS, et al. Restenosis after carotid artery stenting and endarterectomy: a secondary analysis of CREST, a randomised controlled trial. *Lancet Neurol*. Sep 2012; 11(9):755-763. PMID 22857850
10. Lee VH, Brown RD, Jr., Mandrekar JN, et al. Incidence and outcome of cervical artery dissection: a population based study. *Neurology*. Nov 28 2006; 67(10):1809-1812. PMID 17130413
11. Salzler GG, Farber A, Rybin DV, et al. The association of Carotid Revascularization Endarterectomy versus Stent Trial (CREST) and Centers for Medicare and Medicaid Services Carotid Guideline Publication on utilization and outcomes of carotid stenting among "high-risk" patients. *J Vasc Surg*. Jul 2017;66(1):104-111 e101. PMID 28502543
12. Spangler EL, Goodney PP, Schanzer A, et al. Outcomes of carotid endarterectomy versus stenting in comparable medical risk patients. *J Vasc Surg*. Nov 2014; 60(5): 1227- 1231.e1. PMID 24953899
13. Schirmer CM, Atalay B, Malek AM. Endovascular recanalization of symptomatic flow-limiting cervical carotid dissection in an isolated hemisphere. *Neurosurg Focus*. Jun 2011; 30(6): E16. PMID 21631217
14. Bonati LH, Lyrer P, Ederle J, et al. Percutaneous transluminal balloon angioplasty and stenting for carotid artery stenosis. *Cochrane Database Syst Rev*. Sep 12 2012(9):CD000515. PMID 22972047
15. Kuliha M, Roubec M, Prochazka V, et al. Randomized clinical trial comparing neurological outcomes after carotid endarterectomy or stenting. *Br J Surg*. Feb 2015; 102(3): 194-201. PMID 25511816

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

