



**EFFECTIVE DATE:** 12|01|2014  
**POLICY LAST UPDATED:** 05|19|2021

## OVERVIEW

This policy documents coverage guidelines for Medicare Advantage Plans and Commercial Products for enhanced external counterpulsation (EECP) used in outpatient treatment. EECP is a noninvasive treatment used to augment diastolic pressure, decrease left ventricular afterload, and increase venous return. It has been studied primarily as a treatment for patients with refractory angina and heart failure.

## MEDICAL CRITERIA

Not applicable

## PRIOR AUTHORIZATION

Not applicable

## POLICY STATEMENT

### Medicare Advantage Plans

EECP is covered for Medicare Advantage Plan members.

**Note:** Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all Medicare Advantage Plan policies. Therefore, Medicare Advantage Plan policies may differ from Commercial products. In some instances, benefits for Medicare Advantage Plans may be greater than what is allowed by the CMS.

### Commercial Products

EECP used in outpatient treatment is not medically necessary for all indications, including but not limited to, treatment of chronic stable angina pectoris, heart failure, erectile dysfunction, or ischemic stroke as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate section of the Benefit Booklet, Evidence of Coverage or Subscriber Agreement for services not medically necessary.

## BACKGROUND

Enhanced external counterpulsation (EECP) uses timed, sequential inflation of pressure cuffs on the calves, thighs, and buttocks to augment diastolic pressure, decrease left ventricular afterload, and increase venous return. The proposed mechanism of action is the augmentation of diastolic pressure by displacement of a volume of blood backward into the coronary arteries during diastole when the heart is in a state of relaxation and resistance in the coronary arteries is at a minimum. The resulting increase in coronary artery perfusion pressure may enhance coronary collateral development or increase flow through existing collaterals. Also, when the left ventricular contracts, it faces reduced aortic counterpressure, because the counterpulsation has somewhat emptied the aorta. EECP has been primarily investigated as a treatment for chronic stable angina.

Intra-aortic balloon counterpulsation is a more familiar, invasive form of counterpulsation that is used as a method of temporary circulatory assistance for the ischemic heart, often after acute myocardial infarction. In contrast, EECP is thought to provide a permanent effect on the heart by enhancing the coronary collateral development. A full course of therapy usually consists of 35, one-hour treatments, which may be offered once or twice daily, usually 5 days a week. The multiple components of the procedure include the use of the device itself, finger plethysmography to follow the blood flow, continuous electrocardiograms to trigger inflation and deflation, and optional use of pulse oximetry to measure oxygen saturation before and after treatment.

### **Regulatory Status**

A variety of enhanced external counterpulsation (EECP) devices have been cleared for marketing by the Food and Drug Administration (FDA) through the 510(k) process. Examples of EECP devices with FDA clearance and the manufacturers are: External Counterpulsation System (Vamed Medical Instrument), Pure Flow External Counter-Pulsation Device (Xtreem Pulse), Renew® NCP-5 External Counterpulsation System (Renew Group), ECP Health System Model (ECP Health), CardiAssist™ Counter Pulsation System (Cardiomedics), ACS Model NCP-2 External Counterpulsation Device (Applied Cardiac Systems), and the EECP® Therapy System (Vasomedical).

For individuals who have chronic stable angina who receive enhanced external counterpulsation (EECP), the evidence includes randomized controlled trials (RCTs), observational studies, and systematic reviews. Relevant outcomes are overall survival, symptoms, morbid events, and functional outcomes. There is a single blinded RCT that includes clinical outcomes, and it reported benefit on only 1 of 4 main angina outcomes. Additional small RCTs have reported changes in physiologic measures associated with EECP but did not provide relevant evidence on clinical efficacy. Because of the variable natural history of angina, the multiple confounding variables for cardiac outcomes, and the potential for a placebo effect, RCT evidence is needed. Observational studies, including registry studies with large numbers of patients, add little to determinations of efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure who receive EECP, the evidence includes RCTs, observational studies, and systematic reviews. Relevant outcomes are overall survival, symptoms, morbid events, and functional outcomes. One RCT that reported on clinical outcomes found a modest benefit with EECP on some outcomes and not others. A second RCT reported improvements on the 6-minute walk test with EECP, but had methodologic limitations: RCT findings ultimately proved inconclusive. The observational studies on EECP in heart failure have limited ability to inform the evidence on EECP due to the multiple confounding variables for cardiac outcomes and the potential for a placebo effect. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have other conditions related to ischemia or vascular dysfunction who receive EECP, the evidence includes RCTs, registry studies, and systematic reviews. Relevant outcomes are overall survival, symptoms, morbid events, and functional outcomes. Two RCTs have assessed use of EECP for treatment of central retinal artery occlusion; both trials had methodologic limitations. Registry studies of erectile function have reported improvements for some outcomes with EECP but design shortcomings limit conclusions drawn. EECP has also been used to treat acute ischemic stroke, but the evidence base in is not robust. EECP has been used in a small RCT to treat type 2 diabetes. Reported follow-up was short-term. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. Therefore, this service is considered not medically necessary for Commercial products.

### **Medicare Advantage Plans**

Medicare has published a national coverage decision on EECP that allows coverage for the following indications:

“Coverage is provided for the use of EECP for patients who have been diagnosed with disabling angina who, in the opinion of a cardiologist or cardiothoracic surgeon, are not readily amenable to surgical intervention, such as percutaneous transluminal coronary angioplasty or cardiac bypass because: 1) Their condition is inoperable, or at high risk of operative complications or post-operative failure; 2) Their coronary anatomy is not readily amendable to such procedures; or 3) They have co-morbid states which create excessive risk.”

Medicare's coverage decision also noted that while the U.S. Food and Drug Administration has cleared EECP "for use in treating a variety of cardiac conditions, including stable or unstable angina pectoris, acute myocardial infarction and cardiogenic shock, the use of this device to treat cardiac conditions other than stable angina pectoris is not covered.

## CODING

### Medicare Advantage Plans and Commercial Products

The following HCPCS code is covered for Medicare Advantage Plans and not medically necessary for Commercial products:

**G0166** External counterpulsation, per treatment session

## RELATED POLICIES

Not applicable

## PUBLISHED

Provider Update, July 2021

Provider Update, July 2020

Provider Update, July 2019

Provider Update, July 2018

Provider Update, May 2017

## REFERENCES

- 1.Center for Medicare & Medicaid Services (CMS). National Coverage Determination for external counterpulsation (ECP) therapy for severe angina (20.20). 2006; [https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?CALId=185&CalName=PSA+\(Addition+of+ICD-9-CM+600.10%2C+Nodular+prostate+without+urinary+obstruction+and+600.11%2C+with+urinary+obstruction%2C+as+covered+indications\)&ExpandComments=n&CommentPeriod=0&NCDId=97&ncdver=2&CoverageSelection=National&ncd\\_id=20.20&ncd\\_version=2&basket=ncd%2525253A20%2525252E20%2525253A2%2525253AExternal+Counterpulsation+%25252528ECP%25252529+for+Severe+Angin&bc=gAAAABAAQEAAAA%3D%3D&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?CALId=185&CalName=PSA+(Addition+of+ICD-9-CM+600.10%2C+Nodular+prostate+without+urinary+obstruction+and+600.11%2C+with+urinary+obstruction%2C+as+covered+indications)&ExpandComments=n&CommentPeriod=0&NCDId=97&ncdver=2&CoverageSelection=National&ncd_id=20.20&ncd_version=2&basket=ncd%2525253A20%2525252E20%2525253A2%2525253AExternal+Counterpulsation+%25252528ECP%25252529+for+Severe+Angin&bc=gAAAABAAQEAAAA%3D%3D&) Accessed April 20, 2021.
- 2.Arora RR, Chou TM, Jain D, et al. The multicenter study of enhanced external counterpulsation (MUST-EECP): effect of EECP on exercise-induced myocardial ischemia and anginal episodes. *J Am Coll Cardiol.* Jun 1999; 33(7): 1833-40. PMID 10362181
- 3.Arora RR, Chou TM, Jain D, et al. Effects of enhanced external counterpulsation on Health-Related Quality of Life continue 12 months after treatment: a substudy of the Multicenter Study of Enhanced External Counterpulsation. *J Investig Med.* Jan 2002; 50(1): 25-32. PMID 11813825
- 4.Bondesson SM, Edvinsson ML, Pettersson T, et al. Reduced peripheral vascular reactivity in refractory angina pectoris: Effect of enhanced external counterpulsation. *J Geriatr Cardiol.* Dec 2011; 8(4): 215-23. PMID 22783308
- 5.Gloekler S, Meier P, de Marchi SF, et al. Coronary collateral growth by external counterpulsation: a randomised controlled trial. *Heart.* Feb 2010; 96(3): 202-7. PMID 19897461
- 6.Buschmann EE, Utz W, Pagonas N, et al. Improvement of fractional flow reserve and collateral flow by treatment with external counterpulsation (Art.Net.-2 Trial). *Eur J Clin Invest.* Oct 2009; 39(10): 866-75. PMID 19572918

7. Braith RW, Conti CR, Nichols WW, et al. Enhanced external counterpulsation improves peripheral artery flow-mediated dilation in patients with chronic angina: a randomized sham-controlled study. *Circulation*. Oct 19 2010; 122(16): 1612-20. PMID 20921442
8. Casey DP, Beck DT, Nichols WW, et al. Effects of enhanced external counterpulsation on arterial stiffness and myocardial oxygen demand in patients with chronic angina pectoris. *Am J Cardiol*. May 15 2011; 107(10): 1466-72. PMID 21420062
9. Shakouri SK, Razavi Z, Eslamian F, et al. Effect of Enhanced External Counterpulsation and Cardiac Rehabilitation on Quality of Life, Plasma Nitric Oxide, Endothelin 1 and High Sensitive CRP in Patients With Coronary Artery Disease: A Pilot Study. *Ann Rehabil Med*. Apr 2015; 39(2): 191-8. PMID 25932415
10. Holubkov R, Kennard ED, Foris JM, et al. Comparison of patients undergoing enhanced external counterpulsation and percutaneous coronary intervention for stable angina pectoris. *Am J Cardiol*. May 15 2002; 89(10): 1182-6. PMID 12008172
11. Shechter M, Matetzky S, Feinberg MS, et al. External counterpulsation therapy improves endothelial function in patients with refractory angina pectoris. *J Am Coll Cardiol*. Dec 17 2003; 42(12): 2090-5. PMID 14680732
12. Amin F, Al Hajeri A, Civelek B, et al. Enhanced external counterpulsation for chronic angina pectoris. *Cochrane Database Syst Rev*. Feb 17 2010; (2): CD007219. PMID 20166092
13. Shah SA, Shapiro RJ, Mehta R, et al. Impact of enhanced external counterpulsation on Canadian Cardiovascular Society angina class in patients with chronic stable angina: a meta-analysis. *Pharmacotherapy*. Jul 2010; 30(7): 639-45. PMID 20575628
14. McKenna C, McDaid C, Suekarran S, et al. Enhanced external counterpulsation for the treatment of stable angina and heart failure: a systematic review and economic analysis. *Health Technol Assess*. Apr 2009; 13(24): iii-iv, ix-xi, 1-90. PMID 19409154
15. Barsheshet A, Hod H, Shechter M, et al. The effects of external counterpulsation therapy on circulating endothelial progenitor cells in patients with angina pectoris. *Cardiology*. 2008; 110(3): 160-6. PMID 18057883
16. Qin X, Deng Y, Wu D, et al. Does Enhanced External Counterpulsation (EECP) Significantly Affect Myocardial Perfusion?: A Systematic Review Meta-Analysis. *PLoS One*. 2016; 11(4): e0151822. PMID 27045935
17. Soran O, Kennard ED, Bart BA, et al. Impact of external counterpulsation treatment on emergency department visits and hospitalizations in refractory angina patients with left ventricular dysfunction. *Congest Heart Fail*. Jan-Feb 2007; 13(1): 36-40. PMID 17268208
18. Loh PH, Cleland JG, Louis AA, et al. Enhanced external counterpulsation in the treatment of chronic refractory angina: a long-term follow-up outcome from the International Enhanced External Counterpulsation Patient Registry. *Clin Cardiol*. Apr 2008; 31(4): 159-64. PMID 18404725
19. Thakkar BV, Hirsch AT, Satran D, et al. The efficacy and safety of enhanced external counterpulsation in patients with peripheral arterial disease. *Vasc Med*. Feb 2010; 15(1): 15-20. PMID 19841026
20. Kumar A, Aronow WS, Vadnerkar A, et al. Effect of enhanced external counterpulsation on clinical symptoms, quality of life, 6-minute walking distance, and echocardiographic measurements of left ventricular systolic and diastolic function after 35 days of treatment and at 1-year follow up in 47 patients with chronic refractory angina pectoris. *Am J Ther*. Mar-Apr 2009; 16(2): 116-8. PMID 19300038
21. Pettersson T, Bondesson S, Cojocar D, et al. One year follow-up of patients with refractory angina pectoris treated with enhanced external counterpulsation. *BMC Cardiovasc Disord*. Jun 15 2006; 6: 28. PMID 16776842
22. Loh PH, Louis AA, Windram J, et al. The immediate and long-term outcome of enhanced external counterpulsation in treatment of chronic stable refractory angina. *J Intern Med*. Mar 2006; 259(3): 276-84. PMID 16476105
23. Feldman AM, Silver MA, Francis GS, et al. Treating heart failure with enhanced external counterpulsation (EECP): design of the Prospective Evaluation of EECP in Heart Failure (PEECH) trial. *J Card Fail*. Apr 2005; 11(3): 240-5. PMID 15812754
24. Feldman AM, Silver MA, Francis GS, et al. Enhanced external counterpulsation improves exercise tolerance in patients with chronic heart failure. *J Am Coll Cardiol*. Sep 19 2006; 48(6): 1198-205. PMID 16979005

25. Abbott-Smith CW, Chung ES, Varricchio T, et al. Enhanced external counterpulsation improves exercise duration and peak oxygen consumption in older patients with heart failure: a subgroup analysis of the PEECH trial. *Congest Heart Fail.* Nov-Dec 2006; 12(6): 307-11. PMID 17170583
26. Rampengan SH, Prihartono J, Siagian M, et al. The Effect of Enhanced External Counterpulsation Therapy and Improvement of Functional Capacity in Chronic Heart Failure patients: a Randomized Clinical Trial. *Acta Med Indones.* Oct 2015; 47(4): 275-82. PMID 26932695
27. Soran O, Kennard ED, Kelsey SF, et al. Enhanced external counterpulsation as treatment for chronic angina in patients with left ventricular dysfunction: a report from the International EECPP Patient Registry (IEPR). *Congest Heart Fail.* Nov-Dec 2002; 8(6): 297-302. PMID 12461318
28. Lawson WE, Kennard ED, Holubkov R, et al. Benefit and safety of enhanced external counterpulsation in treating coronary artery disease patients with a history of congestive heart failure. *Cardiology.* 2001; 96(2): 78-84. PMID 11740136
29. Lawson WE, Silver MA, Hui JC, et al. Angina patients with diastolic versus systolic heart failure demonstrate comparable immediate and one-year benefit from enhanced external counterpulsation. *J Card Fail.* Feb 2005; 11(1): 61-6. PMID 15704066
30. Vijayaraghavan K, Santora L, Kahn J, et al. New graduated pressure regimen for external counterpulsation reduces mortality and improves outcomes in congestive heart failure: a report from the Cardiomedics External Counterpulsation Patient Registry. *Congest Heart Fail.* May-Jun 2005; 11(3): 147-52. PMID 15947536
31. Soran O, Fleishman B, Demarco T, et al. Enhanced external counterpulsation in patients with heart failure: a multicenter feasibility study. *Congest Heart Fail.* Jul-Aug 2002; 8(4): 204-8, 227. PMID 12147943
32. Fraser SG, Adams W. Interventions for acute non-arteritic central retinal artery occlusion. *Cochrane Database Syst Rev.* Jan 21 2009; (1): CD001989. PMID 19160204
33. Werner D, Michalk F, Harazny J, et al. Accelerated reperfusion of poorly perfused retinal areas in central retinal artery occlusion and branch retinal artery occlusion after a short treatment with enhanced external counterpulsation. *Retina.* Aug 2004; 24(4): 541-7. PMID 15300074
34. Lawson WE, Hui JC, Kennard ED, et al. Effect of enhanced external counterpulsation on medically refractory angina patients with erectile dysfunction. *Int J Clin Pract.* May 2007; 61(5): 757-62. PMID 17493089
35. Han JH, Leung TW, Lam WW, et al. Preliminary findings of external counterpulsation for ischemic stroke patient with large artery occlusive disease. *Stroke.* Apr 2008; 39(4): 1340-3. PMID 18309160
36. Lin S, Liu M, Wu B, et al. External counterpulsation for acute ischaemic stroke. *Cochrane Database Syst Rev.* Jan 18 2012; 1: CD009264. PMID 22259001
37. Sardina PD, Martin JS, Avery JC, et al. Enhanced external counterpulsation (EECP) improves biomarkers of glycemic control in patients with non-insulin-dependent type II diabetes mellitus for up to 3 months following treatment. *Acta Diabetol.* Oct 2016; 53(5): 745-52. PMID 27179825
38. Sardina PD, Martin JS, Dzieza WK, et al. Enhanced external counterpulsation (EECP) decreases advanced glycation end products and proinflammatory cytokines in patients with non-insulin-dependent type II diabetes mellitus for up to 6 months following treatment. *Acta Diabetol.* Oct 2016; 53(5): 753-60. PMID 27278477
39. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* Dec 18 2012; 60(24): e44-e164. PMID 23182125
40. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* Nov 04 2014; 64(18): 1929-49. PMID 25077860

41. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. Oct 15 2013; 128(16): e240-327. PMID 23741058
42. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. Aug 08 2017; 136(6): e137-e161. PMID 28455343

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

