

## Payment Policy | Monitored Anesthesia Care for Gastrointestinal Endoscopic Procedures



**EFFECTIVE DATE:** 07|01|2018

**POLICY LAST UPDATED:** 11|17|2021

### OVERVIEW

Monitored Anesthesia Care (MAC) is anesthesia care involves a drug-induced depression of consciousness during which the patient may respond purposefully to verbal commands (either alone or accompanied by light tactile stimulation), and requires monitoring of the patient by a practitioner who is qualified to administer anesthesia. Typically, cardiovascular function is maintained, and no interventions to maintain a patent airway are required (spontaneous ventilation is usually adequate). Indications for MAC depend on the nature of the procedure, the patient's clinical condition, and/or the potential need to convert to a general or regional anesthetic.

The intent of this policy is to address monitored anesthesia care services for gastrointestinal endoscopic diagnostic or therapeutic procedures performed in the outpatient setting.

### MEDICAL CRITERIA

Not applicable

### PRIOR AUTHORIZATION

#### Medicare Advantage Plans and Commercial Products

Prior authorization is not required.

### POLICY STATEMENT

#### Medicare Advantage Plans and Commercial Products

Monitored anesthesia care for gastrointestinal endoscopic procedures is a covered service for elective upper and lower endoscopy for members with a higher risk for sedation-related complications.

Member's medical records must document that services are medically necessary for the care provided. Blue Cross Blue Shield of Rhode Island (BCBSRI) maintains the right to audit the services provided to our members, regardless of the participation status of the provider. All documentation must be available to BCBSRI upon request. Failure to produce the requested information may result in denial or retraction of payment.

### COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for applicable anesthesia/surgery services coverage/benefits.

### BACKGROUND

Use of monitored anesthesia care for upper or lower gastrointestinal (GI) endoscopy is considered appropriate in the following circumstances:

- Member is under 18 years of age; OR
- Member is over 70 years of age; OR
- Member is pregnant; OR
- Member is acutely agitated and/or uncooperative; OR
- There is an increased risk for complications due to severe co-morbidity corresponding to the American Society of Anesthesiologists (ASA) Physical Status Modifier\* of P2 or greater; OR
- There is an increased risk for airway obstruction due to anatomic variation, such as:
  - History of stridor;
  - Dysmorphic facial features;

- Oral abnormalities (e.g. macroglossia);
- Neck abnormalities (e.g. neck mass);
- Jaw abnormalities (e.g. micrognathia); OR
- Prolonged or therapeutic gastrointestinal endoscopy procedures requiring deep sedation, such as;
  - endoscopy in members with adhesions after abdominal surgery
  - endoscopic retrograde cholangiopancreatography
  - stent placement in the upper gastrointestinal tract
  - complex therapeutic procedures, such as plication of the cardioesophageal junction; OR
- Member has one of the following:
  - History of adverse reaction to sedation;
  - History of inadequate response to sedation;
  - History or anticipated intolerance to standard sedatives, such as
    - Chronic opioid use
    - Chronic benzodiazepine use
  - Documented sleep apnea;
  - Morbid obesity (e.g. BMI > 40kg/m<sup>2</sup>)
  - Active medical problems related to drug or alcohol abuse
  - Inability to follow simple commands (cognitive dysfunction, intoxication, or psychological impairment)
  - Spasticity or movement disorder complicating the procedure

**\*American Society of Anesthesiologists (ASA) Physical Status Modifiers**

P1 – A normal healthy patient

P2 – A patient with mild systemic disease

P3 – A patient with severe systemic disease

P4 – A patient with severe systemic disease that is a constant threat to life

P5 – A moribund patient who is not expected to survive without the operation

P6 – A declared brain-dead patient whose organs are being removed for donor purposes

In 2008, the American Society for Gastrointestinal Endoscopy published a guideline outlining appropriate use of sedation and anesthesia in GI endoscopy. The guideline notes the routine use of MAC for average-risk patients undergoing standard upper and lower GI endoscopy is not appropriate. Recommendations in the guideline detail the clinical situations gastroenterologists should consider when screening patients to determine the appropriate level of sedation or MAC. Recommendations are consistent with positions and guidelines developed and published by the American Society of Anesthesiologists (ASA).

The American Society of Anesthesiologists Position on Monitored Anesthesia Care (2013) states: Monitored anesthesia care is a specific anesthesia service for a diagnostic or therapeutic procedure. Indications for monitored anesthesia care include the nature of the procedure, the patient's clinical condition and/or the potential need to convert to a general or regional anesthetic. Monitored anesthesia care includes all aspects of anesthesia care – a pre-procedure visit, intra-procedure care and post-procedure anesthesia management. During monitored anesthesia care, the anesthesiologist provides or medically directs a number of specific services, including but not limited to:

- Diagnosis and treatment of clinical problems that occur during the procedure
- Support of vital functions
- Administration of sedatives, analgesics, hypnotics, anesthetic agents or other medications as necessary for patient safety
- Psychological support and physical comfort
- Provision of other medical services as needed to complete the procedure safely

Monitored anesthesia care may include varying levels of sedation, analgesia, and anxiolysis as necessary. The provider of monitored anesthesia care must be prepared and qualified to convert to general anesthesia when

necessary. If the patient loses consciousness and the ability to respond purposefully, the anesthesia care is a general anesthetic, irrespective of whether airway instrumentation is required.

“An essential component of MAC is the anesthesia assessment and management of a patient’s actual or anticipated physiological derangements or medical problems that may occur during a diagnostic or therapeutic procedure. While Monitored Anesthesia Care may include the administration of sedatives and/or analgesics often used for Moderate Sedation, the provider of MAC must be prepared and qualified to convert to general anesthesia when necessary. Additionally, a provider’s ability to intervene to rescue a patient’s airway from any sedation-induced compromise is a prerequisite to the qualifications to provide Monitored Anesthesia Care. By contrast, Moderate Sedation is not expected to induce depths of sedation that would impair the patient’s own ability to maintain the integrity of his or her airway. These components of Monitored Anesthesia Care are unique aspects of an anesthesia service that are not part of Moderate Sedation.

## **CODING**

### **Medicare Advantage Plans and Commercial Products:**

The following CPT code(s) are covered:

- 00731** Anesthesia for upper gastrointestinal endoscopic procedures, endoscope introduced proximal to duodenum; not otherwise specified
- 00732** Anesthesia for upper gastrointestinal endoscopic procedures, endoscope introduced proximal to duodenum; endoscopic retrograde cholangiopancreatography (ERCP)
- 00811** Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; not otherwise specified
- 00812** Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; screening colonoscopy
- 00813** Anesthesia for combined upper and lower gastrointestinal endoscopic procedures, endoscope introduced both proximal to and distal to the duodenum

## **RELATED POLICIES**

Anesthesia Services

## **PUBLISHED**

Provider Update January 2022  
Provider Update January 2021  
Provider Update January 2020  
Provider Update, November 2019  
Provider Update, July 2018

## **REFERENCES**

1. American Society of Anesthesiologists (ASA). Position on monitored anesthesia care (Amended October 17, 2018). 2018; <https://www.asahq.org/standards-and-guidelines/position-on-monitored-anesthesia-care>. Accessed October 17, 2020.
2. American Society of Anesthesiologists (ASA). Distinguishing monitored Anesthesia care (MAC) from moderate sedation/analgesia (Last Amended October 17, 2018). 2018; <https://www.asahq.org/standards-and-guidelines/distinguishing-monitored-anesthesia-care-mac-from-moderate-sedationanalgesia-conscious-sedation>. Accessed October 15, 2020.
3. American Society of Anesthesiologists (ASA). Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia (Last Amended on October 23, 2019). 2019; <https://www.asahq.org/standardsand-guidelines/continuum-of-depth-of-sedation-definition-of-general-anesthesia-and-levels-of-sedationanalgesia>. Accessed October 14, 2020.
4. American Society of Anesthesiologists (ASA). Statement on Respiratory Monitoring during Endoscopic Procedures (Amended October 2019). 2019; <https://www.asahq.org/standards-and-guidelines/statement-on-respiratorymonitoring-during-endoscopic-procedures>. Accessed October 18, 2020.
5. Cohen LB, Delegee MH, Aisenberg J, et al. AGA Institute review of endoscopic sedation. *Gastroenterology*. Aug 2007; 133(2): 675-701. PMID 17681185

6. Enestvedt BK, Eisen GM, Holub J, et al. Is the American Society of Anesthesiologists classification useful in risk stratification for endoscopic procedures?. *Gastrointest Endosc.* Mar 2013; 77(3): 464-71. PMID 23410699
7. Agostoni M, Fanti L, Gemma M, et al. Adverse events during monitored anesthesia care for GI endoscopy: an 8-year experience. *Gastrointest Endosc.* Aug 2011; 74(2): 266-75. PMID 21704990
8. Berzin TM, Sanaka S, Barnett SR, et al. A prospective assessment of sedation-related adverse events and patient and endoscopist satisfaction in ERCP with anesthesiologist-administered sedation. *Gastrointest Endosc.* Apr 2011; 73(4): 710-7. PMID 21316669
9. Cote GA, Hovis RM, Ansstas MA, et al. Incidence of sedation-related complications with propofol use during advanced endoscopic procedures. *Clin Gastroenterol Hepatol.* Feb 2010; 8(2): 137-42. PMID 19607937
10. Singh H, Poluha W, Cheung M, et al. Propofol for sedation during colonoscopy. *Cochrane Database Syst Rev.* Oct 08 2008; (4): CD006268. PMID 18843709
11. Poincloux L, Laquiere A, Bazin JE, et al. A randomized controlled trial of endoscopist vs. anaesthetist-administered sedation for colonoscopy. *Dig Liver Dis.* Jul 2011; 43(7): 553-8. PMID 21450542
12. Shen XC, Ao X, Cao Y, et al. Etomidate-remifentanyl is more suitable for monitored anesthesia care during gastroscopy in older patients than propofol-remifentanyl. *Med Sci Monit.* Jan 01 2015; 21: 1-8. PMID 25553506
13. Treeprasertsuk S, Rerknimitr R, Angsuwatcharakon P, et al. The safety of propofol infusion compared to midazolam and meperidine intravenous bolus for patients undergoing double balloon enteroscopy. *J Med Assoc Thai.* May 2014; 97(5): 483-9. PMID 25065086
14. Horiuchi A, Nakayama Y, Hidaka N, et al. Low-dose propofol sedation for diagnostic esophagogastroduodenoscopy: results in 10,662 adults. *Am J Gastroenterol.* Jul 2009; 104(7): 1650-5. PMID 19513021
15. Sieg A, Beck S, Scholl SG, et al. Safety analysis of endoscopist-directed propofol sedation: a prospective, national multicenter study of 24 441 patients in German outpatient practices. *J Gastroenterol Hepatol.* Mar 2014; 29(3): 517-23. PMID 24716213
16. de Paulo GA, Martins FP, Macedo EP, et al. Sedation in gastrointestinal endoscopy: a prospective study comparing nonanesthesiologist-administered propofol and monitored anesthesia care. *Endosc Int Open.* Feb 2015; 3(1): E7- E13. PMID 26134777
17. Bernards CM, Hadzic A, Suresh S, et al. Regional anesthesia in anesthetized or heavily sedated patients. *Reg Anesth Pain Med.* Sep-Oct 2008; 33(5): 449-60. PMID 18774514
18. American Society of Anesthesiologists (ASA). Guidelines for ambulatory anesthesia and surgery (Reaffirmed October 2018). 2018; American Society of Anesthesiologists (ASA). Guidelines for ambulatory anesthesia and surgery. <https://www.asahq.org/standards-and-guidelines/guidelines-for-ambulatory-anesthesia-and-surgery>. Accessed October 16, 2020.
19. Fleisher LA, Pasternak LR, Herbert R, et al. Inpatient hospital admission and death after outpatient surgery in elderly patients: importance of patient and system characteristics and location of care. *Arch Surg.* Jan 2004; 139(1): 67-72. PMID 14718279
20. Whippey A, Kostandoff G, Paul J, et al. Predictors of unanticipated admission following ambulatory surgery: a retrospective case-control study. *Can J Anaesth.* Jul 2013; 60(7): 675-83. PMID 23606232
21. American Society of Anesthesiologists (ASA). Statement on safe use of propofol (Amended October 2019). 2019; <https://www.asahq.org/standards-and-guidelines/statement-on-safe-use-of-propofol>. Accessed October 22, 2020.
22. Early DS, Lightdale JR, Vargo JJ, et al. Guidelines for sedation and anesthesia in GI endoscopy. *GastrointestEndosc.* Feb 2018; 87(2): 327-337. PMID 29306520
23. Rizk MK, Sawhney MS, Cohen J, et al. Quality indicators common to all GI endoscopic procedures. *Am J Gastroenterol.* Jan 2015; 110(1): 48-59. PMID 25448874
24. Chandrasekhara V, Early DS, Acosta RD, et al. Modifications in endoscopic practice for the elderly. *Gastrointest Endosc.* Jul 2013; 78(1): 1-7. PMID 23664042
25. Calderwood AH, Chapman FJ, Cohen J, et al. Guidelines for safety in the gastrointestinal endoscopy unit. *Gastrointest Endosc.* Mar 2014; 79(3): 363-72. PMID 24485393
26. Vargo JJ, Cohen LB, Rex DK, et al. Position statement: nonanesthesiologist administration of propofol for GI endoscopy. *Gastrointest Endosc.* Dec 2009; 70(6): 1053-9. PMID 19962497

27. Dumonceau JM, Riphaus A, Schreiber F, et al. Non-anesthesiologist administration of propofol for gastrointestinal endoscopy: European Society of Gastrointestinal Endoscopy, European Society of Gastroenterology and Endoscopy Nurses and Associates Guideline-Updated June 2015. *Endoscopy*. Dec 2015; 47(12): 1175-89. PMID 26561915

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

