

EFFECTIVE DATE: 06 | 01 | 2020

POLICY LAST UPDATED: 02 | 16 | 2022

OVERVIEW

Radiofrequency (RF) coblation is being evaluated for the treatment of plantar fasciitis, lateral epicondylitis, and various musculoskeletal tendinopathies. When utilized for tenotomy, bipolar RF energy is directed into the tendon to generate a controlled, low-temperature field of ionizing particles that break organic bonds, ablating or debriding target tissue with the goal of relieving pain and restoring function.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

Radiofrequency coblation tenotomy is not covered as a treatment for musculoskeletal conditions, including but not limited to plantar fasciitis, lateral epicondylitis, shoulder or rotator cuff tendinopathy, Achilles tendinopathy, patellar tendinopathy and wrist tendinopathy as the evidence is insufficient to determine the effects of the technology on health outcomes.

Commercial Products

Radiofrequency coblation tenotomy is not medically necessary as a treatment for musculoskeletal conditions, including but not limited to plantar fasciitis, lateral epicondylitis, shoulder or rotator cuff tendinopathy, Achilles tendinopathy, patellar tendinopathy and wrist tendinopathy as the evidence is insufficient to determine the effects of the technology on health outcomes.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Radiofrequency Coblation

Radiofrequency (RF) coblation uses bipolar low-frequency energy in an electrically conductive fluid (eg, saline) to generate a high-density plasma field around the energy source. This creates a low-temperature field of ionizing particles that break organic bonds within the target tissue. Coblation technology is used in a variety of surgical procedures, particularly related to otolaryngology and orthopedics. The proposed advantage of coblation is that the procedure provides for controlled and highly localized ablation, resulting in minimal damage to surrounding tissue. RF coblation was also found to exhibit several properties that may make it an attractive option for addressing the underlying pathophysiology of chronic tendinopathies, namely increased angiogenesis, reduction of inflammatory responses, and increased expression of growth factors. RF coblation surgical wands are utilized by orthopedic surgeons in minimally invasive arthroscopic procedures to facilitate soft tissue debridement, subacromial decompression, meniscal removal and sculpting, or tendon debridement.

Tendinopathy

Tendinopathy is a clinical pain syndrome characterized by tendon thickening due to proliferation and chronic irritation of neovascular repair tissue with a history of repetitive tendon loading. This condition commonly

results from overuse and has a high incidence rate in athletes and laborers. Clinical history should clarify predisposing training or activity and assess the level of functioning. Biomechanical abnormalities during activity should be identified and corrected. Standard treatment may, therefore, consist of biomechanical modification, activity modification, physical therapy (eg, heavy load resistance training), and nonsteroidal anti-inflammatory medication. For chronic tendinopathies, glucocorticoids should only be used in select cases (eg, rotator cuff tendinopathy). Surgical consultation following months of a well-designed physical therapy program with adjunct medical treatments can be considered if there is no improvement in pain or function. Validated and reliable functional assessment scores should be utilized by the clinician to grade symptoms and assess patient function. Examples of suitable scales include the Victoria Institute of Sport Assessment for Achilles tendinopathy. Surgical approaches may involve incisions to the paratendon and removal of adhesions and degenerate tissue. Longitudinal incisions may be made in the tendon to promote a repair response. This latter strategy has also been delivered via minimally invasive arthroscopic approaches. These approaches may also address the debridement of the neovascular supply to the tendon surface. Collectively, a prolonged recovery duration to accommodate tendon healing may be required with these interventions.

Plantar Fasciitis

Plantar fasciitis is a musculoskeletal condition characterized by pain in the plantar region of the foot that worsens upon initiation of walking and with local point tenderness elicited during a clinical examination. Radiographic and ultrasonographic studies are not typically indicated for primary diagnosis but may be useful in ruling out alternative causes and visualizing the thickening of the plantar fascia. Initial standard therapy may consist of stretching exercises, orthotics, activity and lifestyle modification, nonsteroidal anti-inflammatory drugs, splints or casts, and glucocorticoid injections. The vast majority of patients improve without surgery. Surgery is generally considered a last line of therapy and is reserved for individuals who do not respond to at least 6 to 12 months of initial, nonsurgical therapy. Surgical approaches include variations of open or endoscopic, partial or complete, plantar fascia release which may or may not include calcaneal spur resection, excision of abnormal tissue, and nerve decompression. The use of RF microtenotomy during open or percutaneous surgery has been explored alone or in combination with plantar fasciotomy.

Plantar fasciitis is one of the most common causes of foot and heel pain in adults. It is estimated to be responsible for approximately one million patient medical visits per year in the U.S. The peak incidence of the condition in the general population occurs between ages 40 and 60. There is a higher incidence rate among runners with a younger age of onset. The etiology of plantar fasciitis is poorly understood and may be multifactorial in nature. Contributing risk factors may include obesity, prolonged standing or activity, flat feet, and reduced ankle dorsiflexion. Plantar fasciitis has been reported in association with fluoride use for the treatment of osteoporosis. Differential sources of foot and heel pain may include Achilles tendinopathy, stress fractures due to osteoporosis, rheumatoid arthritis, peripheral neuropathies associated with diabetes, extrinsic factors (eg, inappropriate footwear), aging, and structural disorders.

Lateral Epicondylitis

Lateral epicondylitis, also known as tennis elbow, represents chronic tendinosis of the myotendinous group of the lateral epicondyle characterized by pain and disability. The incidence in the general population may approach 1 to 3 percent. Risk factors include smoking, obesity, forceful activity, and repetitive activity for at least two hours daily. Lateral epicondylitis is characterized by injury to the extensor carpi radialis brevis or extensor digitorum communis muscles. The condition is diagnosed through findings of localized tenderness and pain with clinical examination. Initial conservative management includes modification of activity and biomechanics, counterforce bracing or splinting, nonsteroidal anti-inflammatory drugs, and physical therapy. Surgical referral is typically reserved for patients with severe symptoms that do not improve despite compliance with an appropriately designed physical therapy program for at least six months.

Regulatory Status

In 2014, the TOPAZ® EZ Microdebrider Coblation® Wand with Integrated Finger Switch, an electrosurgical cutting and coagulation device (ArthroCare Corporation, K140521), was cleared for marketing by the U.S.

Food and Drug Administration through the 510(k) process, on the basis of an earlier predicate device (ArthroCare Topaz Wand, K080282, 2008). The surgical wands are indicated for debridement, resection, ablation, and coagulation of soft tissue and hemostasis of blood vessels in arthroscopic and orthopedic procedures, including fasciotomy, synovectomy, tenotomy, and capsulotomy of the foot and tenotomy of the knee, wrist, elbow, ankle, shoulder, and rotator cuff.

For individuals with plantar fasciitis who receive RF coblation tenotomy, the evidence includes nonrandomized, comparative cohort studies and case series. The relevant outcomes are symptoms, functional outcomes, quality of life (QOL), medication use, and treatment-related morbidity. The trials reported improved pain and functional scores over 3-12 months, with improved outcomes with open versus percutaneous approaches. However, open RF coblation microtenotomy was associated with a higher incidence of postoperative persistent pain (9.1%) compared to endoscopic plantar fasciotomy (0%) in one study, with a separate study reporting a complication rate of 33% when both interventions were used in combination. A higher number of postoperative pain recurrences at 6 and 12 months were also reported with open RF coblation microtenotomy compared to endoscopic plantar fasciotomy. The durability of this intervention is unknown as no studies have reported long-term outcomes beyond 12 months. Studies are limited by small sample sizes, heterogeneity in surgical technique (open, percutaneous, endoscopic), missing data and/or inappropriate exclusions, lack of randomization, unclear blinding practices for patient outcome assessments, and poor statistical reporting. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with lateral epicondylitis who receive RF coblation tenotomy, the evidence includes small randomized controlled trials (RCTs). The relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trials compared RF microtenotomy to open or arthroscopic elbow release surgery. Clinically meaningful improvements in pain and functional scores were noted for all treatment arms, with no significant differences between groups through one to seven years of follow-up. For disability assessments in one study, open release surgery met the threshold for a clinically meaningful improvement over RF microtenotomy at one year, though this mean difference was not statistically significant. Studies were generally underpowered or demonstrated inconsistent delivery and unclear blinding of outcome assessments and inappropriate handling of missing or crossover data. No studies featuring RF coblation tenotomy for the treatment of wrist tendinopathy were identified. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with Achilles tendinopathy who receive RF coblation tenotomy, the evidence includes a small, single-blinded RCT. The relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trial did not demonstrate an added benefit for RF microdebridement compared to surgical decompression. Pain and functional outcomes improved in both groups but were not statistically different at a six-month follow-up. The study was limited by a control group that showed significantly less severe symptom scores at baseline that did not fully meet the two-point threshold for a clinically meaningful difference in pain score reduction. Larger, adequately controlled studies with longer follow-up durations are lacking. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with shoulder or rotator cuff tendinopathy who receive RF coblation tenotomy, the evidence includes small RCTs. The relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. Trials did not demonstrate an added benefit for RF microdebridement compared to arthroscopic subacromial decompression surgery. Pain and functional outcomes improved in both groups but were not statistically different through one to two years follow-up. Neither study prespecified a clinically meaningful difference in outcome measures nor were harms assessed throughout their course. The loss to follow-up in 1 study was 18.7%. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with patellar tendinopathy who receive radiofrequency coblation tenotomy, the evidence includes one small RCT. The relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. The trial did not demonstrate an added benefit for RF microdebridement compared to mechanical debridement in patients with chondral lesions and patellar tendinopathy. The study lacked reporting with validated pain measures over time and reported a higher incidence of crepitus in patients undergoing RF microdebridement. Furthermore, the study only enrolled female participants, limiting the broader applicability of these findings. Larger studies with validated pain and functional outcome measures are required to adequately assess the technology. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

Medicare Advantage Plans and Commercial Products

There is not a specific code for this service. Claims must be filed with an unlisted CPT code.

RELATED POLICIES

Unlisted Procedures

PUBLISHED

Provider Update, April 2022

Provider Update, March 2021

Provider Update, April 2020

REFERENCES:

1. Tay KS, Ng YC, Singh IR et al. Open technique is more effective than percutaneous technique for TOPAZ radiofrequency coblation for plantar fasciitis.. *Foot Ankle Surg*, 2012 Oct 25;18(4). PMID 23093126
2. Davenport TE, Kulig K, Matharu Y et al. The EdUReP model for nonsurgical management of tendinopathy.. *Phys Ther*, 2005 Sep 27;85(10). PMID 16180958
3. Robinson JM, Cook JL, Purdam C et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy.. *Br J Sports Med*, 2001 Oct 2;35(5). PMID 11579069
4. Lohrer H, David S, Nauck T. Surgical treatment for achilles tendinopathy - a systematic review.. *BMC Musculoskelet Disord*, 2016 May 12;17:207. PMID 27165287
5. Nirschl RP, Pettrone FA. Tennis elbow. The surgical treatment of lateral epicondylitis.. *J Bone Joint Surg Am*, 1979 Sep 1;61(6A). PMID 479229
6. Chou AC, Ng SY, Su DH et al. Radiofrequency microtenotomy is as effective as plantar fasciotomy in the treatment of recalcitrant plantar fasciitis.. *Foot Ankle Surg*, 2016 Nov 5;22(4). PMID 27810027
7. Riddle DL, Schappert SM. Volume of ambulatory care visits and patterns of care for patients diagnosed with plantar fasciitis: a national study of medical doctors.. *Foot Ankle Int*, 2004 May 12;25(5). PMID 15134610
8. Riddle DL, Pulisic M, Pidcoe P et al. Risk factors for Plantar fasciitis: a matched case-control study.. *J Bone Joint Surg Am*, 2003 May 3;85(5). PMID 12728038
9. Rano JA, Fallat LM, Savoy-Moore RT. Correlation of heel pain with body mass index and other characteristics of heel pain.. *J Foot Ankle Surg*, 2002 Jan 5;40(6). PMID 11777230
10. Riggs BL, Hodgson SF, Hoffman DL et al. Treatment of primary osteoporosis with fluoride and calcium. Clinical tolerance and fracture occurrence.. *JAMA*, 1980 Feb 1;243(5). PMID 7351765
11. Shiri R, Viikari-Juntura E, Varonen H et al. Prevalence and determinants of lateral and medial epicondylitis: a population study.. *Am. J. Epidemiol.*, 2006 Sep 14;164(11). PMID 16968862
12. Struijs PA, Kerkhoffs GM, Assendelft WJ et al. Conservative treatment of lateral epicondylitis: brace versus physical therapy or a combination of both-a randomized clinical trial.. *Am J Sports Med*, 2004 Feb 24;32(2). PMID 14977675
13. Hamlin K, Munro C, Barker SL et al. Open release versus radiofrequency microtenotomy in the treatment of lateral epicondylitis: a prospective randomized controlled trial.. *Shoulder Elbow*, 2017 Dec 26;10(1). PMID 29276537

14. Wang W, Rikhranj IS, Chou ACC et al. Endoscopic Plantar Fasciotomy vs Open Radiofrequency Microtenotomy for Recalcitrant Plantar Fasciitis.. *Foot Ankle Int*, 2017 Nov 29;39(1). PMID 29182482
15. Weil L, Glover JP, Weil LS. A new minimally invasive technique for treating plantar fasciosis using bipolar radiofrequency: a prospective analysis.. *Foot Ankle Spec*, 2008 Feb 1;1(1). PMID 19825686
16. Yeap EJ, Chong KW, Yeo W et al. Radiofrequency coblation for chronic foot and ankle tendinosis.. *J Orthop Surg (Hong Kong)*, 2010 Jan 13;17(3). PMID 20065374
17. Sean NY, Singh I, Wai CK. Radiofrequency microtenotomy for the treatment of plantar fasciitis shows good early results.. *Foot Ankle Surg*, 2010 Nov 5;16(4). PMID 21047605
18. Meknas K, Al Hassoni TN, Odden-Miland NA et al. Medium-Term Results After Treatment of Recalcitrant Lateral Epicondylitis: A Prospective, Randomized Study Comparing Open Release and Radiofrequency Microtenotomy.. *Orthop J Sports Med*, 2013 Sep 1;1(4). PMID 26535247
19. Lee JH, Park I, Hyun HS et al. A Comparison of Radiofrequency-Based Microtenotomy and Arthroscopic Release of the Extensor Carpi Radialis Brevis Tendon in Recalcitrant Lateral Epicondylitis: A Prospective Randomized Controlled Study.. *Arthroscopy*, 2018 Jan 26;34(5). PMID 29366739
20. Morrison RJM, Brock TM, Reed MR et al. Radiofrequency Microdebridement Versus Surgical Decompression for Achilles Tendinosis: A Randomized Controlled Trial.. *J Foot Ankle Surg*, 2017 May 13;56(4). PMID 28495412
21. Martin RL, Chimenti R, Cuddeford T et al. Achilles Pain, Stiffness, and Muscle Power Deficits: Midportant Achilles Tendinopathy Revision 2018.. *J Orthop Sports Phys Ther*, 2018 May 2;48(5). PMID 29712543
22. Shibuya N, Thorud JC, Humphers JM et al. Is percutaneous radiofrequency coblation for treatment of Achilles tendinosis safe and effective?. *J Foot Ankle Surg*, 2012 Sep 15;51(6). PMID 22974813
23. Lu Y, Zhang Q, Zhu Y et al. Is radiofrequency treatment effective for shoulder impingement syndrome? A prospective randomized controlled study.. *J Shoulder Elbow Surg*, 2013 Sep 3;22(11). PMID 23994459
24. Al-Ani Z, Jacobsen EW, Kartus JT et al. Radiofrequency microtenotomy: a promising method for treatment of rotator cuff tendinopathy.. *Knee Surg Sports Traumatol Arthrosc*, 2019 Sep 2. PMID 31473769
25. Owens BD, Stickles BJ, Balikian P et al. Prospective analysis of radiofrequency versus mechanical debridement of isolated patellar chondral lesions.. *Arthroscopy*, 2002 Feb 7;18(2). PMID 11830808
26. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination for Thermal Intradiscal Procedures (TIPs) (150.11). 2009; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=324>. Accessed November 3, 2020.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

