

EFFECTIVE DATE: 04|01|2023

POLICY LAST UPDATED: 3|31|2023

OVERVIEW

Laboratory tests have been developed that detect the expression of different genes in pigmented lesions or melanoma tumor tissue. Test results may help providers and patients decide whether to biopsy suspicious pigmented lesions, aid in diagnosis of lesions with indeterminate histopathologic findings or determine whether to perform sentinel lymph node biopsy in patients diagnosed with stage I or II cutaneous melanoma.

The following tests are addressed in this policy:

- DecisionDx-Melanoma (Castle Biosciences)
- Pigmented Lesion Assay (DermTech)
- myPath Melanoma (Castle Biosciences)

MEDICAL CRITERIA

Medicare Advantage Plans

DecisionDx Melanoma - 81529

DecisionDx Melanoma may be considered medically necessary when the following criteria are met:

- Patients diagnosed with cutaneous melanoma tumors with clinically negative sentinel node basins who are being considered for SLNB to determine eligibility for adjuvant therapy, and 1 or 2 below:
 1. Cutaneous melanoma ≥ 0.3 mm in Breslow thickness **without distant metastases** where additional information beyond anatomic and pathologic staging:
 - Will influence the decision to perform Sentinel Lymph Node Biopsy (T1-T2 only)
 - Will aid in the determination of appropriate adjuvant therapy
 2. Cutaneous melanoma < 0.3 mm in Breslow thickness being considered for sentinel lymph node biopsy:
 - in whom there is significant uncertainty about the adequacy of microstaging (positive deep margin), or
 - with other adverse features (e.g. very high mitotic index $\geq 2/\text{mm}^2$), lymphovascular invasion, or combination of these factors)

Pigmented Lesion Assay (PLA) – 0089U

The PLA may be considered medically necessary when all the following criteria are met:

- Melanocytic skin lesions with one or more clinical or historical characteristics suggestive of melanoma, including one or more ABCDE criteria (outlined below) when a clinician trained in the clinical diagnosis of skin cancer is considering the need for biopsy to rule out melanoma:
 - Asymmetry
 - Border
 - Color
 - Diameter
 - Evolving
- Primary melanocytic skin lesions between 5mm and 19mm
- Lesions where the skin is intact (i.e. non-ulcerated or non-bleeding lesions)
- Lesions that do not contain a scar or were previously biopsied

- Lesions not located in areas of psoriasis, eczema or similar skin conditions
- Lesions not clinically diagnosed as melanoma
- Lesions in areas other than palms of hands, soles of feet, nails, mucous membranes and hair covered areas that cannot be trimmed

myPath Melanoma – 0090U

myPath Melanoma may be considered medically necessary when the following criteria are met:

- The test is ordered by a board-certified dermatopathologist and;
- The lesion is considered to be a non-metastatic, melanocytic lesion that has not been previously treated, and;
- Testing is an adjunct to histopathology, as a clear distinction between a benign or malignant lesion cannot be made using clinical characteristics and histopathological features alone, and;
- The patient may be subjected to additional intervention, such as re-excision and/or sentinel lymph node biopsy, as a result of the diagnostic uncertainty, and;
- The results of the gene expression testing will be used in conjunction with the clinical evaluation, histopathological features and other diagnostic procedures to determine and/or alter the treatment plan.

Commercial Products

Not applicable

PRIOR AUTHORIZATION

Medicare Advantage Plans

Prior authorization is required for the following tests:

- DecisionDx-Melanoma
- Pigmented Lesion Assay
- myPath Melanoma

Note: Laboratories are not allowed to obtain clinical authorization or participate in the authorization process on behalf of the ordering physician. Only the ordering physician shall be involved in the authorization, appeal or other administrative processes related to prior authorization/medical necessity.

In no circumstance shall a laboratory or a physician/provider use a representative of a laboratory or anyone with a relationship to a laboratory and/or a third party to obtain authorization on behalf of the ordering physician, to facilitate any portion of the authorization process or any subsequent appeal of a claim where the authorization process was not followed and/or a denial for clinical appropriateness was issued, including any element of the preparation of necessary documentation of clinical appropriateness. If a laboratory or a third party is found to be supporting any portion of the authorization process, BCBSRI will deem the action a violation of this policy and severe action will be taken up to and including termination from the BCBSRI provider network. If a laboratory provides a laboratory service that has not been authorized, the service will be denied as the financial liability of the participating laboratory and may not be billed to the member.

POLICY STATEMENT

Medicare Advantage Plans

The following tests may be considered medically necessary when the medical criteria above are met:

- DecisionDx-Melanoma
- Pigmented Lesion Assay
- myPath Melanoma

Commercial Products

The following tests are not medically necessary as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome:

- DecisionDx-Melanoma
- Pigmented Lesion Assay
- myPath Melanoma

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable laboratory and not medically necessary/not covered benefits/coverage.

BACKGROUND

CUTANEOUS MELANOMA

Cutaneous melanoma accounts for more than 90% of cases of melanoma. For many decades, melanoma incidence was rapidly increasing in the United States. However, recent estimates have suggested the rise may be slowing. In 2018, more than 90,000 new cases of melanoma are expected to be diagnosed and more than 9,000 people are expected to die of melanoma.

Risk Factors

Exposure to solar ultraviolet radiation is a major risk factor for melanoma. Most melanomas occur on sun-exposed skin, particularly those areas most susceptible to sunburn. Likewise, features that are associated with an individual's sensitivity to sunlight, such as light skin pigmentation, red or blond hair, blue or green eyes, freckling tendency, and poor tanning ability are well-known risk factors for melanoma. There is also a strong association between high total body nevus counts and melanoma.

Several genes appear to contribute to melanoma predisposition such as tumor suppressor gene CDKN2A, melanocortin-1 receptor (MC1R) gene, and BAP1 variants. Individuals with either familial or sporadic melanoma have a 2 to 3 times increased risk of developing a subsequent primary melanoma. Several occupational exposures and lifestyle factors, such as body mass index and smoking, have been evaluated as possible risk factors for melanoma.

Gene Expression Profiling (GEP)

GEP measures the activity of thousands of genes simultaneously and creates a snapshot of cellular function. Data for GEP are generated by several molecular technologies including DNA microarrays that measure activity relative to previously identified genes and RNA-Seq that directly sequences and quantifies RNA molecules. Clinical applications of GEP include disease diagnosis, disease classification, prediction of drug response and prognosis.

DecisionDx-Melanoma

The DecisionDx test measures expression of 31 genes using quantitative reverse-transcription polymerase chain reaction. The test includes 28 prognostic gene targets and 3 endogenous control genes. The test is performed on standard tissue sections from an existing formalin-fixed, paraffin-embedded biopsy or wide local excision specimen. The DecisionDx test report provides a 'class' which stratifies tumors as class 1 or class 2. According to the sample report available on the manufacturer website: "The DecisionDx-Melanoma algorithm generates a value between 0 and 1 with a crossover point of 0.5. Subclassification (A or B) is based on proximity of this value to the crossover point."

Pigmented Lesion Assay (PLA)

The Pigmented Lesion Assay test measures expression of 6 genes (PRAME, LINC00518, CMIP, B2M, ACTB, PPIA). The PRAME (PReferentially expressed Antigenin MELanoma) gene encodes an antigen that is preferentially expressed in human melanomas, and that is not expressed in normal tissues (except testis). LINC00518 (Long Intergenic Non-protein Coding RNA518) is a regulatory RNA molecule. The other 4 genes provide normalization values.

The test is performed on skin samples of lesions at least 5 mm in diameter obtained via noninvasive, proprietary adhesive patch biopsies of a stratum corneum specimen. The test does not work on the palms of hands, soles of feet, nails, or mucous membranes, and it should not be used on bleeding or ulcerated lesions.

The Pigmented Lesion Assay test report includes 2 results. The first result is called the PLA MAGE (Melanoma Associated Gene Expression), which indicates low-risk (neither PRAME nor LINC00518 expression was detected), moderate-risk (expression of either PRAME or LINC00518 was detected), or high-risk (expression of both PRAME and LINC00518 was detected). The second result is as an algorithmic Pigmented Lesion Assay score that ranges from 0 to 100, with higher scores indicating higher suspicion of malignant disease.

myPath Melanoma

The myPath test measures expression of 23 genes using quantitative reverse-transcription polymerase chain reaction. Fourteen genes are involved in melanoma pathogenesis and are grouped into 3 components related to cell differentiation, cell signaling, and the immune response, and 9 housekeeper genes are also included. The test is performed on 5 standard tissue sections from an existing formalin-fixed, paraffin-embedded biopsy specimen.

The myPath test report includes an algorithmic myPath score ranging from -16.7 to 11.1, with higher, positive scores indicating higher suspicion of malignant disease. The myPath report also classifies these scores: -16.7 to -2.1 are “benign”; -2.0 to -0.1 are “indeterminate”; and 0.0 to +11.1 are “malignant”. The myPath test is meant as an add-on test to standard histopathology.

Commercial Products

For individuals with American Joint Committee on Cancer (AJCC) stage I or II cutaneous melanoma who receive GEP with the DecisionDx-Melanoma test to inform management decisions regarding enhanced surveillance, the evidence includes retrospective and prospective observational studies. Relevant outcomes are overall survival, disease-specific survival, test validity, change in disease status, resource utilization and treatment-related morbidity. The DecisionDx-Melanoma test has three independent clinical validity studies that have reported five-year recurrence-free survival (RFS) in AJCC stage I or II patients. Gerami et al (2015) reported RFS rates of 37% for DecisionDx class 2 (high-risk) in patients in AJCC stage I and II patients combined. Zager et al (2018) reported RFS rates of 85% (95% confidence interval [CI], 74% to 97%) for DecisionDx class 2 patients in AJCC stage 1 and 55% (95% CI, 44% to 69%) for DecisionDx class 2 in AJCC stage II disease. RFS does not appear to be well-characterized as evidenced by the variation in estimates across studies. This indication is to 'rule-in' patients for enhanced surveillance; therefore, specificity and positive predictive value (PPV) are key performance characteristics. Zager et al (2018) and Greenhaw et al (2018) the specificities were 71% and 87% respectively while the PPV were 48% and 24%, respectively. The PPV suggests that the majority of patients identified as high-risk by the DecisionDx test would not develop metastasis and would be unnecessarily subjected to additional surveillance. Greenhaw et al (2018) also reported that in 219 AJCC stage I patients, 201 had DecisionDx class 1 (low-risk) scores and 18 had DecisionDx class 2 (high-risk) scores. The only metastasis in stage I patients occurred in a patient with a DecisionDx class 1 score. Therefore none of their stage 1 patients benefited from DecisionDx testing but 18 (8%) were incorrectly identified as high-risk for metastasis and could have received unnecessary surveillance. Five-year RFS data are not available for the subgroup of patients for whom a 'rule-out' test would be relevant (class IIB through III). There is no evidence that changes to the frequency and methods for surveillance improve outcomes. Given that the evidence is insufficient to demonstrate test performance and there is no evidence that changes in surveillance improve outcomes, no inferences can be made about clinical utility through a chain of evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with AJCC stage I or II cutaneous melanoma who receive GEP with the DecisionDx-Melanoma test to inform management decisions regarding adjuvant therapy, the evidence includes retrospective and prospective observational studies. Relevant outcomes are overall survival, disease-specific survival, test validity, change in disease status, resource utilization and treatment-related morbidity. The

DecisionDx-Melanoma test has three independent clinical validity studies that have reported five-year RFS in AJCC stage I or II patients. Gerami et al (2015) reported RFS rates of 37% for DecisionDx class 2 (high-risk) in patients in AJCC stage I and II patients combined. Zager et al (2018) reported RFS rates of 85% (95% CI, 74% to 97%) for DecisionDx class 2 patients in AJCC stage 1 and 55% (95% CI, 44% to 69%) for DecisionDx class 2 in AJCC stage II disease. RFS does not appear to be well-characterized as evidenced by the variation in estimates across studies. This indication is to 'rule-in' patients for adjuvant therapy; therefore, specificity and PPV are key performance characteristics. Zager et al (2018) and Greenhaw et al (2018) the specificities were 71% and 87% respectively while the PPV were 48% and 24%, respectively. The PPV suggests that the majority of patients identified as high-risk by the DecisionDx test would not develop metastasis and would be unnecessarily subjected to additional treatment. Greenhaw et al (2018) also reported that in 219 AJCC stage I patients, 201 had DecisionDx class 1 (low-risk) scores and 18 had DecisionDx class 2 (high-risk) scores. The only metastasis in stage I patients occurred in a patient with a DecisionDx class 1 score. Therefore none of their stage 1 patients benefited from DecisionDx testing but 18 (8%) were incorrectly identified as high-risk for metastasis and could have received unnecessary treatment. There is no evidence that adjuvant therapy improves outcomes in these patients. Given that the evidence is insufficient to demonstrate test performance and there is no evidence that adjuvant therapy improves outcomes, no inferences can be made about clinical utility through a chain of evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with stage I or II cutaneous melanoma with clinically negative sentinel node basins who are being considered for sentinel lymph node (SLN) biopsy who receive GEP with the DecisionDx-Melanoma test to determine whether to perform SLN biopsy, the evidence includes retrospective observational studies. Relevant outcomes are overall survival, disease-specific survival, test validity, change in disease status, resource utilization, and treatment-related morbidity. The DecisionDx-Melanoma test has 3 independent clinical validity studies that have reported 5-year RFS in AJCC stage I or II patients. Gerami et al (2015) reported RFS rates of 98% in DecisionDx class 1 (low-risk) without CIs, in AJCC stage I or II patients. Zager et al (2017) reported RFS rates of 96% (95% CI, 94% to 99%) for DecisionDx class 1 in patients with AJCC stage I disease; they also reported RFS rates of 74% (95% CI, 60% to 91%) for DecisionDx class 1 in patients with AJCC stage II disease. Although CIs were not available for the first study, RFS does not appear to be well-characterized as evidenced by the variation in estimates across studies. Zager et al (2017) also reported that in 56 patients who were DecisionDx class 1 (low-risk) but SLN biopsy-positive, 22 recurrences (39%) occurred over 5 years. If the DecisionDx test were used as a triage for SLN biopsy, these patients would not undergo SLN biopsy and would likely not receive adjuvant therapy, which has shown to be effective at prolonging the time to recurrence in node-positive patients. Data on 5-year RFS is not available for the target population (Class 1A patients ≤ 55 years old who have tumors ≤ 2 mm deep [T1 to T2]) outside of the retrospective cohort that was used to identify the target population. No direct evidence of clinical utility was identified. Given that the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility through a chain of evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with suspicious pigmented lesions (based on ABCDE and/or ugly duckling criteria) being considered for biopsy who receive gene expression profiling with the DermTech Pigmented Lesion Assay to determine which lesions should proceed to biopsy, the evidence includes observational studies. Relevant outcomes are overall survival, disease-specific survival, validity, and resource utilization. The Pigmented Lesion Assay has 1 clinical validity study with many methodologic and reporting limitations. Therefore, performance characteristics are not well-characterized. Also, the test has not been compared with dermoscopy, another tool frequently used to make biopsy decisions. No direct evidence of clinical utility was identified. Given that the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility through a chain of evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have melanocytic lesions with indeterminate histopathologic features who receive gene expression profiling with the myPath Melanoma test added to histopathology to aid in the diagnosis of melanoma, the evidence includes observational studies. Relevant outcomes are overall survival, disease-

specific survival, test validity, change in disease status, and treatment-related morbidity. The myPath test has 2 clinical validity studies including long-term follow-up for metastasis as the reference standard. In 1 study, it is not clear whether the study population included lesions that were indeterminate following histopathology. The second study focused on indeterminate lesions but had limitations including a retrospective design and less than 5-year follow-up in 31% of cases. Therefore, performance characteristics are not well-characterized. No direct evidence of clinical utility was identified. Given that the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility through a chain of evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

CODING

The following CPT codes are covered for Medicare Advantage Plans when medical criteria above are met and are not medically necessary for Commercial Products.

This code can be used for DecisionDx-Melanoma:

81529 Oncology (cutaneous melanoma), mRNA, gene expression profiling by real-time RT-PCR of 31 genes (28 content and 3 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk, including likelihood of sentinel lymph node metastasis

This code can be used for Pigmented Lesion Assay (PLA):

0089U Oncology (melanoma), gene expression profiling by RTqPCR, PRAME and LINC00518, superficial collection using adhesive patch(es)

This code can be used for myPath Melanoma:

0090U Oncology (cutaneous melanoma), mRNA gene expression profiling by RT-PCR of 23 genes (14 content and 9 housekeeping), utilizing formalin-fixed paraffin-embedded (FFPE) tissue, algorithm reported as a categorical result (ie, benign, intermediate, malignant)

RELATED POLICIES

Genetic Testing Services

Proprietary Laboratory Analyses (PLA)

PUBLISHED

Provider Update, February 2023

Provider Update, October 2021

Provider Update, November 2020

Provider Update, April 2019

Provider Update, August 2018

REFERENCES

1. Centers for Medicare and Medicaid Services. Local Coverage Determination (LCD): MolDX: Pigmented Lesion Assay (L38151)
2. Centers for Medicare and Medicaid Services. Local Coverage Article: Billing and Coding: MolDX: Pigmented Lesion Assay (A58052)
3. Chang AE, Karnell LH, Menck HR. The National Cancer Data Base report on cutaneous and noncutaneous melanoma: a summary of 84,836 cases from the past decade. The American College of Surgeons Commission on Cancer and the American Cancer Society. *Cancer*. Oct 15 1998; 83(8): 1664-78. PMID 9781962
4. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin*. Jan 2018; 68(1): 7-30. PMID 29313949
5. Gilchrest BA, Eller MS, Geller AC, et al. The pathogenesis of melanoma induced by ultraviolet radiation. *N Engl J Med*. Apr 29 1999; 340(17): 1341-8. PMID 10219070
6. Gandini S, Sera F, Cattaruzza MS, et al. Meta-analysis of risk factors for cutaneous melanoma: III. Family history, actinic damage and phenotypic factors. *Eur J Cancer*. Sep 2005; 41(14): 2040-59. PMID 16125929

7. Caini S, Gandini S, Sera F, et al. Meta-analysis of risk factors for cutaneous melanoma according to anatomical site and clinico-pathological variant. *Eur J Cancer*. Nov 2009; 45(17): 3054-63. PMID 19545997
8. Goldstein AM, Chan M, Harland M, et al. Features associated with germline CDKN2A mutations: a GenoMEL study of melanoma-prone families from three continents. *J Med Genet*. Feb 2007; 44(2): 99-106. PMID 16905682
9. Wendt J, Rauscher S, Burgstaller-Muehlbacher S, et al. Human Determinants and the Role of Melanocortin-1 Receptor Variants in Melanoma Risk Independent of UV Radiation Exposure. *JAMA Dermatol*. Jul 01 2016; 152(7):776-82. PMID 27050141
10. Wiesner T, Obenaus AC, Murali R, et al. Germline mutations in BAP1 predispose to melanocytic tumors. *NatGenet*. Aug 28 2011; 43(10): 1018-21. PMID 21874003
11. Chen T, Fallah M, Forsti A, et al. Risk of Next Melanoma in Patients With Familial and Sporadic Melanoma by Number of Previous Melanomas. *JAMA Dermatol*. Jun 2015; 151(6): 607-15. PMID 25671687
12. Jiang AJ, Rambhatla PV, Eide MJ. Socioeconomic and lifestyle factors and melanoma: a systematic review. *Br J Dermatol*. Apr 2015; 172(4): 885-915. PMID 25354495
13. Abbasi NR, Shaw HM, Rigel DS, et al. Early diagnosis of cutaneous melanoma: revisiting the ABCD criteria. *JAMA*. Dec 08 2004; 292(22): 2771-6. PMID 15585738
14. Grob JJ, Bonerandi JJ. The 'ugly duckling' sign: identification of the common characteristics of nevi in an individual as a basis for melanoma screening. *Arch Dermatol*. Jan 1998; 134(1): 103-4. PMID 9449921
15. Wilson RL, Yentzer BA, Isom SP, et al. How good are US dermatologists at discriminating skin cancers? A number-needed-to-treat analysis. *J Dermatolog Treat*. Feb 2012; 23(1): 65-9. PMID 21756146
16. National Center for Biotechnology Information. PRAME preferentially expressed antigen in melanoma. 2021; <https://www.ncbi.nlm.nih.gov/gene/23532>. Accessed April 8, 2022.
17. DermTech. Pigmented Lesion Assay: Non-invasive gene expression analysis of pigmented skin lesions. Performance and Development Notes. 2015; <http://dermtech.com/wp-content/uploads/2015/10/White-Paper-DermTech-Melanoma-Assay-.pdf>. Accessed April 8, 2022.
18. Wachsman W, Morhenn V, Palmer T, et al. Noninvasive genomic detection of melanoma. *Br J Dermatol*. Apr 2011; 164(4): 797-806. PMID 21294715
19. Gerami P, Alsobrook JP, Palmer TJ, et al. Development of a novel noninvasive adhesive patch test for the evaluation of pigmented lesions of the skin. *J Am Acad Dermatol*. Aug 2014; 71(2): 237-44. PMID 24906614
20. Gerami P, Yao Z, Polsky D, et al. Development and validation of a noninvasive 2-gene molecular assay for cutaneous melanoma. *J Am Acad Dermatol*. Jan 2017; 76(1): 114-120.e2. PMID 27707590
21. Vestergaard ME, Macaskill P, Holt PE, et al. Dermoscopy compared with naked eye examination for the diagnosis of primary melanoma: a meta-analysis of studies performed in a clinical setting. *Br J Dermatol*. Sep 2008; 159(3):669-76. PMID 18616769
22. Murzaku EC, Hayan S, Rao BK. Methods and rates of dermoscopy usage: a cross-sectional survey of US dermatologists stratified by years in practice. *J Am Acad Dermatol*. Aug 2014; 71(2): 393-5. PMID 25037790
23. Engasser HC, Warshaw EM. Dermoscopy use by US dermatologists: a cross-sectional survey. *J Am Acad Dermatol*. Sep 2010; 63(3): 412-9, 419.e1-2. PMID 20619490
24. Bossuyt PM, Irwig L, Craig J, et al. Comparative accuracy: assessing new tests against existing diagnostic pathways. *BMJ*. May 06 2006; 332(7549): 1089-92. PMID 16675820
25. Ferris LK, Gerami P, Skelsey MK, et al. Real-world performance and utility of a noninvasive gene expression assay to evaluate melanoma risk in pigmented lesions. *Melanoma Res*. Oct 2018; 28(5): 478-482. PMID 30004988
26. Ferris LK, Jansen B, Ho J, et al. Utility of a Noninvasive 2-Gene Molecular Assay for Cutaneous Melanoma and Effect on the Decision to Biopsy. *JAMA Dermatol*. Jul 01 2017; 153(7): 675-680. PMID 28445578
27. Myriad. n.d. Understanding the myPath Melanoma Results; <https://mypathmelanoma.com/about-mypath-melanoma/understanding-the-mypath-melanoma-results/>. Accessed April 8, 2022
28. Clarke LE, Warf MB, Flake DD, et al. Clinical validation of a gene expression signature that differentiates benign nevi from malignant melanoma. *J Cutan Pathol*. Apr 2015; 42(4): 244-52. PMID 25727210

29. Clarke LE, Flake DD, Busam K, et al. An independent validation of a gene expression signature to differentiate malignant melanoma from benign melanocytic nevi. *Cancer*. Feb 15 2017; 123(4): 617-628. PMID 27768230
30. Reimann JDR, Salim S, Velazquez EF, et al. Comparison of melanoma gene expression score with histopathology, fluorescence in situ hybridization, and SNP array for the classification of melanocytic neoplasms. *Mod Pathol*. Nov2018; 31(11): 1733-1743. PMID 29955141
31. Gaiser T, Kutzner H, Palmedo G, et al. Classifying ambiguous melanocytic lesions with FISH and correlation with clinical long-term follow up. *Mod Pathol*. Mar 2010; 23(3): 413-9. PMID 20081813
32. Vergier B, Prochazkova-Carlotti M, de la Fouchardiere A, et al. Fluorescence in situ hybridization, a diagnostic aid in ambiguous melanocytic tumors: European study of 113 cases. *Mod Pathol*. May 2011; 24(5): 613-23. PMID21151100
33. Ko JS, Clarke LE, Minca EC, et al. Correlation of melanoma gene expression score with clinical outcomes on a series of melanocytic lesions. *Hum Pathol*. Apr 2019; 86: 213-221. PMID 30566894
34. Clarke LE, Pimentel JD, Zalaznick H, et al. Gene expression signature as an ancillary method in the diagnosis of desmoplastic melanoma. *Hum Pathol*. Dec 2017; 70: 113-120. PMID 29079183
35. Minca EC, Al-Rohil RN, Wang M, et al. Comparison between melanoma gene expression score and fluorescence in situ hybridization for the classification of melanocytic lesions. *Mod Pathol*. Aug 2016; 29(8): 832-43. PMID27174586
36. Ko JS, Matharoo-Ball B, Billings SD, et al. Diagnostic Distinction of Malignant Melanoma and Benign Nevi by a Gene Expression Signature and Correlation to Clinical Outcomes. *Cancer Epidemiol Biomarkers Prev*. Jul 2017;26(7): 1107-1113. PMID 28377414
37. Clarke LE, Mabey B, Flake Li DD, et al. Clinical validity of a gene expression signature in diagnostically uncertain neoplasms. *Per Med*. Sep 2020; 17(5): 361-371. PMID 32915688
38. Cockerell C, Tschen J, Billings SD, et al. The influence of a gene-expression signature on the treatment of diagnostically challenging melanocytic lesions. *Per Med*. Mar 2017; 14(2): 123-130. PMID 28757886
39. Cockerell CJ, Tschen J, Evans B, et al. The influence of a gene expression signature on the diagnosis and recommended treatment of melanocytic tumors by dermatopathologists. *Medicine (Baltimore)*. Oct 2016; 95(40):e4887. PMID 27749545
40. Gershenwald JES, R.A.; Hess, K.R.; et al. *Melanoma of the Skin*. Chicago, IL: American Joint Committee on Cancer; 2017.
41. Eggermont AM, Chiarion-Sileni V, Grob JJ, et al. Prolonged Survival in Stage III Melanoma with Ipilimumab Adjuvant Therapy. *N Engl J Med*. Nov 10 2016; 375(19): 1845-1855. PMID 27717298
42. Weber J, Mandala M, Del Vecchio M, et al. Adjuvant Nivolumab versus Ipilimumab in Resected Stage III or IV Melanoma. *N Engl J Med*. Nov 09 2017; 377(19): 1824-1835. PMID 28891423
43. Long GV, Hauschild A, Santinami M, et al. Adjuvant Dabrafenib plus Trametinib in Stage III BRAF-Mutated Melanoma. *N Engl J Med*. Nov 09 2017; 377(19): 1813-1823. PMID 28891408
44. Gerami P, Cook RW, Wilkinson J, et al. Development of a prognostic genetic signature to predict the metastatic risk associated with cutaneous melanoma. *Clin Cancer Res*. Jan 01 2015; 21(1): 175-83. PMID 25564571
45. Zager JS, Gastman BR, Leachman S, et al. Performance of a prognostic 31-gene expression profile in an independent cohort of 523 cutaneous melanoma patients. *BMC Cancer*. Feb 05 2018; 18(1): 130. PMID 29402264
46. Wrightson WR, Wong SL, Edwards MJ, et al. Complications associated with sentinel lymph node biopsy for melanoma. *Ann Surg Oncol*. Jul 2003; 10(6): 676-80. PMID 12839853
47. Soong SJ, Ding S, Coit DG, et al. AJCC: Individualized melanoma patient outcome prediction tools. n.d.;<http://www.melanomaprognosis.net/>. Accessed April 8, 2022.
48. Callender GG, Gershenwald JE, Egger ME, et al. A novel and accurate computer model of melanoma prognosis for patients staged by sentinel lymph node biopsy: comparison with the American Joint Committee on Cancer model. *J Am Coll Surg*. Apr 2012; 214(4): 608-17; discussion 617-9. PMID 22342785
49. Dicker TJ, Kavanagh GM, Herd RM, et al. A rational approach to melanoma follow-up in patients with primary cutaneous melanoma. Scottish Melanoma Group. *Br J Dermatol*. Feb 1999; 140(2): 249-54. PMID 10233217

50. Garbe C, Paul A, Kohler-Spath H, et al. Prospective evaluation of a follow-up schedule in cutaneous melanoma patients: recommendations for an effective follow-up strategy. *J Clin Oncol.* Feb 01 2003; 21(3): 520-9. PMID12560444
51. Faries MB, Steen S, Ye X, et al. Late recurrence in melanoma: clinical implications of lost dormancy. *J Am CollSurg.* Jul 2013; 217(1): 27-34; discussion 34-6. PMID 23643694
52. Hsueh EC, DeBloom JR, Lee J, et al. Interim analysis of survival in a prospective, multi-center registry cohort of cutaneous melanoma tested with a prognostic 31-gene expression profile test. *J Hematol Oncol.* Aug 29 2017;10(1): 152. PMID 28851416
53. Podlipnik S, Carrera C, Boada A, et al. Early outcome of a 31-gene expression profile test in 86 AJCC stage IB-II melanoma patients. A prospective multicentre cohort study. *J Eur Acad Dermatol Venereol.* May 2019; 33(5): 857-862. PMID 30702163
54. Hsueh EC, DeBloom JR, Lee JH, et al. Long-Term Outcomes in a Multicenter, Prospective Cohort Evaluating the Prognostic 31-Gene Expression Profile for Cutaneous Melanoma. *JCO Precis Oncol.* 2021; 5. PMID 34036233
55. Gastman BR, Gerami P, Kurley SJ, et al. Identification of patients at risk of metastasis using a prognostic 31-gene expression profile in subpopulations of melanoma patients with favorable outcomes by standard criteria. *J Am Acad Dermatol.* Jan 2019; 80(1): 149-157.e4. PMID 30081113
56. Gastman BR, Zager JS, Messina JL, et al. Performance of a 31-gene expression profile test in cutaneous melanomas of the head and neck. *Head Neck.* Apr 2019; 41(4): 871-879. PMID 30694001
57. Vetto JT, Hsueh EC, Gastman BR, et al. Guidance of sentinel lymph node biopsy decisions in patients with T1-T2melanoma using gene expression profiling. *Future Oncol.* Apr 2019; 15(11): 1207-1217. PMID 30691297
58. Marks, Etan et al. Establishing an evidence-based decision point for clinical use of the 31-gene expression profile test in cutaneous melanoma. *SKIN The Journal of Cutaneous Medicine, [S.l.],* July 2019, v. 3, n. 4, p. 239-249.
59. Greenhaw BN, Zitelli JA, Brodland DG. Estimation of Prognosis in Invasive Cutaneous Melanoma: An Independent Study of the Accuracy of a Gene Expression Profile Test. *Dermatol Surg.* Dec 2018; 44(12): 1494-1500. PMID29994951
60. Keller J, Schwartz TL, Lizalek JM, et al. Prospective validation of the prognostic 31-gene expression profiling test in primary cutaneous melanoma. *Cancer Med.* May 2019; 8(5): 2205-2212. PMID 30950242
61. Gerami P, Cook RW, Russell MC, et al. Gene expression profiling for molecular staging of cutaneous melanoma inpatients undergoing sentinel lymph node biopsy. *J Am Acad Dermatol.* May 2015; 72(5): 780-5.e3. PMID25748297
62. Ferris LK, Farberg AS, Middlebrook B, et al. Identification of high-risk cutaneous melanoma tumors is improved when combining the online American Joint Committee on Cancer Individualized Melanoma Patient Outcome Prediction Tool with a 31-gene expression profile-based classification. *J Am Acad Dermatol.* May 2017; 76(5): 818-825.e3. PMID 28110997
63. Berger AC, Davidson RS, Poitras JK, et al. Clinical impact of a 31-gene expression profile test for cutaneous melanoma in 156 prospectively and consecutively tested patients. *Curr Med Res Opin.* Sep 2016; 32(9): 1599-604.PMID 27210115
64. Farberg AS, Glazer AM, White R, et al. Impact of a 31-gene Expression Profiling Test for Cutaneous Melanoma on Dermatologists' Clinical Management Decisions. *J Drugs Dermatol.* May 01 2017; 16(5): 428-431. PMID 28628677
65. Schuitevoerder D, Heath M, Cook RW, et al. Impact of Gene Expression Profiling on Decision-Making in Clinically Node Negative Melanoma Patients after Surgical Staging. *J Drugs Dermatol.* Feb 01 2018; 17(2): 196-199. PMID29462228
66. Dillon LD, Gadzia JE, Davidson RS, et al. Prospective, multicenter clinical impact evaluation of a 31-geneexpression profile test for management of melanoma patients. *Skin.* 2018;2(2):111-121.
67. Hyams DM, Covington KR, Johnson CE, et al. Integrating the melanoma 31-gene expression profile test with surgical oncology practice within national guideline and staging recommendations. *Future Oncol.* Feb 2021; 17(5):517-527. PMID 33021104
68. Dillon LD, McPhee M, Davidson RS, et al. Expanded evidence that the 31-gene expression profile test provides clinical utility for melanoma management in a multicenter study. *Curr Med Res Opin.* Feb 15 2022: 1-8. PMID35081854

69. National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology. Cutaneous Melanoma. Version 2.2022.
https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf. Accessed April 8, 2022.
70. Swetter SM, Tsao H, Bichakjian CK, et al. Guidelines of care for the management of primary cutaneous melanoma. *J Am Acad Dermatol*. Jan 2019; 80(1): 208-250. PMID 30392755
71. American Academy of Dermatology. Choosing Wisely. 2019. <https://www.choosingwisely.org/clinician-lists/american-academy-dermatology-sentinel-lymph-node-biopsy-early-melanoma-evaluation/>. Accessed April 8, 2022.
72. Warf MB, Flake DD, Adams D, Gutin A, et al. Analytical validation of a melanoma diagnostic gene signature using formalin-fixed paraffin-embedded melanocytic lesions. *Biomark Med*. 2015;9(5):407-16.
73. Leachman SA, Koon SM, Korcheva VB, White KP. Assessing Genetic Expression Profiles in Melanoma Diagnosis. *Dermatol Clin* 2017;35: 537–544.
74. Cassarino DS, Lewine N, Cole D, Wade B, Gustavsen G. Budget impact analysis of a novel gene expression assay for the diagnosis of malignant melanoma. *Journal of medical economics*. 2014;17(11):782-791.
75. National Comprehensive Cancer Network (NCCN) – Melanoma v2.2021
76. Hayes Genetic Testing Evaluation (GTE) Synopsis: DecisionDx- Melanoma
77. Renzetti M, Farma J, Handor B et al. Combined experience of two tertiary referral centers with DecisionDx-Melanoma GEP testing. In Society of Surgical Oncology Seattle, WA: 2017.
78. Fleming MD, Johnson C, Covington KR et al. Clinical impact of a 31-gene expression profile test on physician recommendations for management of melanoma patients in a prospectively tested cohort. In Society of Melanoma Research. Brisbane, Australia: 2017
79. Moody JA, Ali RF, Carbone AC et al. Complications of sentinel lymph node biopsy for melanoma - A systematic review of the literature. *Eur J Surg Oncol* 2017; 43: 270-277.
80. Wong SL, Faries MB, Kennedy EB et al. Sentinel Lymph Node Biopsy and Management of Regional Lymph Nodes in Melanoma: American Society of Clinical Oncology and Society of Surgical Oncology Clinical Practice Guideline Update. *J Clin Oncol* 2017;
81. Svoboda RM, Glazer AM, et al. Factors affecting dermatologists' use of a 31-gene expression profiling test as an adjunct for predicting metastatic risk in cutaneous melanoma. *J Drugs Dermatol*. 2018;17(5):544-547.
82. Mirsky RS, Prado G, et al. management decisions made by physician assistants and nurse practitioners in cutaneous malignant melanoma patients: Impact of a 31-gene expression profile test. *J Drugs Dermatol*. 2018;17(11):1220-1223.
83. Cook RW, Middlebrook B, et al. Analytic validity of DecisionDx-Melanoma, a gene expression profile test for determining metastatic risk in melanoma patients. *Diagnostic Pathology* (2018) 13:13
84. NIH, National cancer Institute. Moles to Melanoma: Recognizing the ABCDE Features. <https://moles-melanomatool.cancer.gov/>
85. Yao Z, Moy R, Allen T, Jansen B. An Adhesive Patch-Based Skin Biopsy Device for Molecular Diagnostics and Skin Microbiome Studies. *J Drugs Dermatol*. 2017;16(10):979-986.
86. Kwatra S.G., H.H., Semenov Y.R., Trotter S.C., Holland E., Leachman S. A Dermatologist's Guide to Implementation of Gene Expression Profiling in the Management of Melanoma. *J Clin Aesthet Dermatol* 2020 13, S3-S14

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

