

**EFFECTIVE DATE:** 7|01|2026

**POLICY LAST REVIEWED:** 05|06|2026

## **OVERVIEW**

Microprocessor-controlled prostheses use feedback from sensors to adjust joint movement on a real-time as-needed basis. Active joint control is intended to improve safety and function, particularly for patients who can maneuver on uneven terrain and with variable gait.

## **MEDICAL CRITERIA**

### **Medicare Advantage Plans**

Blue Cross & Blue Shield of Rhode Island (BCBSRI) follows the medical necessity criteria from the Centers for Medicare and Medicaid Services (CMS) National and Local Coverage Determinations (NCD/LCD) for L5615. Please use the online tool for participating providers. See the Related Policies section.

### **Commercial Products**

The medical criteria below is only applicable for the following HCPCS code:

- L5615
- 

A microprocessor-controlled knee may be considered medically necessary in individuals with transfemoral amputation who meet the following requirements:

- demonstrated need for long-distance ambulation at variable rates (use of the limb in the home or for basic community ambulation is not sufficient to justify provision of the computerized limb over standard limb applications) OR demonstrated patient need for regular ambulation on uneven terrain or for regular use on stairs (use of the limb for limited stair climbing in the home or employment environment is not sufficient evidence for prescription of this device over standard prosthetic application); AND
- physical ability, including adequate cardiovascular and pulmonary reserve, for ambulation at faster than normal walking speed; AND
- adequate cognitive ability to master use and care requirements for the technology

Amputees should be evaluated by an independent, qualified professional to determine the most appropriate prosthetic components and control mechanism. A trial period may be indicated to evaluate the tolerability and efficacy of the prosthesis in a real-life setting. Decisions about the potential benefits of microprocessor knees involve multiple factors including activity levels and the patient's physical and cognitive ability. A patient's need for daily ambulation of at least 400 continuous yards, daily and frequent ambulation at variable cadence or on uneven terrain (eg, gravel, grass, curbs), and daily and frequent use of ramps and/or stairs (especially stair descent) should be considered as part of the decision. Typically, the daily and frequent need of 2 or more of these activities would be needed to show benefit.

### **Individual Selection and Identification**

For individuals in whom the potential benefits of the microprocessor knees are uncertain, individuals may first be fitted with a standard prosthesis to determine their level of function with the standard device.

- A. Contraindications for the use of the microprocessor knee should include the following:
- Any condition that prevents socket fitting, such as a complicated wound or intractable pain which precludes socket wear

- Inability to tolerate the weight of the prosthesis
- Medicare level K0-no ability or potential to ambulate or transfer
- Medicare level K1-limited ability to transfer or ambulate on level ground at fixed cadence
- Medicare level K2-limited community ambulator who does not have the cardiovascular reserve, strength, and balance to improve stability in stance to permit increased independence, less risk of falls, and potential to advance to a less restrictive walking device
- Inability to use swing and stance features of the knee unit
- Poor balance or ataxia that limits ambulation
- Significant hip flexion contracture (>20°)
- Significant deformity of remaining limb that would impair the ability to stride
- Limited cardiovascular and/or pulmonary reserve or profound weakness
- Limited cognitive ability to understand gait sequencing or care requirements
- Long-distance or competitive running
- Falls outside of recommended weight or height guidelines of the manufacturer
- Specific environmental factors such as excessive moisture or dust, or inability to charge the prosthesis
- Extremely rural conditions where maintenance ability is limited.

B. Indications for the use of the microprocessor knee should include the following:

- Adequate cardiovascular and pulmonary reserve to ambulate at variable cadence
- Adequate strength and balance in stride to activate the knee unit
- Should not exceed the weight or height restrictions of the device
- Adequate cognitive ability to master technology and gait requirements of the device
- Hemi-pelvectomy through knee-disarticulation level of amputation, including bilateral; lower-extremity amputees are candidates if they meet functional criteria as listed
- The individual is an active walker and requires a device that reduces energy consumption to permit longer distances with less fatigue
- Daily activities or job tasks that do not permit full focus of concentration on knee control and stability-such as uneven terrain, ramps, curbs, stairs, repetitive lifting, and/or carrying
- Medicare level K2-limited community ambulator, but only if improved stability in stance permits increased independence, less risk of falls, and potential to advance to a less restrictive walking device, and the individual has the cardiovascular reserve, strength, and balance to use the prosthesis. The microprocessor enables fine-tuning and adjustment of the hydraulic mechanism to accommodate the unique motor skills and demands of the functional level K2 ambulator.
- Medicare level K3-unlimited community ambulator
- Medicare level K4-active adult athlete who needs to function as a K3 level in daily activities
- Potential to lessen back pain by providing more secure stance control, using less muscle control to keep the knee stable
- Potential to unload and decrease stress on remaining limb
- Potential to return to an active lifestyle.

C. Physical and Functional Fitting Criteria for New Amputees:

- New amputees may be considered if they meet certain criteria as outlined above
- Premorbid and current functional assessment important determinant
- Requires stable wound and ability to fit the socket
- Immediate postoperative fit is possible
- Must have potential to return to an active lifestyle

## PRIOR AUTHORIZATION

Prior authorization for microprocessor-controlled prostheses (L5615) for lower limb is required for Medicare Advantage Plans and recommended for Commercial Products.

## **POLICY STATEMENT**

### **Medicare Advantage Plans**

A microprocessor-controlled knee in individuals with transfemoral amputation is considered medically necessary when the criteria in the online authorization tool is met.

A powered knee is considered medically necessary when the criteria in the online authorization tool is met.

### **Commercial Products**

A microprocessor-controlled knee in individuals with transfemoral amputation is considered medically necessary when the criteria above are met.

A powered knee is considered medically necessary when the criteria in the online authorization tool is met.

## **COVERAGE**

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

## **BACKGROUND**

### **Lower-Extremity Prosthetics**

More than 100 different prosthetic ankle-foot and knee designs are currently available. The choice of the most appropriate design may depend on the patient's underlying activity level. For example, the requirements of a prosthetic knee in elderly, largely homebound individual will differ from those of a younger, active person. Key elements of prosthetic knee design involve providing stability during both the stance and swing phase of the gait. Prosthetic knees vary in their ability to alter the cadence of the gait, or the ability to walk on rough or uneven surfaces. In contrast to more simple prostheses, which are designed to function optimally at 1 walking cadence, fluid and hydraulic-controlled devices are designed to allow amputees to vary their walking speed by matching the movement of the shin portion of the prosthesis to the movement of the upper leg. For example, the rate at which the knee flexes after "toe-off" and then extends before heel strike depends in part on the mechanical characteristics of the prosthetic knee joint. If the resistance to flexion and extension of the joint does not vary with gait speed, the prosthetic knee extends too quickly or too slowly relative to the heel strike if the cadence is altered. When properly controlled, hydraulic or pneumatic swing-phase controls allow the prosthetist to set a pace adjusted to the individual amputee, from very slow to a race-walking pace. Hydraulic prostheses are heavier than other options and require gait training; for these reasons, these prostheses are prescribed for athletic or fit individuals. Other design features include multiple centers of rotation, referred to as "polycentric knees." The mechanical complexity of these devices allows engineers to optimize selected stance and swing-phase features.

### **Regulatory Status**

According to the manufacturers, microprocessor-controlled prostheses are considered a class I device by the FDA and are exempt from 510(k) requirements. This classification does not require submission of clinical data regarding efficacy but only notification of FDA prior to marketing. FDA product codes: ISW, KFX.

For individuals who have a transfemoral amputation who receive a prosthesis with a microprocessor-controlled knee, the evidence includes a number of within-subject comparisons of microprocessor-controlled knees versus non-microprocessor-controlled knee joints and systematic reviews of these studies. Relevant outcomes are functional outcomes, health status measures, and quality of life. For K3- and K4-level amputees, studies have shown an objective improvement in function on some outcome measures, particularly for hill and ramp descent, and strong patient preference for microprocessor-controlled prosthetic knees. Benefits include a more normal gait, increased stability, and a decrease in falls. The evidence in Medicare level K2 ambulators suggests that a prosthesis with stance control only can improve activities that require balance

and improve walking in this population. For these reasons, a microprocessor-controlled knee may provide incremental benefit for these individuals. The potential to achieve a higher functional level with a microprocessor-controlled knee includes having the appropriate physical and cognitive ability to use the advanced technology. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a transfemoral amputation who receive a prosthesis with a powered knee, the evidence includes 2 small (n<10) within-subject comparisons. Relevant outcomes are functional outcomes, health status measures, and quality of life. The limited evidence available to date does not support an improvement in functional outcomes with powered knee prostheses compared with microprocessor-controlled knees in individuals with unilateral above-knee amputation. Additionally, no studies were identified that directly compared powered knee prostheses with standard prosthetic knees in patients with transfemoral amputation. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## **CODING**

### **Medicare Advantage Plans**

The following HCPCS code(s) are medically necessary for Medicare Advantage Plans when the medical criteria in the in the online authorization tool for participating providers is met:

**L5615** Addition, endoskeletal knee-shin system, 4 bar linkage or multiaxial, fluid swing and stance phase control

**L5827** Endoskeletal knee-shin system, single axis, electromechanical swing and stance phase control, with or without shock absorption and stance extension damping (New Code Effective 4/1/2025)

### **Commercial Products**

The following HCPCS code(s) is medically necessary for Commercial Products when the medical criteria above has been met:

**L5615** Addition, endoskeletal knee-shin system, 4 bar linkage or multiaxial, fluid swing and stance phase control

The following HCPCS code(s) are medically necessary for Medicare Advantage Plans when the medical criteria in the in the online authorization tool for participating providers is met:

**L5827** Endoskeletal knee-shin system, single axis, electromechanical swing and stance phase control, with or without shock absorption and stance extension damping (New Code Effective 4/1/2025)

## **RELATED POLICIES**

Prior Authorization via Web-Based Tool for Durable Medical Equipment (DME)

Medicare Advantage Plans National and Local Coverage Determinations

## **PUBLISHED**

Provider Update, June 2026

Provider Update, September 2025

Provider Update, June 2025

Provider Update, June 2024

Provider Update, May 2023

## **REFERENCES**

Flynn K. Short Report: Computerized lower limb prosthesis (VA Technology Assessment Program). No. 2. Boston, MA: Veterans Health Administration; 2000.

Thibaut A, Beaudart C, Maertens DE Noordhout B, et al. Impact of microprocessor prosthetic knee on mobility and quality of life in patients with lower limb amputation: a systematic review of the literature. Eur J Phys Rehabil Med. Jun 2022; 58(3): 452-461. PMID 35148043

Hahn A, Bueschges S, Prager M, et al. The effect of microprocessor controlled exo-prosthetic knees on limited community ambulators: systematic review and meta-analysis. *Disabil Rehabil.* Dec 2022; 44(24): 7349-7367. PMID 34694952

Morgan SJ, Friedly JL, Nelson IK, et al. The effects of microprocessor prosthetic knee use in early rehabilitation: A pilot randomized controlled trial. *PM R.* Apr 2025; 17(4): 371-383. PMID 39895150

Wurdeman SR, Hafner BJ, Sawers A, et al. ASsessing Clinical outcomes with microprocEssor kNee uTilization in a K2 population (ASCENT K2): randomized controlled trial results for above-knee prosthesis users over age 65. *Disabil Rehabil.* Mar 2026; 48(5): 1476-1493. PMID 40686456

Theeven P, Hemmen B, Rings F, et al. Functional added value of microprocessor-controlled knee joints in daily life performance of Medicare Functional Classification Level-2 amputees. *J Rehabil Med.* Oct 2011; 43(10): 906-15. PMID 21947182

Theeven PJ, Hemmen B, Geers RP, et al. Influence of advanced prosthetic knee joints on perceived performance and everyday life activity level of low-functional persons with a transfemoral amputation or knee disarticulation. *J Rehabil Med.* May 2012; 44(5): 454-61. PMID 22549656

Burnfield JM, Eberly VJ, Gronely JK, et al. Impact of stance phase microprocessor-controlled knee prosthesis on ramp negotiation and community walking function in K2 level transfemoral amputees. *Prosthet Orthot Int.* Mar 2012; 36(1): 95-104. PMID 22223685

Orendurff MS, Segal AD, Klute GK, et al. Gait efficiency using the C-Leg. *J Rehabil Res Dev.* 2006; 43(2): 239-46. PMID 16847790

Klute GK, Berge JS, Orendurff MS, et al. Prosthetic intervention effects on activity of lower-extremity amputees. *Arch Phys Med Rehabil.* May 2006; 87(5): 717-22. PMID 16635636

Williams RM, Turner AP, Orendurff M, et al. Does having a computerized prosthetic knee influence cognitive performance during amputee walking?. *Arch Phys Med Rehabil.* Jul 2006; 87(7): 989-94. PMID 16813788

Hafner BJ, Smith DG. Differences in function and safety between Medicare Functional Classification Level-2 and -3 transfemoral amputees and influence of prosthetic knee joint control. *J Rehabil Res Dev.* 2009; 46(3): 417-33. PMID 19675993

Highsmith MJ, Kahle JT, Miro RM, et al. Ramp descent performance with the C-Leg and interrater reliability of the Hill Assessment Index. *Prosthet Orthot Int.* Oct 2013; 37(5): 362-8. PMID 23327837

Howard CL, Wallace C, Perry B, et al. Comparison of mobility and user satisfaction between a microprocessor knee and a standard prosthetic knee: a summary of seven single-subject trials. *Int J Rehabil Res.* Mar 2018; 41(1): 63-73. PMID 29293160

Hafner BJ, Willingham LL, Buell NC, et al. Evaluation of function, performance, and preference as transfemoral amputees transition from mechanical to microprocessor control of the prosthetic knee. *Arch Phys Med Rehabil.* Feb 2007; 88(2): 207-17. PMID 17270519

Kaufman KR, Bernhardt KA, Symms K. Functional assessment and satisfaction of transfemoral amputees with low mobility (FASTK2): A clinical trial of microprocessor-controlled vs. non-microprocessor-controlled knees. *Clin Biomech (Bristol).* Oct 2018; 58: 116-122. PMID 30077128

Kaufman KR, Levine JA, Brey RH, et al. Gait and balance of transfemoral amputees using passive mechanical and microprocessor-controlled prosthetic knees. *Gait Posture.* Oct 2007; 26(4): 489-93. PMID 17869114

Kaufman KR, Levine JA, Brey RH, et al. Energy expenditure and activity of transfemoral amputees using mechanical and microprocessor-controlled prosthetic knees. *Arch Phys Med Rehabil.* Jul 2008; 89(7): 1380-5. PMID 18586142

Johansson JL, Sherrill DM, Riley PO, et al. A clinical comparison of variable-damping and mechanically passive prosthetic knee devices. *Am J Phys Med Rehabil.* Aug 2005; 84(8): 563-75. PMID 16034225

Carse B, Scott H, Brady L, et al. Evaluation of gait outcomes for individuals with established unilateral transfemoral amputation following the provision of microprocessor controlled knees in the context of a clinical service. *Prosthet Orthot Int.* Jun 01 2021; 45(3): 254-261. PMID 34016870

Alzeer AM, Bhaskar Raj N, Shahine EM, et al. Impacts of Microprocessor-Controlled Versus Non-microprocessor-Controlled Prosthetic Knee Joints Among Transfemoral Amputees on Functional Outcomes: A Comparative Study. *Cureus.* Apr 2022; 14(4): e24331. PMID 35607529

Best TK, Seelhoff CA, Wensman J, et al. The clinical effects of the Össur Power Knee with phase-based and default control during sitting, standing, and walking. *J Neuroeng Rehabil.* Sep 29 2025; 22(1): 200. PMID 41024213

Zhou S, Kestur S, Maldonado J, et al. "Comparing the biomechanical response of users of an open-source powered knee and ankle prosthesis versus a passive prosthesis during ramp and stair ambulation". J Biomech. Jun 2025; 186: 112732. PMID 40300430

Hofstad C, Linde H, Limbeek J, et al. Prescription of prosthetic ankle-foot mechanisms after lower limb amputation. Cochrane Database Syst Rev. 2004; 2004(1): CD003978. PMID 14974050

Alimusaj M, Fradet L, Braatz F, et al. Kinematics and kinetics with an adaptive ankle foot system during stair ambulation of transtibial amputees. Gait Posture. Oct 2009; 30(3): 356-63. PMID 19616436

Fradet L, Alimusaj M, Braatz F, et al. Biomechanical analysis of ramp ambulation of transtibial amputees with an adaptive ankle foot system. Gait Posture. Jun 2010; 32(2): 191-8. PMID 20457526

Darter BJ, Wilken JM. Energetic consequences of using a prosthesis with adaptive ankle motion during slope walking in persons with a transtibial amputation. Prosthet Orthot Int. Feb 2014; 38(1): 5-11. PMID 23525888

Gailey RS, Gaunaud I, Agrawal V, et al. Application of self-report and performance-based outcome measures to determine functional differences between four categories of prosthetic feet. J Rehabil Res Dev. 2012; 49(4): 597-612. PMID 22773262

Delussu AS, Brunelli S, Paradisi F, et al. Assessment of the effects of carbon fiber and bionic foot during overground and treadmill walking in transtibial amputees. Gait Posture. Sep 2013; 38(4): 876-82. PMID 23702342

Thomas-Pohl M, Villa C, Davot J, et al. Microprocessor prosthetic ankles: comparative biomechanical evaluation of people with transtibial traumatic amputation during standing on level ground and slope. Disabil Rehabil Assist Technol. Jan 2021; 16(1): 17-26. PMID 31535903

Colas-Ribas C, Martinet N, Audat G, et al. Effects of a microprocessor-controlled ankle-foot unit on energy expenditure, quality of life, and postural stability in persons with transtibial amputation: An unblinded, randomized, controlled, cross-over study. Prosthet Orthot Int. Dec 01 2022; 46(6): 541-548. PMID 36515900

Au S, Berniker M, Herr H. Powered ankle-foot prosthesis to assist level-ground and stair-descent gaits. Neural Netw. May 2008; 21(4): 654-66. PMID 18499394

Ferris AE, Aldridge JM, Rábago CA, et al. Evaluation of a powered ankle-foot prosthetic system during walking. Arch Phys Med Rehabil. Nov 2012; 93(11): 1911-8. PMID 22732369

Herr HM, Grabowski AM. Bionic ankle-foot prosthesis normalizes walking gait for persons with leg amputation. Proc Biol Sci. Feb 07 2012; 279(1728): 457-64. PMID 21752817

Mancinelli C, Patrilli BL, Tropea P, et al. Comparing a passive-elastic and a powered prosthesis in transtibial amputees. Annu Int Conf IEEE Eng Med Biol Soc. 2011; 2011: 8255-8. PMID 22256259

Cacciola CE, Kannenberg A, Hibler KD, Howell J. Impact of a Powered Prosthetic Ankle-Foot Component on Musculoskeletal Pain in Individuals with Transtibial Amputation: A Real-World Cross-Sectional Study with Concurrent and Recalled Pain and Functional Ratings. J Prosthet Orthot. 2022. doi: 10.1097/JPO.0000000000000442.

VA/DOD Clinical Practice Guideline, Work Group. VA/DOD Clinical Practice Guideline for Rehabilitation of Individuals With Lower Limb Amputation. U.S. Government Printing Office; 2024:2-162.  
[https://www.healthquality.va.gov/guidelines/Rehab/amp/LLA-CPG\\_2024-Guideline\\_final\\_20250110.pdf](https://www.healthquality.va.gov/guidelines/Rehab/amp/LLA-CPG_2024-Guideline_final_20250110.pdf)

**CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS**

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

