

EFFECTIVE DATE: 10|01|2026

POLICY LAST REVIEWED: 06|03|2026

OVERVIEW

Microvolt T-wave alternans (MTWA) testing has been investigated as a noninvasive test to identify a patient's risk for sudden cardiac death. The test measures the beat-to-beat microvolt variation in the amplitude of the electrocardiogram tracing. Some research indicates a positive test has a greater risk of developing ventricular tachyarrhythmias than a negative test.

MEDICAL CRITERIA

Medicare Advantage Plans

Not applicable

Commercial Products

MTWA testing using the spectral analytic method is considered medically necessary for the evaluation of persons at risk of sudden cardiac death for medically necessary implantable cardioverter-defibrillator placement.

PRIOR AUTHORIZATION

Medicare Advantage Plans

Not applicable

Commercial Products

Prior authorization is recommended for Commercial Products.

POLICY STATEMENT

Medicare Advantage Plans

MTWA testing is covered.

Note: Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all Medicare Advantage Plans policies. Therefore, Medicare Advantage Plans policies may differ from Commercial Products. In some instances, benefits for Medicare Advantage Plans may be greater than what is allowed by the CMS.

Commercial Products

MTWA testing using the spectral analytic method is considered medically necessary when the medical criteria above is met.

MTWA testing using the spectral analytic method is considered not medically necessary when the medical criteria above is not met.

MTWA testing using the spectral analytic method is considered not medically necessary for the following indications as the evidence is insufficient to determine the effects of the technology on health outcomes:

1. Diagnosis and risk assessment of acute coronary syndrome
2. Diagnosis of reversible myocardial ischemia in individuals without structural heart disease
3. Evaluation of non-pathological preterm infants
4. Evaluation of children and adolescents with Eisenmenger syndrome

5. Evaluation of malignant ventricular arrhythmias in Chagas disease
6. Evaluation of the adequacy of medical therapy
7. Guidance of anti-arrhythmic therapy
8. Judgement of the severity of ischemic cardiomyopathy
9. Prediction of major adverse cardiac events in ischemic heart failure
10. Prediction of post-operative mortality in cardiac surgery
11. Prognosis of myocardial function in newborns with hypoxic-ischemic encephalopathy
12. Prognosis of pulmonary arterial hypertension
13. Prognostic evaluation in hypertrophic cardiomyopathy
14. Risk assessment of sudden cardiac death in children with chronic renal failure
15. Risk stratification of cardiac events (e.g., sudden cardiac death) in members following repair of tetralogy of Fallot
16. Risk stratification in Brugada syndrome
17. Tracking changes in risk during cardiac disease progression

MTWA combined with electrophysiologic study for prediction of ventricular tachyarrhythmias in individuals with arrhythmogenic right ventricular cardiomyopathy is considered not medically necessary as the evidence is insufficient to determine the effects of the technology on health outcomes.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable machine tests or limitations of benefits/coverage when services are not medically necessary.

BACKGROUND

Microvolt T-wave alternans (MTWA) refers to a beat-to-beat variability in T-wave amplitude. Because a routine electrocardiogram (EKG) cannot detect these small fluctuations, this test requires specialized sensors to detect the fluctuations and computer algorithms to evaluate the results. T-wave alternans is measured by a provocative test that requires gradual elevation of the heart rate to more than 110 beats per minute. The test can be performed in conjunction with an exercise tolerance stress test. Test results are reported as the number of standard deviations (SDs) by which the peak signal of the T-wave exceeds the background noise. This number is referred to as the alternans ratio. An alternans ratio of 3 or greater is typically considered a positive result, an absent alternans ratio is considered a negative result, and other values are indeterminate.

The presence of T-wave alternans has been investigated as a risk factor for fatal arrhythmias and sudden cardiac death in patients with a history of myocardial infarction (MI), heart failure, or cardiomyopathy. Patients with these disorders at high-risk for sudden cardiac death may be treated with medications to suppress the emergence of arrhythmias or undergo implantation of cardiac defibrillators to terminate tachyarrhythmias when they occur. Since sudden cardiac death is one of the most common causes of death after a MI or in patients with dilated cardiomyopathy, there is substantial interest in risk stratification to target therapy.

Patient groups are categorized into those who have not experienced a life-threatening arrhythmia (i.e., primary prevention) and those who have (i.e., secondary prevention). Those who have experienced a life-threatening arrhythmia are already at high risk and would not be considered for testing. T-wave alternans is one of many risk factors that have been investigated for identifying candidates for primary prevention. Others include left ventricular ejection fraction (LVEF), arrhythmias detected on Holter monitor or electrophysiologic studies, heart rate variability, and baroreceptor sensitivity. Signal-averaged electrocardiography (SAECG) is another technique for risk stratification. It measures beat-averaged conduction, while T-wave alternans measures beat-to-beat variability.

MTWA testing has primarily been used for defining the risk of ventricular arrhythmias in persons at risk for sudden cardiac death and determining which patients are most likely to benefit from implantable cardioverter-defibrillators. The quality of evidence is adequate to conclude that MTWA testing using a spectral analysis

algorithm can improve net health outcomes for patients who are candidates for implantable cardioverter defibrillator (ICD) placement.

Commercial Products

There are many established tools for diagnosis, prognosis, and clinical decision making for acute coronary syndrome. Understanding the advantages and limitations of each tool according to the clinical scenario is essential. Several emerging tools, such as novel biomarkers (e.g., high-sensitivity troponin and growth differentiation factor-15), electroencephalographic (ECG) techniques (e.g., heart rate turbulence or MTWA), and imaging modalities (computed tomography angiography and cardiac magnetic resonance) may potentially improve clinical care; however, they must be fully evaluated and validated in different scenarios and patient cohorts before they are incorporated into clinical practice. The evidence is insufficient to determine the effects of the technology on health outcomes.

Findings in one study showed that MTWA could be the new non-invasive tool for the detection of reversible ischemia in patients with suspected coronary artery disease without structural heart disease. Furthermore, MTWA can detect ischemia earlier and with greater accuracy compared with exercise ECG testing. However, this study had several drawbacks. The study was conducted on a relatively small number of patients. The research could be seen as a pilot study and encouragement for large multi-center study with a view to a definitive confirmation of criteria and values of MTWA for the diagnosis of reversible ischemia in patients without structural heart disease. The evidence is insufficient to determine the effects of the technology on health outcomes.

A preliminary retrospective study suggested that non-pathological preterm infants showed MTWA could potentially indicate a condition of cardiac risk possibly related to the low development status of the infant but further investigations are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Studies with longer follow-up periods investigating the risks of development of sudden death and arrhythmia or cardiovascular mortality in pediatric individuals with Eisenmenger syndrome are needed to obtain definite results. The evidence is insufficient to determine the effects of the technology on health outcomes.

MTWA is a direct measure of ventricular repolarization instability and has emerged as a potentially useful way of determining arrhythmia vulnerability in members with Chagas disease. However, this methodology has not been evaluated in patients with Chagas disease. Further assessment in a prospective study is needed to establish the causality and clinical application of the test. The evidence is insufficient to determine the effects of the technology on health outcomes.

Studies have resulted in questions currently undergoing investigation, to include MTWA's potential to guide ICD implantation, track changes in risk during cardiac disease progression, and evaluate the adequacy of medical therapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

MTWA represents a potential useful tool judge severity following ischemic cardiomyopathy. However, the evidence is insufficient to determine the effects of the technology on health outcomes.

The prognostic value of MTWA for the prediction of major adverse cardiac events in ischemic heart failure has not been identified to-date. The evidence is insufficient to determine the effects of the technology on health outcomes.

MTWA at the end of cardiac surgery could potentially predict in-hospital mortality. This predictability was more robust when combined with the EuroSCORE II. Additional studies on the robustness of MTWA as predictive marker in a larger cohort are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Global cardiac functions and myocardial performance of newborns with hypoxic-ischemic encephalopathy might be improved with therapeutic hypothermia which can be determined by using MTWA and tissue-

Doppler measurements. However, further studies are needed to evaluate whether MTWA is prognostic in determining the myocardial dysfunction and arrhythmias. The evidence is insufficient to determine the effects of the technology on health outcomes.

The role of MTWA in pulmonary arterial hypertension remains obscure. Further research is needed to elucidate the clinical significance and prognostic value of MTWA. The evidence is insufficient to determine the effects of the technology on health outcomes.

For the risk assessment of sudden cardiac death in children with chronic renal failure, studies for MTWA are lacking but might be used for early risk assessment in pediatric patients with CRF in the future. The evidence is insufficient to determine the effects of the technology on health outcomes.

There is a significant risk of arrhythmia and sudden death after repair of congenital heart disease. Studies to date indicate further studies of the role of MTWA following repair of tetralogy of Fallot are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For risk stratification in Brugada syndrome, studies were found to be hypothetical (i.e., not based on experimental data); thus, further rigorous studies are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

One study showed that the predicted value of MTWA for ventricular tachyarrhythmia in patients with arrhythmogenic right ventricular cardiomyopathy remains unclear. Despite a follow-up period as long as 8 years, this study was limited by a small sample size affiliated with a single center. The evidence is insufficient to determine the effects of the technology on health outcomes.

MTWA is associated with the occurrence of ventricular arrhythmias in several heart diseases; however, its role in HCM remains uncertain. One study showed that altered MTWA was not associated with the occurrence of SCD or potentially fatal ventricular arrhythmias in HCM patients, and the low rate of these events during long-term follow-up suggested the good prognosis of this heart disease. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

Medicare Advantage Plans

The following code is covered:

93025 Microvolt T-wave alternans for assessment of ventricular arrhythmias

Commercial Products

The following code is covered when the medical criteria above is met:

93025 Microvolt T-wave alternans for assessment of ventricular arrhythmias

RELATED POLICIES

Prior Authorization of Services, Treatments or Procedures

PUBLISHED

Provider Update, August 2026

Provider Update, June 2025

Provider Update, April 2024

Provider Update, May 2023

Provider Update June 2022

REFERENCES

1. Centers for Medicare and Medicaid Services (CMS) National Coverage Determination (NCD), 20.30, Microvolt T-Wave Alternans (MTWA)
2. Adachi K, Ohnishi Y, Shima T, et al. Determinant of microvolt-level T-wave alternans testing specifically in patients with dilated cardiomyopathy. *J Am Coll Cardiol.* 1999;34:374-380.

3. Adachi K, Ohnishi Y, Yokoyama M. Risk stratification for sudden cardiac death in dilated cardiomyopathy using microvolt-level T-wave alternans. *Jpn Circ J*. 2001;65(2):76-80.
4. Adler A, Rosso R, Chorin E, et al. Risk stratification in Brugada syndrome: Clinical characteristics, electrocardiographic parameters, and auxiliary testing. *Heart Rhythm*. 2016;13(1):299-310.
5. Almeida BCS, Carmo AALD, Barbosa MPT, et al. Association between microvolt T-wave alternans and malignant ventricular arrhythmias in Chagas disease. *Arq Bras Cardiol*. 2018;110(5):412-417. 5.
6. Antman EM, Anbe DT, Armstrong PW, et al.; American College of Cardiology; American Heart Association Task Force on Practice Guidelines. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction--executive summary: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction). *Circulation*. 2004;110(5):588-636.
7. Antunes MO, Arteaga-Fernandez E, Samesima N, et al. Prognostic evaluation of microvolt T-wave alternans in hypertrophic cardiomyopathy: 9-year clinical follow-up. *Bras Cardiol*. 2023;120(8):e20220833.
8. Barbosa MP, da Costa Rocha MO, Neto ES, et al. Usefulness of microvolt T-wave alternans for predicting outcome inpatients with Chagas disease with implantable cardioverter defibrillators. *Int J Cardiol*. 2016;222:80-85.
9. Bartczak A, Trojnarzka O, Cieplucha A, et al. Microvolt T-wave alternans in adult patients with repaired tetralogy of Fallot. *2015;10(2):E89-E97*.
10. BlueCross BlueShield Association (BCBSA), Technology Evaluation Center (TEC). Microvolt T-wave alternans testing to risk stratify patients being considered for ICD therapy for primary prevention of sudden death. TEC Assessment Program. Chicago, IL: BCBSA; October 2005;20(9).
11. BlueCross BlueShield Association (BCBSA), Technology Evaluation Center (TEC). Microvolt T-wave alternans testing to risk-stratify patients being considered for ICD therapy for primary prevention of sudden death. TEC Assessment Program. Chicago, IL: BCBSA; June 2007;21(14). 10.
12. Cain ME, Arthur RM, Trobaugh JW. Detection of the fingerprint of the electrophysiological abnormalities that increase vulnerability to life-threatening ventricular arrhythmias. *J Interv Card Electrophysiol*. 2003;9(2):103-118.
13. California Technology Assessment Forum (CTAF). Microvolt T-wave alternans testing to risk-stratify patients for implantable cardioverter-defibrillator placement for prevention of sudden cardiac death. A Technology Assessment. San Francisco, CA:CTAF; October 18, 2006.
14. Centers for Medicare and Medicaid Services (CMS). Decision memo for microvolt T-wave alternans (CAG-00293R). Medicare Coverage Database. Baltimore, MD: CMS; May 12, 2008.
15. Centers for Medicare and Medicaid Services (CMS). Decision memo for implantable defibrillators (CAG-00157R3). Baltimore, MD: CMS; January 27, 2005.
16. Centers for Medicare and Medicaid Services (CMS). Decision memo for microvolt T-wave alternans (CAG-00293N). Medicare Coverage Database. Baltimore, MD: CMS; March 21, 2006.
17. Chauhan VS, Selvaraj RJ. Utility of microvolt T-wave alternans to predict sudden cardiac death in patients with cardiomyopathy. *Curr Opin Cardiol*. 2007;22(1):25-32.
18. Chen Z, Shi Y, Hou X, et al. Microvolt T-wave alternans for risk stratification of cardiac events in ischemic cardiomyopathy: A meta-analysis. *Int J Cardiol*. 2013;167(5):2061-2065. 17.
19. Cheung MM, Weintraub RG, Cohen RJ, et al. T wave alternans threshold late after repair of tetralogy of Fallot. *JCardiovasc Electrophysiol*. 2002;13(7):657-661.
20. Chow T, Kereiakes DJ, Onufer J, et al; MASTER Trial Investigators. Does microvolt T-wave alternans testing predict ventricular tachyarrhythmias in patients with ischemic cardiomyopathy and prophylactic defibrillators? The MASTER(Microvolt T Wave Alternans Testing for Risk Stratification of Post-Myocardial Infarction Patients) trial. *J Am Coll Cardiol*. 2008;52(20):1607-1615.
21. Cleland JG, Coletta AP, Nikitin N, et al. Update of clinical trials from the American College of Cardiology 2003. EPHEBUS, SPORTIF-III, ASCOT, COMPANION, UK-PACE and T-wave alternans. *Eur J Heart Fail*. 2003;5(3):391-398.
22. Costantini O, Hohnloser SH, Kirk MM, et al; ABCD Trial Investigators. The ABCD (Alternans Before Cardioverter Defibrillator) Trial: Strategies using T-wave alternans to improve efficiency of sudden cardiac death prevention. *J Am Coll Cardiol*. 2009;53(6):471-479.

23. Cutler MJ, Rosenbaum DS. Risk stratification for sudden cardiac death: Is there a clinical role for T wave alternans? *HeartRhythm*. 2009;6(8 Suppl):S56-S61.
24. Danilowicz-Szymanowicz L, Lewicka E, Dabrowska-Kugacka A, et al. Microvolt T-wave alternans profiles in patients with pulmonary arterial hypertension compared to patients with left ventricular systolic dysfunction and a group of healthy volunteers. *Anatol J Cardiol*. 2016;16(11):825-830.
25. Doyle T, Kavanaugh-McHugh A. Pathophysiology, clinical features, and diagnosis of tetralogy of Fallot. UpToDate [onlineserial]. Waltham, MA: UpToDate; reviewed May 2015.
26. Engel G, Beckerman JG, Froelicher VF, et al. Electrocardiographic arrhythmia risk testing. *Curr Probl Cardiol*. 2004;29(7):365-432.
27. Francis DP, Salukhe TV. Who needs a defibrillator after myocardial infarction? *Lancet*. 2003;362(9378):91-92.
28. Gehi AK, Stein RH, Metz LD, Gomes JA. Microvolt T-wave alternans for the risk stratification of ventricular tachyarrhythmic events: A meta-analysis. *J Am Coll Cardiol*. 2005;46(1):75-82.
29. Gold MR, Bloomfield DM, Anderson KP, et al. A comparison of T-wave alternans, signal averaged electrocardiography and programmed ventricular stimulation for arrhythmia risk stratification. *J Am Coll Cardiol* 2000;36:2247-2253.
30. Gold MR, Ip JH, Costantini O, et al. Role of microvolt T-wave alternans in assessment of arrhythmia vulnerability among patients with heart failure and systolic dysfunction: Primary results from the T-wave alternans sudden cardiac death in heart failure trial sub study. *Circulation*. 2008;118(20):2022-2028.
31. Goldberger JJ, Subacius H, Patel T, et al. Sudden cardiac death risk stratification in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol*. 2014;63(18):1879-1889.
32. Grimm W, Hoffmann J, Menz V, Maisch B. Relation between microvolt level T wave alternans and other potential noninvasive predictors of arrhythmic risk in the Marburg Cardiomyopathy Study. *Pacing Clin Electrophysiol*. 2000;23(11 Pt2):1960-1964.
33. Gupta A, Hoang DD, Karliner L, et al. Ability of microvolt T-wave alternans to modify risk assessment of ventricular tachyarrhythmic events: A meta-analysis. *Am Heart J*. 2012;163(3):354-364.
34. Haghjoo M, Arya A, Sadr-Ameli MA. Value of microvolt T-wave alternans for predicting patients who would benefit from implantable cardioverter-defibrillator therapy. *Cardiol Rev*. 2006;14(4):173-179.
35. Halliöglu O, Keceli M, Bozlu G, et al. Evaluation of T-wave alternans in pediatric patients with chronic renal failure. *J Electrocardiol*. 2018;51(4):622-627.
36. Hennersdorf MG, Niebch V, Perings C, Strauer BE. T wave alternans and ventricular arrhythmias in arterial hypertension. *Hypertension*. 2001;37(2):199-203.
37. Hennersdorf MG, Perings C, Niebch V, et al. T wave alternans as a risk predictor in patients with cardiomyopathy and mild-to-moderate heart failure. *Pacing Clin Electrophysiol*. 2000;23(9):1386-1391.
38. Hohnloser SH, Ikeda T, Bloomfield DM, et al. T-wave alternans negative coronary patients with low ejection and benefit from defibrillator implantation. *Lancet*. 2003a;362(9378):125-126.
39. Hohnloser SH, Klingenheben T, Bloomfield D, et al. Usefulness of microvolt T-wave alternans for prediction of ventricular tachyarrhythmic events in patients with dilated cardiomyopathy: Results from a prospective observational study. *J Am Coll Cardiol*. 2003b;41(12):2220-2224.
40. Hombach V. Electrocardiogram of the failing heart. *Card Electrophysiol Rev*. 2002;6(3):209-214.
41. Ikeda T, Saito H, Tanno K, et al. T-wave alternans as a predictor for sudden cardiac death after myocardial infarction. *Am J Cardiol*. 2002;89(1):79-82.
42. Ikeda T, Sakata T, Takami M, et al. Combined assessment of T-wave alternans and late potentials used to predict arrhythmic events after myocardial infarction. *J Am Coll Cardiol*. 2000;35:722-730.
43. Ikeda T, Sakurada H, Sakabe K, et al. Assessment of noninvasive markers in identifying patients at risk in the Brugada syndrome: Insight into risk stratification. *J Am Coll Cardiol*. 2001;37(6):1628-1634.
44. Kadish, AH, Buxton AE, Kennedy HL, et al. ACC/AHA clinical competence statement on electrocardiography and ambulatory electrocardiography. *J Am Coll Cardiol*. 2001;38(7):2091-2100.
45. Karpuz D, Celik Y, Giray D, et al. Therapeutic hypothermia and myocardium in perinatal asphyxia: A microvolt T-wave alternans and Doppler echocardiography study. *Bratisl Lek Listy*. 2017;118(12):765-771.
46. Karpuz D, Halliöglu O, Yilmaz DC. Increased microvolt T-wave alternans in children and adolescents with Eisenmenger syndrome. *Anatol J Cardiol*. 2018;19(5):303-310.

47. Kaufmann DK, Raczak G, Szwoch M, et al. Baroreflex sensitivity but not microvolt T-wave alternans can predict major adverse cardiac events in ischemic heart failure. *Cardiol J*. 2022;29(6):1004-1012.
48. Kitamura H, Ohnishi Y, Okajima K, et al. Onset heart rate of microvolt-level T-wave alternans provides clinical and prognostic value in nonischemic dilated cardiomyopathy. *J Am Coll Cardiol*. 2002;39(2):295-300.
49. Klingenheben T, Hohnloser SH. Clinical value of T-wave alternans assessment. *Card Electrophysiol Rev*. 2002;6(3):323-328.
50. Klingenheben T, Ptaszynski P. Clinical significance of microvolt T-wave alternans. *Herzschrittmacherther Elektrophysiol*. 2007;18(1):39-44.
51. Klingenheben T, Zabel M, D'Agostino RB, et al. Predictive value of T-wave alternans for arrhythmic events in patients with congestive heart failure. *Lancet*. 2000;356(9230):651-652.
52. Koo CH, Lee HC, Kim TK, et al. Microvolt T-wave alternans at the end of surgery is associated with postoperative mortality in cardiac surgery patients. *Sci Rep*. 2019;9(1):17351.
53. Kulkarni K, Walton RD, Armoundas AA, Tolkacheva EG. Clinical potential of beat-to-beat diastolic interval control in preventing cardiac arrhythmias. *J Am Heart Assoc*. 2021;10(11):e020750.
54. Kusmirek SL, Gold MR. Sudden cardiac death: The role of risk stratification. *Am Heart J*. 2007;153(4 Suppl):25-33.
55. Marcantoni I, Sbröllini A, Agostinelli G, et al. T-wave alternans in nonpathological preterm infants. *Ann Noninvasive Electrocardiol*. 2020;25(4):e12745.
56. Narayan SM. T wave (repolarization) alternans: Clinical aspects. UpToDate [online serial]. Waltham, MA:UpToDate; reviewed May 2015.
57. National Institutes of Health (NIH), National Heart, Lung, and Blood Institute (NHLBI). Prognostic significance of T wave alternans. ClinicalTrials.gov Abstract. Study ID Numbers 942. NLM Identifier NCT00006501. Bethesda, MD: NIH; January 29, 2001. .
58. Pedretti RF, Sarzi Braga S. Non-invasive sudden death risk stratification. *Ital Heart J*. 2005;6(3):180-189.
59. Pelli A, Kentta TV, Junttila MJ, et al; EU-CERT-ICD Study Investigators. Lack of prognostic value of T-wave alternans for implantable cardioverter-defibrillator benefit in primary prevention. *J Am Heart Assoc*. 2024;13(11):e032465.
60. Puljevic M, Danilowicz-Szymanowicz L, Molon G, et al. TWARDMI pilot trial: The value and optimal criteria of microvolt T-wave alternans in the diagnosis of reversible myocardial ischemia in patients without structural cardiac disease. *Ann Noninvasive Electrocardiol* 2019;24(2): e12610.
61. Quan XQ, Zhou HL, Ruan L, et al. Ability of ambulatory ECG-based T-wave alternans to modify risk assessment of cardiac events: A systematic review. *BMC Cardiovasc Disord*. 2014;14:198.
62. Sakamoto S, Takagi M, Kakihara J, et al. The utility of T-wave alternans during the morning in the summer for the risk stratification of patients with Brugada syndrome. *Heart Vessels*. 2017;32(3):341-351.
63. Scirica BM. Acute coronary syndrome: Emerging tools for diagnosis and risk assessment. *J Am Coll Cardiol*. 2010;55(14):1403-1415.
64. Sturdivant JL, Gold MR. T wave alternans for ventricular arrhythmia risk stratification. *Minerva Cardioangiol*. 2003;51(1):15-19.
65. Sullivan T, Hiller J. Microvolt T-wave alternans; Horizon scanning prioritizing summary - volume 15. Adelaide, SA: AdelaideHealth Technology Assessment (AHTA) on behalf of National Horizon Scanning Unit (HealthPACT and MSAC); 2007
66. U.S. Food and Drug Administration (FDA). 510(k) Final Decisions Rendered for April 1999. Cambridge Heart, Inc. Cardiac Diagnostic System Model CH 2000. 510(k) No. K983102. Rockville, MD: FDA; April 12, 1999.
67. van der Avoort CJ, Filion KB, Dendukuri N, Brophy JM. Microvolt T-wave alternans as a predictor of mortality and severe arrhythmias in patients with left-ventricular dysfunction: A systematic review and meta-analysis. *BMC Cardiovascular Disorders*. 2009;9(5):1-9.
68. Verrier RL, Klingenheben T, Malik M, et al. Microvolt T-wave alternans physiological basis, methods of measurement, and clinical utility -- consensus guideline by International Society for Holter and Noninvasive Electrocardiology. *J Am Coll Cardiol*. 2011;58(13):1309-1324.
69. Verrier RL, Nieminen T. T-wave alternans as a therapeutic marker for antiarrhythmic agents. *J Cardiovasc Pharmacol*. 2010;55(6):544-554.

70. Verrier RL, Sroubek J. Quantitative T-wave alternans analysis for sudden cardiac death risk assessment and guiding therapy: Answered and unanswered questions: For: Proceedings of ICE2015 Comandatuba, Brazil, Sudden Death Symposium. *J Electrocardiol.* 2016;49(3):429-438.
71. Verrier RL, Tolat AV, Josephson ME. T-Wave alternans for arrhythmia risk stratification in patients with idiopathic dilated cardiomyopathy. *J Am Coll Cardiol.* 2003;41(12):2225-2227.
72. Vitali F, Brieda A, Balla C, et al. Standard ECG in Brugada syndrome as a marker of prognosis: From risk stratification to pathophysiological insights. *J Am Heart Assoc.* 2021;10(10):e020767.
73. Walker ML, Rosenbaum DS. Repolarization alternans: Implications for the mechanism and prevention of sudden cardiac death. *Cardiovasc Res.* 2003;57(3):599-614.
74. Windhagen-Mahnert B, Kadish AH. Ventricular arrhythmias. Application of noninvasive and invasive tests for risk assessment in patients with ventricular arrhythmias. *Cardiol Clin.* 2000;18(2):243-263, vii.
75. Wylie JV, Garlitski AC. Brugada syndrome: Clinical presentation, diagnosis, and evaluation. UpToDate [online serial]., Waltham, MA: UpToDate; reviewed March 2017.
76. Xue S-L, Hou X-F, Sun K-Y, et al. Microvolt T-wave alternans complemented with electrophysiologic study for prediction of ventricular tachyarrhythmias in patients with arrhythmogenic right ventricular cardiomyopathy: A long-term follow-up study. *Chin Med J (Engl)* 2019;132(12):1406-1413.
77. Yang J, Luo J, Li K, et al. T-wave alternans measured by 24-hour ambulatory recordings rather than exercise stress tests as a risk stratification marker in patients with long QT syndrome. *J Am Heart Assoc.* 2024;13(14):e033619.

DRAFT

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

