DRAFT Medical Coverage Policy | Minimally Invasive Procedures for Back Pain

EFFECTIVE DATE: 12|01|2026
POLICY LAST REVIEWED: 08|19|2026

OVERVIEW

This policy addresses a variety of minimally invasive techniques that have been investigated over the years as treatment of low back pain related to disc disease. Surgical management of herniated intervertebral discs most commonly involves discectomy or microdiscectomy, performed manually through an open incision. Automated percutaneous discectomy involves placement of a probe within the intervertebral disc under image guidance with aspiration of disc material using a suction cutting device. Electrothermal intradiscal annuloplasty therapies use radiofrequency energy sources to treat discogenic low back pain arising from annular tears. These annuloplasty techniques are designed to decrease pain arising from the annulus by thermocoagulating nerves in the disc and tightening annular tissue. Laser energy (laser discectomy) and radiofrequency coblation (nucleoplasty) are being evaluated for decompression of the intervertebral disc. For laser discectomy under fluoroscopic guidance, a needle or catheter is inserted into the disc nucleus, and a laser beam is directed through it to vaporize tissue. For disc nucleoplasty, bipolar radiofrequency energy is directed into the disc to ablate tissue.

MEDICAL CRITERIA

Medicare Advantage Plans

Effective 10/1/2026, the following CPTs 64628/64629 may be considered medically necessary for Medicare Advantage Plans when the medical criteria in the online authorization tool for participating providers is met.

Commercial Products

Thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system) will be considered medically reasonable and necessary for the treatment of chronic low back pain in patients who meet ALL the following criteria and do not meet any of the contraindications (#6) below:

1. Chronic lumbar back pain of ≥6 months duration that causes functional deficit measured on a pain or disability scale*, AND
2. Documented failure to respond to ≥6 months of non-surgical management**, AND
3. Absence of non-vertebrogenic pathology per clinical assessment or radiology studies that could explain the source of the patient's pain, including but not limited to fracture, tumor, infection, or significant deformity, AND
4. Evidence of Type 1 or Type 2 Modic changes on MRI, such as inflammation, edema, vertebral endplate changes, disruption and fissuring of the endplate, vascularized fibrous tissues within the adjacent marrow, hypotensive signals (Type 1 Modic change), and changes to the vertebral body marrow including replacement of normal bone marrow by fat, and hypertensive signals (Type 2 Modic change), in 1 or more vertebrae from L3-S1, AND
5. Individuals must have undergone careful screening, evaluation, and diagnosis by a multidisciplinary team prior to thermal destruction of the intraosseous BVN (such screening must include psychological, as well as, physical evaluation). Documentation of the history and careful screening must be available in the patient chart if requested, AND
6. None of the following contraindications listed below are present at the time thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system) is performed:
    1. Skeletally immature patients (≤18 years old);
    2. Severe cardiac or pulmonary compromise;
    3. Active systemic infection or local infection at the intended treatment level;
    4. Bleeding diathesis;
    5. Pregnancy;
    6. Primary radicular pain into the lower extremities (defined as nerve pain following a dermatomal

distribution and that correlates with nerve compression on imaging);
7. Previous lumbar/lumbosacral spine surgery at the intended treatment level (with the exception of discectomy/laminectomy if performed >6 months prior to BVN nerve ablation and radicular pain resolved);
8. Primary symptomatic lumbar or lumbosacral spinal stenosis (defined as the presence of neurogenic claudication and confirmed by imaging);
9. Diagnosed osteoporosis (T-score of -2.5 or less), spine fragility fracture history, trauma/compression fracture at the intended treatment level, or spinal cancer;
10. Radiographic evidence of any of the following that correlates with predominant physical complaints:
    a. Lumbar/lumbosacral disc extrusion or protrusion >5mm at levels L3-S1;
    b. Lumbar/lumbosacral spondylolisthesis > 2mm at any level;
    c. Lumbar/lumbosacral spondylolysis at levels L3-S1;
    d. Lumbar/lumbosacral facet arthrosis/effusion correlated with facet-mediated pain at levels L3-S1.
11. BMI >40;
12. Advanced generalized systemic disease that limits quality-of-life (QOL) improvements would require a statement of the objective of treatment in such cases;
13. Active, untreated substance abuse disorder;
14. Individual is a tobacco user OR there is no clinical documentation that the individual has been abstinent from tobacco use based on attestation.

NOTE: Thermal destruction of the intraosseous BVN must only be performed once per vertebral body from L3-S1 per lifetime. Up to 4 vertebral bodies may be treated during 1 procedure.

* Pain assessment and a disability scale must be obtained at baseline to be used for functional assessment.

** Non-surgical management may include but is not limited to:

- Avoidance of activities that aggravate pain;
- Trial of Chiropractic manipulation
- Trial of Physical Therapy;
- Cognitive support and recovery reassurance;
- Injection therapy – epidural and/or facet;
- Spine biomechanics education;
- Specific lumbar exercise program;
- Home use of heat/cold modalities;
- Low impact aerobic exercise as tolerated;
- Pharmacotherapy (e.g., non-narcotic analgesics, NSAIDs, muscle relaxants, neuroleptics, and narcotics).

PRIOR AUTHORIZATION

Medicare Advantage Plans

Prior authorization is required for Medicare Advantage Plans via the online tool for participating providers for thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system).

Commercial Products

Prior authorization is recommended for Commercial Products for thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system).

POLICY STATEMENT

Medicare Advantage Plans

Effective 10/1/2026, CPTs 64628/64629 when used for thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system) may be considered medically necessary for Medicare Advantage Plans when the medical criteria in the online authorization tool for participating providers is met.

DRAFT

Commercial Products
Thermal destruction of the intraosseous basivertebral nerve (e.g., Intracept® system) will be considered medically necessary when the criteria above have been met.

Medicare Advantage Plans and Commercial Products
The following services are not covered for Medicare Advantage Plans and not medically necessary for Commercial Products as the evidence is insufficient to determine that the technology results in an improvement in the net health outcomes:

- Percutaneous annuloplasty (e.g., intradiscal electrothermal annuloplasty, intradiscal radiofrequency annuloplasty, or intradiscal biacuplasty) for the treatment of chronic discogenic back pain
- Laser discectomy and radiofrequency coblation (disc nucleoplasty) as techniques of disc decompression and treatment of associated pain.
- Automated percutaneous discectomy as a technique of intervertebral disc decompression in individuals with back pain and/or radiculopathy related to disc herniation in the lumbar, thoracic, or cervical spine.

COVERAGE
Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND
Medicare Advantage Plans
Percutaneous thermal intradiscal procedures (TIPs) involve the insertion of a catheter(s)/probe(s) in the spinal disc under fluoroscopic guidance for the purpose of producing or applying heat and/or disruption within the disc to relieve low back pain.

The scope of the Centers for Medicare and Medicaid Services national coverage determination on TIPs includes percutaneous intradiscal techniques that employ the use of a radiofrequency energy source or electrothermal energy to apply or create heat and/or disruption within the disc for coagulation and/or decompression of disc material to treat symptomatic patients with annular disruption of a contained herniated disc, to seal annular tears or fissures, or destroy nociceptors for the purpose of relieving pain. This includes techniques that use single or multiple probe(s)/catheter(s), which utilize a resistance coil or other delivery system technology, are flexible or rigid, and are placed within the nucleus, the nuclear-annular junction, or the annulus.

Although not intended to be an all-inclusive list, TIPs are commonly identified as intradiscal electrothermal therapy (IDET), intradiscal thermal annuloplasty (IDTA), percutaneous intradiscal radiofrequency thermocoagulation (PIRFT), radiofrequency annuloplasty (RA), intradiscal biacuplasty (IDB), percutaneous (or plasma) disc decompression (PDD) or coblation, or targeted disc decompression (TDD). At times, TIPs are identified or labeled based on the name of the catheter/probe that is used (e.g., SpineCath, discTRODE, SpineWand, Accutherm, or TransDiscal electrodes). Each technique or device has its own protocol for application of the therapy.

The Centers for Medicare and Medicaid Services has determined that TIPs are not reasonable and necessary for the treatment of low back pain. Therefore, TIPs, which include procedures that employ the use of a radiofrequency energy source or electrothermal energy to apply or create heat and/or disruption within the disc for the treatment of low back pain, are noncovered. Therefore, these services are not covered for Medicare Advantage Plans.

Medicare Advantage Plans and Commercial Products
Low back pain (LBP) is the most expensive occupational disorder in the United States and the leading cause of disability worldwide. Chronic low back pain (cLBP) is defined as persistent pain in the lumbar region lasting for >12 weeks. cLBP has many different etiologies. Research shows evidence that one etiology is associated with degeneration of the vertebral body or vertebral body endplates, resulting in inflammation. The

inflammatory response is perceived by the basivertebral nerve (BVN), a sensory nerve that enters the posterior vertebral body and branches out to the superior and inferior endplates. The pain signals are then transmitted to the central nervous system, causing what is known as vertebrogenic pain.

Clinically, vertebrogenic pain is generally described as a midline, deep, aching, burning pain that is progressive. Also, it is often associated with an intermittent electrical shock sensation. Vertebrogenic pain is also characterized by absence of radicular expression, lower extremity weakness, or sensory deficits, and the neural tension sign and pain is generally worse with spinal flexion, sitting, standing and general physical activity, when compared to extension.

Diagnosis of vertebrogenic cLBP focuses on the chronic inflammatory response caused by endplate damage, which is visible on MRI. These signal changes, known as Modic changes (MC), are found in the vertebral body bone marrow that is adjacent to the degenerative endplates. Modic 1 changes indicate inflammation and edema, and Modic 2 changes occur in the setting of marrow ischemia when the red hematopoietic bone marrow has converted into yellow fatty marrow.

Thermal destruction (i.e., ablation) of the intraosseous BVN (Intracept® Procedure) is a therapeutic, interventional surgical procedure used to treat cLBP of vertebrogenic origin. The procedure is performed using fluoroscopic imaging under moderate/conscious sedation or general anesthesia. Radiofrequency energy is applied for 15 minutes at 85 degrees Celsius to produce a lesion to destroy the BVN within the vertebral body. At a minimum, the BVN is ablated in at least 1 vertebral body.

A systematic review of the published literature on the efficacy, effectiveness, and complications associated with BVN ablation for the treatment of cLBP was published by Conger, et al. (2021). Evidence was evaluated using the Grades of Recommendation, Assessment, Development and Evaluation (GRADE) system of appraisal to determine the quality of the evidence of the effectiveness of intraosseous BVN radiofrequency neurotomy (RFN). According to GRADE, there is moderate-quality evidence that BVN RFN is both an effective treatment compared to sham procedure and superior to continued standard care management for reduction of pain and disability in stringently selected patients with cLBP and corresponding Modic type 1 and 2 changes at a minimum of 3 months.

The International Society for the Advancement of Spine Surgery (ISASS) 2020 guideline – Intraosseous Ablation of the Basivertebral Nerve for the Relief of Chronic Low Back Pain concluded that "The procedure is supported by level 1 evidence including 2 RCTs demonstrating a statistically significant decrease in pain and an improvement in function with outcomes sustained to at least 24 months in a limited number of studies." BVN ablation may be indicated as a treatment option for cLBP for patients that fail nonsurgical treatment and their cLBP is diagnosed using well-established clinical and MRI findings.

The American Society of Pain and Neuroscience (ASPN) identified evidence-based guidelines from the available literature for the proper identification and selection of patients with vertebrogenic low back pain for BVN ablation. The systematic review was conducted using United States Preventive Services Task Force Criteria Modified for Interventional Spine Procedures and assigned a Grade A rating of the quality of evidence for BVN ablation indicating a high certainty that the net benefit is substantial in appropriately selected individuals.

Additional reviews are supportive of BVN ablation as an effective treatment and more beneficial than current standard of care for a subset of patients with cLBP based on moderate-quality evidence.

cLBP is a significant clinical problem and a socioeconomic burden that affects approximately 30 million people in the United States. Basic science and clinical trials have validated the diagnosis of vertebrogenic pain. The associations of Modic changes (MC) types 1 and 2 with cLBP and vertebral body endplate (VEP) injury have also been established.

DRAFT

Promising short- and long-term follow-up results support the safety, durability, and efficacy of radiofrequency ablation of the BVN for a subset of patients with vertebrogenic cLBP, corresponding with Modic change types 1 and 2 in the L3-S1 VEPs who have failed to respond to conservative treatment for at least 6 months. Published RCTs as well as prospective pre-post intervention studies have resulted in moderate-quality evidence that BVN RFN is safe and both an effective treatment compared to sham procedure and superior to continued standard care management for reduction of pain and disability when proper patient selection and procedural techniques are implemented.

Therefore, thermal destruction of the intraosseous BVN for vertebrogenic cLBP is considered to be medically reasonable and necessary when performed as outlined in the current published literature.
For individuals who have discogenic back pain who receive intradiscal electrothermal annuloplasty, the evidence includes a small number of randomized controlled trials (RCTs). Relevant outcomes are symptoms, functional outcomes, quality of life (QOL), and treatment-related morbidity. Two RCTs on intradiscal electrothermal annuloplasty reported conflicting results, with 1 reporting benefit for intradiscal electrothermal annuloplasty and the other reporting no benefit. Further study in a sham-controlled trial with a representative population of patients is needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have discogenic back pain who receive intradiscal radiofrequency annuloplasty, the evidence includes 2 RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. Neither RCT found evidence of benefit with the treatment. More sham-controlled trials are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have discogenic back pain who receive intradiscal biacuplasty, the evidence includes 2 industry-sponsored RCTs. Relevant outcomes are symptoms, functional outcomes, QOL, and treatment-related morbidity. One trial reported significant improvements at 6 months post-treatment, but not at 1 and 3months. The other trial also showed a significant reduction in visual analog scale scores at 6 months that appeared to continue to the 12-month follow-up; however, it is unclear whether this trial was sufficiently powered. More sham-controlled trials are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have herniated intervertebral disc(s) who receive automated percutaneous discectomy, the evidence includes randomized controlled trials (RCTs) and systematic reviews of observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The published evidence from small RCTs is insufficient to evaluate the impact of automated percutaneous discectomy on the net health outcome. Well-designed and executed RCTs are needed to determine the benefits and risks of this procedure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have discogenic back pain or radiculopathy who receive laser discectomy, the evidence includes systematic reviews of observational studies. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. While numerous case series and uncontrolled studies have reported improvements in pain levels and functioning following laser discectomy, the lack of well-designed and conducted controlled trials limits the interpretation of reported data. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have discogenic back pain or radiculopathy who receive disc nucleoplasty with radiofrequency coblation, the evidence includes randomized controlled trials (RCTs), systematic reviews, and prospective and retrospective nonrandomized studies. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. For nucleoplasty, there are 3 RCTs in addition to several uncontrolled studies. These RCTs are limited by the lack of blinding, an inadequate control condition in 1, inadequate data reporting in the second, and low enrollment with early study termination in the third. The available evidence is insufficient to permit conclusions concerning the effect of these procedures on health

DRAFT

outcomes due to multiple confounding factors that may bias results. High-quality randomized trials with adequate follow-up (at least 1year), which control for selection bias, the placebo effect, and variability in the natural history of low back pain, are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

CODING

Medicare Advantage Plans

The following CPT code(s) are covered for Medicare Advantage Plans and prior authorization is not required:

62330 Decompression, percutaneous, with partial removal of the ligamentum flavum, including laminotomy for access, epidurography, and imaging guidance (ie, ct or fluoroscopy), bilateral; one interspace, lumbar (New Code Effective 1/1/2026)

62331 Decompression, percutaneous, with partial removal of the ligamentum flavum, including laminotomy for access, epidurography, and imaging guidance (ie, ct or fluoroscopy), bilateral; additional interspace(s), lumbar (list separately in addition to code for primary procedure) (New Code Effective 1/1/2026)

The following CPT codes may be considered medically necessary for Medicare Advantage Plans when the medical criteria in the online authorization tool for participating providers is met:

64628 Thermal destruction of intraosseous basivertebral nerve, including all imaging guidance; first 2 vertebral bodies, lumbar or sacral

64629 Thermal destruction of intraosseous basivertebral nerve, including all imaging guidance; each additional vertebral body, lumbar or sacral (List separately in addition to code for primary procedure)

The following CPT/HCPCS code(s) are not covered for Medicare Advantage Plans:

62287 Decompression, percutaneous, of nucleus pulposus of intervertebral disc, any method utilizing needle-based technique to remove disc material under fluoroscopic imaging or other form of indirect visualization, with discography and/or epidural injection(s) at the treated level(s), when performed, single or multiple levels, lumbar (Text Revision Effective 1/1/2026)

S2348 Decompression procedure, percutaneous, of nucleus pulposus of intervertebral disc, using radiofrequency energy, single or multiple levels, lumbar

Commercial Products

The following CPT code(s) are considered medically necessary for Commercial Products when the above medical criteria have been met:

64628 Thermal destruction of intraosseous basivertebral nerve, including all imaging guidance; first 2 vertebral bodies, lumbar or sacral

64629 Thermal destruction of intraosseous basivertebral nerve, including all imaging guidance; each additional vertebral body, lumbar or sacral (List separately in addition to code for primary procedure)

The following code(s) are not medically necessary for Commercial Products:

62287 Decompression, percutaneous, of nucleus pulposus of intervertebral disc, any method utilizing needle-based technique to remove disc material under fluoroscopic imaging or other form of indirect visualization, with discography and/or epidural injection(s) at the treated level(s), when performed, single or multiple levels, lumbar (Text Revision Effective 1/1/2026)

62330 Decompression, percutaneous, with partial removal of the ligamentum flavum, including laminotomy for access, epidurography, and imaging guidance (ie, ct or fluoroscopy), bilateral; one interspace, lumbar (New Code Effective 1/1/2026)

62331 Decompression, percutaneous, with partial removal of the ligamentum flavum, including laminotomy for access, epidurography, and imaging guidance (ie, ct or fluoroscopy), bilateral; additional interspace(s), lumbar (list separately in addition to code for primary procedure) (New Code Effective 1/1/2026)

S2348 Decompression procedure, percutaneous, of nucleus pulposus of intervertebral disc, using radiofrequency energy, single or multiple levels, lumbar

DRAFT

Medicare Advantage Plans and Commercial Products

The following CPT code(s) are not covered for Medicare Advantage Plans, when identified as Percutaneous Thermal Intradiscal Procedures (TIPs), and not medically necessary for Commercial Products:

22526 Percutaneous intradiscal electrothermal annuloplasty, unilateral or bilateral including fluoroscopic guidance; single level

22527 Percutaneous intradiscal electrothermal annuloplasty, unilateral or bilateral including fluoroscopic guidance; one or more additional levels (List separately in addition to code for primary procedure)

RELATED POLICIES

Centers for Medicare and Medicaid Services (CMS) National and Local Coverage Determinations
Prior Authorization of Services, Treatments or Procedures

PUBLISHED

Provider Update, February/October 2026
Provider Update, January 2025
Provider Update, August/December 2023
Provider Update, November 2022
Provider Update, September 2021

REFERENCES

1. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Laser Procedures (140.5). 1997; https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCDId=69&ncdver=1&DocID=140.5&bc=gAAAAAgAAAAAAA%3D%3.
2. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Thermal Intradiscal Procedures (TIPs) (150.11). 2009;https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCDId=324.
3. Centers for Medicare and Medicaid Services (CMS). Palmetto Local Coverage Determination (LCD) for Thermal Destruction of The Intraosseous Basivertebral Nerve (BVN) for Vertebrogenic Lower Back Pain (L39420). Accessed October 22, 2025.
4. Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. Spine J. Jun 01 2015; 15(6): 1461-77. PMID 24412033
5. Haines SJ, Jordan N, Boen JR, et al. Discectomy strategies for lumbar disc herniation: results of the LAPDOG trial. J Clin Neurosci. Jul 2002; 9(4): 411-7. PMID 12217670
6. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniquesin chronic spinal pain. Part II: guidance and recommendations. Pain Physician. Apr 2013; 16(2 Suppl): S49-283. PMID23615883
7. Zhao XM, Chen AF, Lou XX, et al. Comparison of Three Common Intervertebral Disc Discectomies in the Treatment of Lumbar Disc Herniation: A Systematic Review and Meta-Analysis Based on Multiple Data. J Clin Med. Nov 08 2022; 11(22). PMID 36431083
8. Bai X, Lian Y, Wang J, et al. Percutaneous endoscopic lumbar discectomy compared with other surgeries for lumbar disc herniation: A meta-analysis. Medicine (Baltimore). Mar 05 2021; 100(9): e24747. PMID 33655938
9. Gadjradj PS, Harhangi BS, Amelink J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus Open Microdiscectomy for Lumbar Disc Herniation: A Systematic Review and Meta-analysis. Spine (Phila Pa 1976). Apr 15 2021; 46(8): 538-549. PMID 33290374
10. Xu J, Li Y, Wang B, et al. Minimum 2-Year Efficacy of Percutaneous Endoscopic Lumbar Discectomy versus Microendoscopic Discectomy: A Meta-Analysis. World Neurosurg. Jun 2020; 138: 19-26. PMID 32109644
11. Phan K, Xu J, Schultz K, et al. Full-endoscopic versus micro-endoscopic and open discectomy: A systematic review andmeta-analysis of outcomes and complications. Clin Neurol Neurosurg. Mar 2017; 154: 1-12. PMID 28086154

DRAFT

12. Yu P, Qiang H, Zhou J, et al. Percutaneous Transforaminal Endoscopic Discectomy versus Micro-Endoscopic Discectomyfor Lumbar Disc Herniation. Med Sci Monit. Mar 30 2019; 25: 2320-2328. PMID 30927349
13. Zhao XM, Yuan QL, Liu L, et al. Is It Possible to Replace Microendoscopic Discectomy with Percutaneous TransforaminalDiscectomy for Treatment of Lumbar Disc Herniation? A Meta-Analysis Based on Recurrence and Revision Rate. JKorean Neurosurg Soc. Jul 2020; 63(4): 477-486. PMID 32380585
14. Yu P, Qiang H, Zhou J, et al. Percutaneous Transforaminal Endoscopic Discectomy versus Micro-Endoscopic Discectomy for Lumbar Disc Herniation. Med Sci Monit. Mar 30 2019; 25: 2320-2328. PMID 30927349
15. Shi R, Wang F, Hong X, et al. Comparison of percutaneous endoscopic lumbar discectomy versus microendoscopic discectomy for the treatment of lumbar disc herniation: a meta-analysis. Int Orthop. Apr 2019; 43(4): 923-937. PMID 30547214
16. Gadjradj PS, Harhangi BS, Amelink J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus OpenMicrodiscectomy for Lumbar Disc Herniation: A Systematic Review and Meta-analysis. Spine (Phila Pa 1976). Apr 152021; 46(8): 538-549. PMID 33290374
17. Ran B, Wei J, Yang J, et al. Quantitative Evaluation of the Trauma of CT Navigation PELD and OD in the Treatment of HLDH: A Randomized, Controlled Study. Pain Physician. Jul 2021; 24(4): E433-E441. PMID 34213868
18. Zhao XM, Chen AF, Lou XX, et al. Comparison of Three Common Intervertebral Disc Discectomies in the Treatment ofLumbar Disc Herniation: A Systematic Review and Meta-Analysis Based on Multiple Data. J Clin Med. Nov 08 2022;11(22). PMID 36431083
19. Wang SF, Hung SF, Tsai TT, et al. Better Functional Outcome and Pain Relief in the Far-Lateral-Outside-in Percutaneous Endoscopic Transforaminal Discectomy. J Pain Res. 2021; 14: 3927-3934. PMID 35002312
20. Rajamani PA, Goparaju P, Kulkarni AG, et al. A 2-Year Outcomes and Complications of Various Techniques of Lumbar Discectomy: A Multicentric Retrospective Study. World Neurosurg. Dec 2021; 156: e319-e328. PMID 34555576
21. Jing Z, Li L, Song J. Percutaneous transforaminal endoscopic discectomy versus micro endoscopic discectomy for upper lumbar disc herniation: a retrospective comparative study. Am J Transl Res. 2021; 13(4): 3111-3119. PMID 34017479
22. Chen Z, Zhang L, Dong J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus Microendoscopic Discectomy for Lumbar Disc Herniation: Two-Year Results of a Randomized Controlled Trial. Spine (Phila Pa 1976). Apr15 2020; 45(8): 493-503. PMID 31703056
23. Kim CH, Chung CK, Choi Y, et al. The Long-term Reoperation Rate Following Surgery for Lumbar Herniated Intervertebral Disc Disease: A Nationwide Sample Cohort Study With a 10-year Follow-up. Spine (Phila Pa 1976). Oct 01 2019; 44(19): 1382-1389. PMID 30973508
24. Ahn Y, Lee SG, Son S, et al. Transforaminal Endoscopic Lumbar Discectomy Versus Open Lumbar Microdiscectomy: A Comparative Cohort Study with a 5-Year Follow-Up. Pain Physician. May 2019; 22(3): 295-304. PMID 31151337
25. Akçakaya MO, Yörükoğlu AG, Aydoseli A, et al. Serum creatine phosphokinase levels as an indicator of muscle injury following lumbar disc surgery: Comparison of fully endoscopic discectomy and microdiscectomy. Clin Neurol Neurosurg. Jun 2016; 145: 74-8. PMID 27101087
26. Choi KC, Shim HK, Hwang JS, et al. Comparison of Surgical Invasiveness Between Microdiscectomy and 3 Different Endoscopic Discectomy Techniques for Lumbar Disc Herniation. World Neurosurg. Aug 2018; 116: e750-e758. PMID 29787880
27. Dai HJ, Zhang X, Wang LT, et al. The effect of percutaneous transforaminal endoscopic discectomy (PTED) on serum inflammatory factors and pain in patients with lumbar disc herniation after surgery. Int J Clin Exp Med 2020;13:597603.
28. Krappel FA, Schmitz R, Bauer E, et al. Open or endoscopic nucleotomy?. Orthopadische Praxis 2001;37:1649.
29. Tacconi L, Giordan E. Endoscopic transforaminal discectomy vs. far lateral discectomy for extraforaminal disc protrusions: our experience. NeuroQuantology 2019;17:1822.

30. Tacconi L, Signorelli F, Giordan E. Is Full Endoscopic Lumbar Discectomy Less Invasive Than Conventional Surgery? A Randomized MRI Study. World Neurosurg. Jun 2020; 138: e867-e875. PMID 32251813
31. Tao XZ, Jing L, Li JH. Therapeutic effect of transforaminal endoscopic spine system in the treatment of prolapse of lumbar intervertebral disc. Eur Rev Med Pharmacol Sci. Jul 2018; 22(1 Suppl): 103-110. PMID 30004561
32. Wang H, Song Y, Cai L. Effect of percutaneous transforaminal lumbar spine endoscopic discectomy on lumbar disc herniation and its influence on indexes of oxidative stress. Biomed Res 2017;28:.
33. Xu G, Zhang C, Zhu K, et al. Endoscopic removal of nucleus pulposus of intervertebral disc on lumbar intervertebral disc protrusion and the influence on inflammatory factors and immune function. Exp Ther Med. Jan 2020; 19(1): 301-307. PMID 31853303
34. Pan Z, Ha Y, Yi S, et al. Efficacy of Transforaminal Endoscopic Spine System (TESSYS) Technique in Treating Lumbar Disc Herniation. Med Sci Monit. Feb 18 2016; 22: 530-9. PMID 26887645
35. Gibson JNA, Subramanian AS, Scott CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. Eur Spine J. Mar 2017; 26(3): 847-856. PMID 27885470
36. Hsu HT, Chang SJ, Yang SS, et al. Learning curve of full-endoscopic lumbar discectomy. Eur Spine J. Apr 2013; 22(4): 727-33. PMID 23076645
37. Kim MJ, Lee SH, Jung ES, et al. Targeted percutaneous transforaminal endoscopic diskectomy in 295 patients: comparison with results of microscopic diskectomy. Surg Neurol. Dec 2007; 68(6): 623-631. PMID 18053857
38. Qu JX, Li QZ, Chen M : Comparative study of PTED and MED for monosegmentnlumbar disc herniation. Chin J Bone Joint Inj 32 : 70-71,2017
39. Wang H, Cheng J, Xiao H, et al. Adolescent lumbar disc herniation: experience from a large minimally invasive treatment centre for lumbar degenerative disease in Chongqing, China. Clin Neurol Neurosurg. Aug 2013; 115(8): 1415-9. PMID 23419406
40. Zhao W, Li CQ, Zhou Y, Wang J, Zheng WJ : Surgical treatment of thenlumbar disc herniated discs using transforaminal endoscopic surgerynsystem. Orthop J China 20 : 1191-1195, 2012
41. Li M, Yang H, Yang Q. Full-Endoscopic Technique Discectomy Versus Microendoscopic Discectomy for the Surgical Treatment of Lumbar Disc Herniation. Pain Physician. 2015; 18(4): 359-63. PMID 26218939
42. Sinkemani A, Hong X, Gao ZX, et al. Outcomes of Microendoscopic Discectomy and Percutaneous Transforaminal Endoscopic Discectomy for the Treatment of Lumbar Disc Herniation: A Comparative Retrospective Study. Asian Spine J. Dec 2015; 9(6): 833-40. PMID 26713113
43. Song HP, Sheng HF, Xu WX. A case-control study on the treatment of protrusion of lumbar intervertebral disc through PELD and MED. Exp Ther Med. Oct 2017; 14(4): 3708-3712. PMID 29042967
44. Lee DY, Lee SH. Learning curve for percutaneous endoscopic lumbar discectomy. Neurol Med Chir (Tokyo). Sep 2008; 48(9): 383-8; discussion 388-9. PMID 18812679
45. Lee SH, Chung SE, Ahn Y, et al. Comparative radiologic evaluation of percutaneous endoscopic lumbar discectomy and open microdiscectomy: a matched cohort analysis. Mt Sinai J Med. Sep 2006; 73(5): 795-801. PMID 17008941
46. Wang B, Lü G, Patel AA, et al. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. Spine J. Feb 2011; 11(2): 122-30. PMID 21296295
47. Tu Z, Li YW, Wang B, et al. Clinical Outcome of Full-endoscopic Interlaminar Discectomy for Single-level Lumbar Disc Herniation: A Minimum of 5-year Follow-up. Pain Physician. Mar 2017; 20(3): E425-E430. PMID 28339442
48. Li H, Jiang C, Mu X, et al. Comparison of MED and PELD in the Treatment of Adolescent Lumbar Disc Herniation: A 5-Year Retrospective Follow-Up. World Neurosurg. Apr 2018; 112: e255-e260. PMID 29325949
49. Chen Z, Zhang L, Dong J, et al. Percutaneous transforaminal endoscopic discectomy compared with microendoscopic discectomy for lumbar disc herniation: 1-year results of an ongoing randomized controlled trial. J Neurosurg Spine. Mar 2018; 28(3): 300-310. PMID 29303469

DRAFT

50. Liu T, Zhou Y, Wang J, et al. Clinical efficacy of three different minimally invasive procedures for far lateral lumbar disc herniation. Chin Med J (Engl). Mar 2012; 125(6): 1082-8. PMID 22613535
51. Wu XC, Zhou Y, Li CQ. Percutaneous tranforaminal endoscopic discectomy versus microendoscopic discectomy for lumbar disc herniation: a prospective randomized controlled study. J Third Mil Med Univ. 2009;31(9):843-846.
52. Yang L, Liao XQ, Zhao XJ, et al. Comparison of surgical outcomes between percutaneous transforaminal endoscopic discectomy and micro-endoscopic discectomy for lumbar disc herniation. China J Endosc. 2015;21(9):962-965
53. Li ZY, Guo PG, Han D, et al. Analysis of curative effects and prognosis in different procedures of discectomy for patients with lumbar disc herniation. J Clin Med Pract. 2017;21(15):149-150,158
54. Liu HP, Hao DJ, Wang XD, et al. Comparison of two surgeries in treatment of lumbar disc herniation. Chin J Pain Med. 2017;23(6):438-442
55. Luo DK, Zhou NX, Zhao HW, et al. Clinical effectiveness of minimally invasive treatment for lumbar disc herniation. Orthopaedics. 2017;8(6):439-444
56. Qu JX, Li QZ, Chem M, et al. Comparison of the efficacies between percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy for the treatment of single-segmental lumbar disc herniation. Chin J Bone Jt Inj. 2017;32(1):70-71
57. Chen Q, Qin L, Li MW, et al. Comparison of the therapeutic effect of percutaneous transforaminal endoscopic discectomy and posterior discectomy on senile single segmental lumbar disc herniation. Chin J Front Med Sci. 2018;10(2):60-64
58. Wu YM, Bai M, Yin HP, et al. Comparison of the efficacies between two kinds of minimally invasive procedures for the treatment of simple lumbar disc herniation. J Pract Orthop. 2018;24(4):357-360
59. Belykh E, Giers MB, Preul MC, et al. Prospective Comparison of Microsurgical, Tubular-Based Endoscopic, and Endoscopically Assisted Diskectomies: Clinical Effectiveness and Complications in Railway Workers. World Neurosurg. Jun 2016; 90: 273-280. PMID 26898494
60. Garg B, Nagraja UB, Jayaswal A. Microendoscopic versus open discectomy for lumbar disc herniation: a prospective randomised study. J Orthop Surg (Hong Kong). Apr 2011; 19(1): 30-4. PMID 21519072
61. Hermantin FU, Peters T, Quartararo L, et al. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. J Bone Joint Surg Am. Jul 1999; 81(7): 958-65. PMID 10428127
62. Huang TJ, Hsu RW, Li YY, et al. Less systemic cytokine response in patients following micro endoscopic versus open lumbar discectomy. J Orthop Res. Mar 2005; 23(2): 406-11. PMID 15734255
63. Hussein M, Abdeldayem A, Mattar MM. Surgical technique and effectiveness of microendoscopic discectomy for large uncontained lumbar disc herniations: a prospective, randomized, controlled study with 8 years of follow-up. Eur Spine J. Sep 2014; 23(9): 1992-9. PMID 24736930
64. Martín-Láez R, Martínez-Agüeros JA, Suárez-Fernández D, et al. Complications of endoscopic microdiscectomy using the EASYGO! system: is there any difference with conventional discectomy during the learning-curve period?. Acta Neurochir (Wien). Jun 2012; 154(6): 1023-32. PMID 22446750
65. Ohya J, Oshima Y, Chikuda H, et al. Does the microendoscopic technique reduce mortality and major complications in patients undergoing lumbar discectomy? A propensity score-matched analysis using a nationwide administrative database. Neurosurg Focus. Feb 2016; 40(2): E5. PMID 26828886
66. Pan L, Zhang P, Yin Q. Comparison of tissue damages caused by endoscopic lumbar discectomy and traditional lumbar discectomy: a randomised controlled trial. Int J Surg. 2014; 12(5): 534-7. PMID 24583364
67. Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. Neurosurgery. Sep 2007; 61(3): 545-9; discussion 549. PMID 17881967
68. Ruetten S, Komp M, Merk H, et al. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. Spine (Phila Pa 1976). Apr 20 2008; 33(9): 931-9. PMID 18427312
69. Ruetten S, Komp M, Merk H, et al. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision. J Spinal Disord Tech. Apr 2009; 22(2): 122-9. PMID 19342934

DRAFT

70. Sasaoka R, Nakamura H, Konishi S, et al. Objective assessment of reduced invasiveness in MED. Compared with conventional one-level laminotomy. Eur Spine J. May 2006; 15(5): 577-82. PMID 15926058
71. Schizas C, Tsiridis E, Saksena J. Microendoscopic discectomy compared with standard microsurgical discectomy for treatment of uncontained or large contained disc herniations. Neurosurgery. Oct 2005; 57(4 Suppl): 357-60; discussion 357-60. PMID 16234685
72. Teli M, Lovi A, Brayda-Bruno M, et al. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy. Eur Spine J. Mar 2010; 19(3): 443-50. PMID 20127495
73. Ruetten S, Komp M, Merk H, et al. Use of newly developed instruments and endoscopes: full-endoscopic resection of lumbar disc herniations via the interlaminar and lateral transforaminal approach. J Neurosurg Spine. Jun 2007; 6(6): 521-30. PMID 17561740
74. Ruetten S, Komp M, Merk H, et al. Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9-mm endoscopes: a prospective, randomized, controlled study. Spine (Phila Pa 1976). Apr 20 2008; 33(9): 940-8. PMID 18427313
75. Gadjradj PS, Rubinstein SM, Peul WC, et al. Full endoscopic versus open discectomy for sciatica: randomised controlled non-inferiority trial. BMJ. Feb 21 2022; 376: e065846. PMID 35190388
76. Ran B, Wei J, Yang J, et al. Quantitative Evaluation of the Trauma of CT Navigation PELD and OD in the Treatment of HLDH: A Randomized, Controlled Study. Pain Physician. Jul 2021; 24(4): E433-E441. PMID 34213868
77. Wang F, Guo D, Sun T, et al. A comparative study on short-term therapeutic effects of percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy on lumbar disc herniation. Pak J Med Sci. 2019; 35(2): 426-431. PMID 31086527
78. Liu Y, Kim Y, Park CW, et al. Interlaminar Endoscopic Lumbar Discectomy Versus Microscopic Lumbar Discectomy: A Preliminary Analysis of L5-S1 Lumbar Disc Herniation Outcomes in Prospective Randomized Controlled Trials. Neurospine. Dec 2023; 20(4): 1457-1468. PMID 38171312
79. Yang X, Zhang S, Su J, et al. Comparison of Clinical and Radiographic Outcomes Between Transforaminal Endoscopic Lumbar Discectomy and Microdiscectomy: A Follow-up Exceeding 5 Years. Neurospine. Mar 2024; 21(1): 303-313. PMID 38317550
80. Saghebdoust S, Khadivar F, Ekrami M, et al. Transforaminal Endoscopic Lumbar Diskectomy versus Open Microdiskectomy for Symptomatic Lumbar Disk Herniation: A Comparative Cohort Study on Costs and Long-Term Outcomes. J Neurol Surg A Cent Eur Neurosurg. Oct 25 2023. PMID 37879346
81. Wang B, Lü G, Patel AA, et al. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. Spine J. Feb 2011; 11(2): 122-30. PMID 21296295
82. Tenenbaum S, Arzi H, Herman A, et al. Percutaneous Posterolateral Transforaminal Endoscopic Discectomy: Clinical Outcome, Complications, and Learning Curve Evaluation. Surg Technol Int. Dec 2011; 21: 278-83. PMID 22505002
83. Casal-Moro R, Castro-Menéndez M, Hernández-Blanco M, et al. Long-term outcome after microendoscopic diskectomy for lumbar disk herniation: a prospective clinical study with a 5-year follow-up. Neurosurgery. Jun 2011; 68(6): 1568-75; discussion 1575. PMID 21311384
84. Wang M, Zhou Y, Wang J, et al. A 10-year follow-up study on long-term clinical outcomes of lumbar microendoscopic discectomy. J Neurol Surg A Cent Eur Neurosurg. Aug 2012; 73(4): 195-8. PMID 22825836
85. National Institute for Health and Care Excellence (NICE). Automated percutaneous mechanical lumbar discectomy-guidance [IPG141]. 2005; http://guidance.nice.org.uk/IPG141/Guidance/pdf/English. Accessed April 30, 2025.
86. National Institute for Health and Care Excellence (NICE). Percutaneous interlaminar endoscopic lumbar discectomy for sciatica [IPG555]. 2016; https://www.nice.org.uk/guidance/ipg555. Accessed April 29, 2025.
87. National Institute for Health and Care Excellence (NICE). Percutaneous transforaminal endoscopic lumbar discectomy for sciatica [IPG556]. 2016; https://www.nice.org.uk/guidance/ipg556. Accessed April 28, 2025.
88. Kreiner DS, Hwang SW, Easa JE, et al. An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. Spine J. Jan 2014; 14(1): 180-91. PMID 24239490

89. Ma C, Li H, Zhang T, et al. Comparison of Percutaneous Endoscopic Interlaminar Discectomy and Open Fenestration Discectomy for Single-Segment Huge Lumbar Disc Herniation: A Two-year Follow-up Retrospective Study. J Pain Res. 2022; 15: 1061-1070. PMID 35444463
90. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. Spine (Phila Pa 1976). May 01 2009; 34(10): 1066-77. PMID 19363457
91. Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain. J Pain Res. 2022; 15: 3729-3832. PMID 36510616
92. Singh V, Manchikanti L, Benyamin RM, et al. Percutaneous lumbar laser disc decompression: a systematic review of current evidence. Pain Physician. 2009; 12(3): 573-88. PMID 19461824
93. Singh V, Manchikanti L, Calodney AK, et al. Percutaneous lumbar laser disc decompression: an update of current evidence. Pain Physician. Apr 2013; 16(2 Suppl): SE229-60. PMID 23615885
94. Gibson JN, Waddell G. Surgical interventions for lumbar disc prolapse. Cochrane Database Syst Rev. Apr 18 2007; 2007(2): CD001350. PMID 17443505
95. Seddighi AS, Seddighi A, Hosseini S. Clinical Outcomes at 2-Year Follow-Up Comparing Open Surgery and Percutaneous Laser Disc Decompression for Radicular Sciatic Pain Patients. World Neurosurg. Feb 2025; 194: 123392. PMID 39491619
96. Tassi GP. Comparison of results of 500 microdiscectomies and 500 percutaneous laser disc decompression procedures for lumbar disc herniation. Photomed Laser Surg. Dec 2006; 24(6): 694-7. PMID 17199468
97. Choy DS. Percutaneous laser disc decompression: an update. Photomed Laser Surg. Oct 2004; 22(5): 393-406. PMID 15671712
98. Menchetti PP, Canero G, Bini W. Percutaneous laser discectomy: experience and long term follow-up. Acta Neurochir Suppl. 2011; 108: 117-21. PMID 21107947
99. Manchikanti L, Falco FJ, Benyamin RM, et al. An update of the systematic assessment of mechanical lumbar disc decompression with nucleoplasty. Pain Physician. Apr 2013; 16(2 Suppl): SE25-54. PMID 23615886
100. Gerszten PC, Smuck M, Rathmell JP, et al. Plasma disc decompression compared with fluoroscopy-guided transforaminal epidural steroid injections for symptomatic contained lumbar disc herniation: a prospective, randomized, controlled trial. J Neurosurg Spine. Apr 2010; 12(4): 357-71. PMID 20201654
101. Chitragran R, Poopitaya S, Tassanawipas W. Result of percutaneous disc decompression using nucleoplasty in Thailand: a randomized controlled trial. J Med Assoc Thai. Oct 2012; 95 Suppl 10: S198-205. PMID 23451463
102. de Rooij J, Harhangi B, Aukes H, et al. The Effect of Percutaneous Nucleoplasty vs Anterior Discectomy in Patients with Cervical Radicular Pain due to a Single-Level Contained Soft-Disc Herniation: A Randomized Controlled Trial. Pain Physician. Nov 2020; 23(6): 553-564. PMID 33185372
103. Chen CH, Chiu YP, Ji HR, et al. Analysis of the clinical and radiological outcomes of percutaneous cervical nucleoplasty: A case-control study. PLoS One. 2022; 17(12): e0278883. PMID 36508407
104. Bokov A, Skorodumov A, Isrelov A, et al. Differential treatment of nerve root compression pain caused by lumbar disc herniation applying nucleoplasty. Pain Physician. 2010; 13(5): 469-80. PMID 20859316
105. Birnbaum K. Percutaneous cervical disc decompression. Surg Radiol Anat. Jun 2009; 31(5): 379-87. PMID 19190848
106. Cuellar VG, Cuellar JM, Vaccaro AR, et al. Accelerated degeneration after failed cervical and lumbar nucleoplasty. J Spinal Disord Tech. Dec 2010; 23(8): 521-4. PMID 21131800
107. Manchikanti L, Derby R, Benyamin RM, et al. A systematic review of mechanical lumbar disc decompression with nucleoplasty. Pain Physician. 2009; 12(3): 561-72. PMID 19461823
108. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. Pain Physician. Apr 2013; 16(2 Suppl): S49-283. PMID 23615883
109. National Institute for Health and Care Excellence (NICE). Percutaneous coblation of the intervertebral disc for low back pain and sciatica [IPG543]. 2016; https://www.nice.org.uk/guidance/ipg543. Accessed February 19, 2025.

DRAFT

110. North American Spine Society. Clinical guidelines for diagnosis and treatment of lumbar disc herniation with radiculopathy. 2012; https://www.spine.org/Portals/0/Assets/Downloads/ResearchClinicalCare/Guidelines/LumbarDiscHerniation.pdf. Accessed February 19, 2025.
111. U.S. Food & Drug Administration. K213836 Intracept Intraosseous Nerve Ablation System 510k Summary. 2022. https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K213836. Accessed March 1, 2025.
112. Pauza KJ, Howell S, Dreyfuss P, et al. A randomized, placebo-controlled trial of intradiscal electrothermal therapy for the treatment of discogenic low back pain. Spine J. 2004; 4(1): 27-35. PMID 14749191
113. Freeman BJ, Fraser RD, Cain CM, et al. A randomized, double-blind, controlled trial: intradiscal electrothermal therapy versus placebo for the treatment of chronic discogenic low back pain. Spine (Phila Pa 1976). Nov 01 2005; 30(21): 2369-77; discussion 2378. PMID 16261111
114. Barendse GA, van Den Berg SG, Kessels AH, et al. Randomized controlled trial of percutaneous intradiscal radiofrequency thermocoagulation for chronic discogenic back pain: lack of effect from a 90-second 70 C lesion. Spine (Phila Pa 1976). Feb 01 2001; 26(3): 287-92. PMID 11224865
115. Kvarstein G, Måwe L, Indahl A, et al. A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy--a 12-month follow-up. Pain. Oct 2009; 145(3): 279-286. PMID 19647940
116. Kapural L, Vrooman B, Sarwar S, et al. A randomized, placebo-controlled trial of transdiscal radiofrequency, biacuplasty for treatment of discogenic lower back pain. Pain Med. Mar 2013; 14(3): 362-73. PMID 23279658
117. Kapural L, Vrooman B, Sarwar S, et al. Radiofrequency intradiscal biacuplasty for treatment of discogenic lower back pain: a 12-month follow-up. Pain Med. Mar 2015; 16(3): 425-31. PMID 25339501
118. Desai MJ, Kapural L, Petersohn JD, et al. A Prospective, Randomized, Multicenter, Open-label Clinical Trial Comparing Intradiscal Biacuplasty to Conventional Medical Management for Discogenic Lumbar Back Pain. Spine (Phila Pa 1976). Jul 01 2016; 41(13): 1065-1074. PMID 26689579
119. Desai MJ, Kapural L, Petersohn JD, et al. Twelve-Month Follow-up of a Randomized Clinical Trial Comparing Intradiscal Biacuplasty to Conventional Medical Management for Discogenic Lumbar Back Pain. Pain Med. Apr 01 2017; 18(4): 751-763. PMID 27570246
120. Fischgrund JS, Rhyne A, Franke J, et al. Intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: a prospective randomized double-blind sham-controlled multi-center study. Eur Spine J. May 2018; 27(5): 1146-1156. PMID 29423885
121. Fischgrund JS, Rhyne A, Franke J, et al. Intraosseous Basivertebral Nerve Ablation for the Treatment of Chronic Low Back Pain: 2-Year Results From a Prospective Randomized Double-Blind Sham-Controlled Multicenter Study. Int J Spine Surg. Apr 2019; 13(2): 110-119. PMID 31131209
122. Fischgrund JS, Rhyne A, Macadaeg K, et al. Long-term outcomes following intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: 5-year treatment arm results from a prospective randomized double-blind sham-controlled multi-center study. Eur Spine J. Aug 2020; 29(8): 1925-1934. PMID 32451777
123. Khalil JG, Smuck M, Koreckij T, et al. A prospective, randomized, multicenter study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain. Spine J. Oct 2019; 19(10): 1620-1632. PMID 31229663
124. Smuck M, Khalil J, Barrette K, et al. Prospective, randomized, multicenter study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: 12-month results. Reg Anesth Pain Med. Aug 2021; 46(8): 683-693. PMID 34031220
125. Koreckij T, Kreiner S, Khalil JG, et al. Prospective, randomized, multicenter study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: 24-Month treatment arm results. N Am Spine Soc J. Dec 2021; 8: 100089. PMID 35141653
126. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. Pain Physician. Apr 2013; 16(2 Suppl): S49-283. PMID 23615883
127. Boswell MV, Trescot AM, Datta S, et al. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. Pain Physician. Jan 2007; 10(1): 7-111. PMID 17256025

DRAFT

128.Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain. J Pain Res. 2022; 15: 3729-3832. PMID 36510616
129.Lorio M, Clerk-Lamalice O, Rivera M, et al. ISASS Policy Statement 2022: Literature Review of Intraosseous Basivertebral Nerve Ablation. Int J Spine Surg. Dec 2022; 16(6): 1084-1094. PMID 36266051
130.National Institute for Health and Care Excellence. Percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain [IPG545]. 2016; https://www.nice.org.uk/guidance/ipg545. Accessed March 10, 2025.
131.National Institute for Health and Care Excellence. Percutaneous electrothermal treatment of the intervertebral disc annulus for low back pain and sciatica [IPG544]. 2016; https://www.nice.org.uk/guidance/IPG544. Accessed March 9, 2025.

DRAFT

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.