Medical Coverage Policy | Whole Body Dual X-Ray Absorptiometry to Determine Body Composition

EFFECTIVE DATE: 03|01|2025
POLICY LAST REVIEWED: 10|07|2026

OVERVIEW
Using low-dose x-rays of 2 different energy levels, whole-body dual-energy x-ray absorptiometry (DXA) measures lean tissue mass, total and regional body fat, as well as bone density. DXA scans have become a tool for research on body composition (eg, as a more convenient replacement for underwater weighing). This evidence review addresses potential applications in clinical care rather than research use of the technology.

MEDICAL CRITERIA
Not applicable

PRIOR AUTHORIZATION
Not applicable

POLICY STATEMENT
Medicare Advantage Plans and Commercial Products
Dual-energy x-ray absorptiometry body composition studies are considered not covered for Medicare Advantage Plans and not medically necessary for Commercial Products as the evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

COVERAGE
Benefits may vary between groups/contracts. Please refer to the Evidence of Coverage or Subscriber Agreement for applicable not medically necessary/not covered Radiology/Diagnostic Imaging benefits/coverage.

BACKGROUND
Body Composition Measurement
Body composition measurements can be used to quantify and assess the relative proportions of specific body compartments such as fat and lean mass (eg, bones, tissues, organs, muscles). These measurements may be more useful in informing diagnosis, prognosis, or therapy than standard assessments (eg, body weight, body mass index) that do not identify the contributions of individual body compartments or their particular relationships with health and disease. While these body composition measurements have been most frequently utilized for research purposes, they may be useful in clinical settings to:

- Evaluate the health status of undernourished patients, those impacted by certain disease states (eg, anorexia nervosa, cachexia), or those undergoing certain treatments (eg, antiretroviral therapy, bariatric surgery).
- Evaluate the risk of heart disease or diabetes by measuring visceral fat versus total body fat.
- Assess body composition changes related to growth and development (eg, infancy, childhood), aging(eg, sarcopenia), and certain disease states (eg, HIV, diabetes).
- Evaluate patients in situations where body mass index is suspected to be discordant with total fat mass(eg, body-building, edema).

A variety of techniques have been researched, including most commonly, anthropomorphic measures, bioelectrical impedance, and dual-energy x-ray absorptiometry (DXA). All of these techniques are based in part on assumptions about the distribution of different body compartments and their density, and all rely on formulas to convert the measured parameter into an estimate of body composition. Therefore, all techniques

will introduce variation based on how the underlying assumptions and formulas apply to different populations of subjects (ie, different age groups, ethnicities, or underlying conditions). Techniques using anthropomorphics, bioelectrical impedance, underwater weighing, and DXA are briefly reviewed below.

Anthropomorphic Techniques
Anthropomorphic techniques for the estimation of body composition include measurements of skin fold thickness at various sites, bone dimensions, and limb circumference. These measurements are used in various equations to predict body density and body fat. Due to its ease of use, measurement of skinfold thickness is one of the most common techniques. The technique is based on the assumption that the subcutaneous adipose layer reflects total body fat but this association may vary with age and sex. Skinfold thickness measurement precision and utility can also be affected by operator experience and a lack of applicable reference data for specific patient populations or percentile extremes.

Bioelectrical Impedance
Bioelectrical impedance analysis is based on the relation between the volume of the conductor (ie, human body), the conductor's length (ie, height), the components of the conductor (ie, fat and fat-free mass), and its impedance. The technique involves attaching surface electrodes to various locations on the arm and foot. Alternatively, the patient can stand on pad electrodes. Estimates of body composition are based on the assumption that the overall conductivity of the human body is closely related to lean tissue. The impedance value is then combined with anthropomorphic data and certain other patient-specific parameters (eg, age, gender, ethnicity) to give body compartment measures. These measures are calculated based on device manufacturer-specific regression models, which are generally proprietary. Bioelectrical impedance measures can be affected by fat distribution patterns, hydration status, ovulation, and temperature.

Underwater Weighing
Underwater weighing requires the use of a specially constructed tank in which the subject is seated on a suspended chair. The subject is then submerged in the water while exhaling; the difference between weight in air and weight in water is used to estimate total body fat percentage. While valued as a research tool, weighing people underwater is typically not suitable for routine clinical use. This technique is based on the assumption that the body can be divided into 2 compartments with constant densities: adipose tissue, with a density of 0.9g/cm^3, and lean body mass (ie, muscle and bone), with a density of 1.1 g/cm^3. One limitation of the underlying assumption is the variability in density between muscle and bone; eg, bone has a higher density than muscle, and bone mineral density varies with age and other conditions. Also, the density of body fat may vary, depending on the relative components of its constituents (eg, glycerides, sterols, glycolipids).

Dual-energy X-ray Absorptiometry
While the cited techniques assume 2 body compartments, DXA can estimate 3 body compartments consisting of fat mass, lean body mass, and bone mass. DXA systems use a source that generates x-rays at 2 energies. The differential attenuation of the 2 energies is used to estimate the bone mineral content and soft tissue composition. When 2 x-ray energies are used, only 2 tissue compartments can be measured; therefore, soft tissue measurements (ie, fat and lean body mass) can only be measured in areas in which no bone is present. DXA can also determine body composition in defined regions (ie, the arms, legs, and trunk). DXA measurements are based in part on the assumption that the hydration of fat-free mass remains constant at73%. Hydration, however, can vary from 67% to 85% and can vary by disease state. Other assumptions used to derive body composition estimates are considered proprietary by DXA manufacturers. The use of DXA for bone mineral density assessment in patients diagnosed with or at risk of osteoporosis vertebral fracture assessment with densitometry by DXA are not addressed in this policy.

Regulatory Status
Body composition software for several bone densitometer systems has been approved by the U.S. Food and Drug Administration (FDA) through the premarket approval process. They include the Lunar iDXA systems (GE Healthcare), Hologic DXA systems (Hologic), Mindways Software, Inc. systems (Mindways Software, Inc.), and Norland DXA systems (Swissray).

For individuals who have a clinical condition associated with abnormal body composition who receive dual-energy x-ray absorptiometry (DXA) body composition studies, the evidence includes systematic reviews and several cross-sectional studies comparing DXA with other techniques. Relevant outcomes are symptoms and change in disease status. The available studies were primarily conducted in research settings and often used DXA body composition studies as a reference standard. Systematic reviews with meta-analyses exploring the clinical validity of DXA measurements against reference methods for the quantification of fat mass indicate strong overall agreement between these modalities, but raise concerns regarding precision and reliability in some populations, particularly those without existing clinical conditions for which risk of adverse outcomes is influenced by abnormal visceral adiposity. More importantly, no studies were identified in which DXA body composition measurements were actively used in patient management. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a clinical condition managed by monitoring changes in body composition over time who receive serial DXA body composition studies, the evidence includes several prospective studies monitoring patients over time. Relevant outcomes are symptoms and change in disease status. The studies used DXA as a tool to measure body composition and were not designed to assess the accuracy of DXA. None of the studies used DXA findings to make patient management decisions or addressed how serial body composition assessment might improve health outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

CODING

The following CPT code(s) are not covered for Medicare Advantage Plans and not medically necessary for Commercial Products:

76499 Unlisted diagnostic radiographic procedure

RELATED POLICIES

Bone Mineral Density Studies
Dynamic Spinal Visualization and Vertebral Motion Analysis
Vertebral Fracture Assessment

PUBLISHED

Provider Update, January/December 2026
Provider Update, January 2025

REFERENCES

1. Borga M, West J, Bell JD, et al. Advanced body composition assessment: from body mass index to body composition profiling. J Investig Med. Jun 2018; 66(5): 1-9. PMID 29581385
2. Barone M, Losurdo G, Iannone A, et al. Assessment of body composition: Intrinsic methodological limitations and statistical pitfalls. Nutrition. Oct 2022; 102: 111736. PMID 35810580
3. Murphy J, Bacon SL, Morais JA, et al. Intra-Abdominal Adipose Tissue Quantification by Alternative Versus Reference Methods: A Systematic Review and Meta-Analysis. Obesity (Silver Spring). Jul 2019;27(7): 1115-1122. PMID 31131996
4. Sheean P, Gonzalez MC, Prado CM, et al. American Society for Parenteral and Enteral Nutrition Clinical Guidelines: The Validity of Body Composition Assessment in Clinical Populations. JPEN J Parenter Enteral Nutr. Jan 2020; 44(1): 12-43. PMID 31216070
5. Calella P, Valerio G, Brodlie M, et al. Tools and Methods Used for the Assessment of Body Composition in Patients With Cystic Fibrosis: A Systematic Review. Nutr Clin Pract. Oct 2019; 34(5): 701-714. PMID30729571
6. Calella P, Valerio G, Brodlie M, et al. Cystic fibrosis, body composition, and health outcomes: a systematic review. Nutrition. Nov 2018; 55-56: 131-139. PMID 29981489
7. Bundred J, Kamarajah SK, Roberts KJ. Body composition assessment and sarcopenia in patients with pancreatic cancer: a systematic review and meta-analysis. HPB (Oxford). Dec 2019; 21(12): 1603-1612.PMID 31266698

8. Alves FD, Souza GC, Biolo A, et al. Comparison of two bioelectrical impedance devices and dual-energy X-ray absorptiometry to evaluate body composition in heart failure. J Hum Nutr Diet. Dec 2014;27(6): 632-8. PMID 24684316
9. Ziai S, Coriati A, Chabot K, et al. Agreement of bioelectric impedance analysis and dual-energy X-ray absorptiometry for body composition evaluation in adults with cystic fibrosis. J Cyst Fibros. Sep 2014;13(5): 585-8. PMID 24522087
10. Elkan AC, Engvall IL, Tengstrand B, et al. Malnutrition in women with rheumatoid arthritis is not revealed by clinical anthropometrical measurements or nutritional evaluation tools. Eur J Clin Nutr. Oct 2008;62(10): 1239-47. PMID 17637600
11. Jensky-Squires NE, Dieli-Conwright CM, Rossuello A, et al. Validity and reliability of body composition analysers in children and adults. Br J Nutr. Oct 2008; 100(4): 859-65. PMID 18346304
12. Kullberg J, Brandberg J, Angelhed JE, et al. Whole-body adipose tissue analysis: comparison of MRI,CT and dual energy X-ray absorptiometry. Br J Radiol. Feb 2009; 82(974): 123-30. PMID 19168691
13. Liem ET, De Lucia Rolfe E, L'Abée C, et al. Measuring abdominal adiposity in 6 to 7-year-old children. Eur J Clin Nutr. Jul 2009; 63(7): 835-41. PMID 19127281
14. Bedogni G, Agosti F, De Col A, et al. Comparison of dual-energy X-ray absorptiometry, air displacement plethysmography and bioelectrical impedance analysis for the assessment of body composition in morbidly obese women. Eur J Clin Nutr. Nov 2013; 67(11): 1129-32. PMID 24022260
15. Monteiro PA, Antunes Bde M, Silveira LS, et al. Body composition variables as predictors of NAFLD by ultrasound in obese children and adolescents. BMC Pediatr. Jan 29 2014; 14: 25. PMID 24476029
16. Tompuri TT, Lakka TA, Hakulinen M, et al. Assessment of body composition by dual-energy X-ray absorptiometry, bioimpedance analysis and anthropometrics in children: the Physical Activity and Nutrition in Children study. Clin Physiol Funct Imaging. Jan 2015; 35(1): 21-33. PMID 24325400
17. Alves Junior CAS, de Lima LRA, de Souza MC, et al. Anthropometric measures associated with fat mass estimation in children and adolescents with HIV. Appl Physiol Nutr Metab. May 2019; 44(5): 493-498. PMID 30286302
18. Barr RD, Inglis D, Athale U, et al. The Influence of Body Composition on Bone Health in Long-term Survivors of Acute Lymphoblastic Leukemia in Childhood and Adolescence: Analyses by Dual-energy Radiograph Absorptiometry and Peripheral Quantitative Computed Tomography. J Pediatr Hematol Oncol. Nov 01 2022; 44(8): 423-431. PMID 35482464
19. Chang CC, Chen YK, Chiu HC, et al. Assessment of Sarcopenia and Obesity in Patients with Myasthenia Gravis Using Dual-Energy X-ray Absorptiometry: A Cross-Sectional Study. J Pers Med. Nov03 2021; 11(11). PMID 34834491
20. Smoot BJ, Mastick J, Shepherd J, et al. Use of Dual-Energy X-Ray Absorptiometry to Assess Soft Tissue Composition in Breast Cancer Survivors With and Without Lymphedema. Lymphat Res Biol. Aug 2022; 20(4): 391-397. PMID 34793255
21. Dandache C, Confavreux CB, Gavoille A, et al. Peripheral but not axial muscle mass is associated with early mortality in bone metastatic lung cancer patients at diagnosis. Joint Bone Spine. Sep 2023; 90(5):105613. PMID 37442335
22. Wang LH, Leung DG, Wagner KR, et al. Lean tissue mass measurements by dual-energy X-ray absorptiometry and associations with strength and functional outcome measures in facioscapulohumeral muscular dystrophy. Neuromuscul Disord. Sep 2023; 33(9): 63-68. PMID37400350
23. Dawra S, Gupta P, Yadav N, et al. Association between the Distribution of Adipose Tissue and Outcomes in Acute Pancreatitis: A Comparison of Methods of Fat Estimation. Indian J Radiol Imaging. Jan 2023; 33(1): 12-18. PMID 36855725
24. Woolcott OO, Bergman RN. Defining cutoffs to diagnose obesity using the relative fat mass (RFM): Association with mortality in NHANES 1999-2014. Int J Obes (Lond). Jun 2020; 44(6): 1301-1310. PMID31911664
25. Staunstrup LM, Nielsen HB, Pedersen BK, et al. Cancer risk in relation to body fat distribution, evaluated by DXA-scans, in postmenopausal women - the Prospective Epidemiological Risk Factor(PERF) study. Sci Rep. Mar 29 2019; 9(1): 5379. PMID 30926844
26. Reina D, Gómez-Vaquero C, Díaz-Torné C, et al. Assessment of nutritional status by dual X-Ray absorptiometry in women with rheumatoid arthritis: A case-control study. Medicine (Baltimore). Feb2019; 98(6): e14361. PMID 30732168

27. Sinclair M, Hoermann R, Peterson A, et al. Use of Dual X-ray Absorptiometry in men with advanced cirrhosis to predict sarcopenia-associated mortality risk. Liver Int. Jun 2019; 39(6): 1089-1097. PMID30746903
28. Lindqvist C, Brismar TB, Majeed A, et al. Assessment of muscle mass depletion in chronic liver disease: Dual-energy x-ray absorptiometry compared with computed tomography. Nutrition. May 2019; 61: 93-98. PMID 30703575
29. Dordevic AL, Bonham M, Ghasem-Zadeh A, et al. Reliability of Compartmental Body Composition Measures in Weight-Stable Adults Using GE iDXA: Implications for Research and Practice. Nutrients. Oct 12 2018; 10(10). PMID 30321991
30. Bazzocchi A, Ponti F, Cariani S, et al. Visceral fat and body composition changes in a female population after RYGBP: a two-year follow-up by DXA. Obes Surg. Mar 2015; 25(3): 443-51. PMID 25218013
31. Franzoni E, Ciccarese F, Di Pietro E, et al. Follow-up of bone mineral density and body composition in adolescents with restrictive anorexia nervosa: role of dual-energy X-ray absorptiometry. Eur J Clin Nutr. Feb 2014; 68(2): 247-52. PMID 24346474
32. Iyengar NM, Arthur R, Manson JE, et al. Association of Body Fat and Risk of Breast Cancer in Postmenopausal Women With Normal Body Mass Index: A Secondary Analysis of a Randomized Clinical Trial and Observational Study. JAMA Oncol. Feb 01 2019; 5(2): 155-163. PMID 30520976
33. Ashby-Thompson M, Heshka S, Rizkalla B, et al. Validity of dual-energy x-ray absorptiometry for estimation of visceral adipose tissue and visceral adipose tissue change after surgery-induced weight loss in women with severe obesity. Obesity (Silver Spring). May 2022; 30(5): 1057-1065. PMID35384351
34. Sherlock SP, Palmer J, Wagner KR, et al. Dual-energy X-ray absorptiometry measures of lean body mass as a biomarker for progression in boys with Duchenne muscular dystrophy. Sci Rep. Nov 05 2022;12(1): 18762. PMID 36335191
35. Nguyen AL, Herath M, Burns M, et al. The value of whole-body dual-energy x-ray absorptiometry in assessing body composition in patients with inflammatory bowel disease: a prospective study. Eur J Gastroenterol Hepatol. Jan 01 2024; 36(1): 52-61. PMID 37942750
36. Arthur RS, Xue X, Kamensky V, et al. The association between DXA-derived body fat measures and breast cancer risk among postmenopausal women in the Women's Health Initiative. Cancer Med. Feb2020; 9(4): 1581-1599. PMID 31875358
37. Garvey WT, Mechanick JI, Brett EM, et al. AMERICAN ASSOCIATION OF CLINICALENDOCRINOLOGISTS AND AMERICAN COLLEGE OF ENDOCRINOLOGY COMPREHENSIVECLINICAL PRACTICE GUIDELINES FOR MEDICAL CARE OF PATIENTS WITH OBESITY. Endocr Pract. Jul 2016; 22 Suppl 3: 1-203. PMID 27219496
38. Nadolsky K, Garvey WT, Agarwal M, et al. American Association of Clinical Endocrinology Consensus Statement: Algorithm for the Evaluation and Treatment of Adults with Obesity/Adiposity-Based Chronic Disease - 2025 Update. Endocr Pract. Nov 2025; 31(11): 1351-1394. PMID 40956256
39. American College of Radiology. ACR-SPR-SSR Practice Parameter for the Performance of Dual-Energy X-ray Absorptiometry (DXA). 2018; https://www.acr.org/-/media/ACR/Files/Practice-Parameters/DXA.pdf. Accessed July 22, 2026.
40. International Society for Clinical Densitometry. 2019 ISCD Official Positions - Adult. 2019; https://iscd.org/wp-content/uploads/2021/09/2019-Official-Positions-Adult-1.pdf. Accessed July 22, 2026.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.