

## ACITRETIN (S)

### Products Affected

- Acitretin

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	Severely impaired liver or kidney function. Chronic abnormally elevated blood lipid values. Concomitant use of methotrexate or tetracyclines. Pregnancy. Females of child-bearing potential who intend to become pregnant during therapy or at any time for at least 3 years after discontinuing therapy. Females of child-bearing potential who will not use reliable contraception while undergoing treatment and for at least 3 years following discontinuation. Females of child-bearing potential who drink alcohol during treatment or for two months after cessation of therapy.
Required Medical Information	Diagnosis of severe psoriasis
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

# ACTEMRA (BCBS RI)

---

## Products Affected

- Actemra INJ 162MG/0.9ML, 200MG/10ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis).
<b>Required Medical Information</b>	Diagnosis of one of the following: A) moderate to severe rheumatoid arthritis and patient had inadequate response to, intolerance to, or contraindication to one or more non-biologic disease-modifying anti-rheumatic drugs for at least 3 consecutive months, or B) systemic juvenile idiopathic arthritis and patient had inadequate response or intolerance to at least one oral systemic agent (i.e. NSAID, corticosteroid) C) Polyarticular Juvenile Idiopathic Arthritis AND Patient has had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs for at least 3 consecutive months
<b>Age Restrictions</b>	RA - 18 years of age or older. sJIA and pJIA - 2 years of age or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

# ACTIQ (BCBS RI)

## Products Affected

- Fentanyl Citrate Oral Transmucosal

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Management of acute or post-operative pain, including headache/migraine, dental pain, or use in the emergency room. Opioid non-tolerant patients.
<b>Required Medical Information</b>	Patient meets the following: A) Diagnosis of cancer and use is for breakthrough cancer pain, B) patient is opioid tolerant and taking at least 60 mg morphine/day, at least 25 mcg transdermal fentanyl/hour, at least 30 mg of oxycodone daily, at least 8 mg oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer, C) at least one other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, D) prescriber is registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

## ADAGEN (S)

---

### Products Affected

- Adagen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Severe thrombocytopenia
<b>Required Medical Information</b>	Diagnosis of adenosine deaminase (ADA) deficiency in a patient with severe combined immunodeficiency disease (SCID) AND patient is not a suitable candidate for, or who has failed, bone marrow transplantation.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ADCIRCA (S)

---

### Products Affected

- Adcirca

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Receiving nitrate therapy (includes intermittent use)
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) Functional Class II or III that was confirmed by right heart catheterization
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	N/A

## ADEMPAS (S)

---

### Products Affected

- Adempas

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form. Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline). Pregnancy.
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO group I with New York Heart Association Functional Class II or III AND diagnosis was confirmed by right heart catheterization OR Patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) AND patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable AND female patients are enrolled in the ADEMPAS REMS program
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	For renewal, medication was effective (i.e. improved 6 minute walk distance, oxygen saturation, etc.)

# AFINITOR (S)

## Products Affected

- Afinitor

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced metastatic renal cell carcinoma and patient has failed therapy (disease progressed) with Sutent or Nexavar OR Diagnosis of progressive pancreatic neuroendocrine tumors (pNET) that are unresectable, locally advanced or metastatic OR Diagnosis of renal angiomyolipoma with tuberous sclerosis complex (TSC) and patient does not require immediate surgery OR Diagnosis of advanced hormone receptor-positive, HER2-negative breast cancer and patient is a postmenopausal woman and patient has failed treatment with Femara or Arimidex and the medication will be used in combination with Aromasin OR Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with TSC that requires therapeutic intervention but is not a candidate for curative surgical resection.
<b>Age Restrictions</b>	18 years of age or older for RCC, pNET, and renal angiomyolipoma with TSC. 1 year of age or older for SEGA
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## AFINITOR DISPERZ (S)

---

### Products Affected

- Afinitor Disperz

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex (TSC) that requires therapeutic intervention but is not a candidate for curative surgical resection.
<b>Age Restrictions</b>	1 year of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## ALDURAZYME (S)

---

### Products Affected

- Aldurazyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Hurler or Hurler-Scheie form of Mucopolysaccharidosis I (MPS I) or Diagnosis of Scheie form of MPS I with moderate to severe symptoms.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## AMITIZA (S)

---

### Products Affected

- Amitiza

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Mechanical gastrointestinal obstruction.
<b>Required Medical Information</b>	Diagnosis of irritable bowel syndrome-constipation for at least 12 non-consecutive weeks and patient has tried and failed increasing fluid and fiber intake and patient has tried and failed or has an intolerance to osmotic laxatives, stimulant laxatives or probiotics and patient is female OR Diagnosis of chronic idiopathic constipation for at least 3 months and patient has tried and failed increasing fluid and fiber intake and patient has tried and failed or has an intolerance to osmotic laxatives, stimulant laxatives or stool softeners OR Diagnosis of chronic opioid-induced constipation due to non-cancer pain and patient has tried and failed a stool softener and a stimulant laxative AND Patient is not being treated with methadone
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 4 months. Renewal - 12 months
<b>Other Criteria</b>	For renewal, the patient has experienced an increase in the number of bowel movements.

## AMPYRA (BCBS RI)

---

### Products Affected

- Ampyra

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	History of seizure. Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute). Patient currently using any other forms of 4-aminopyridine.
<b>Required Medical Information</b>	Diagnosis of multiple sclerosis AND patient is ambulatory (able to walk at least 25 feet) AND patient has walking impairment
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 1 month. Renewal - 12 months
<b>Other Criteria</b>	For renewal, walking speed has improved from baseline.

## ANADROL-50 (S)

---

### Products Affected

- Anadrol-50

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Breast or prostate cancer in men. Breast cancer in women with hypercalcemia. Pregnancy. Nephrosis or nephrotic phase of nephritis. Severe hepatic dysfunction.
<b>Required Medical Information</b>	Diagnosis of anemia caused by deficient red cell production AND patient has tried and had an inadequate response or intolerance to standard therapies for anemia such as: erythropoietin stimulating agents, immunosuppressants, blood transfusions, etc. AND treatment will not replace other supportive measures (e.g., transfusion, iron supplementation, immunosuppressive therapy, corticosteroids, erythropoiesis-stimulating agents)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient experienced an objective improvement in anemia (e.g., increased hemoglobin, increased reticulocyte count, reduction/elimination for need of blood transfusions)

# ARANESP (S)

## Products Affected

- Aranesp Albumin Free INJ  
100MCG/0.5ML, 100MCG/ML,  
150MCG/0.3ML, 200MCG/0.4ML,  
200MCG/ML, 25MCG/0.42ML,  
25MCG/ML, 300MCG/0.6ML,  
300MCG/ML, 40MCG/0.4ML,  
40MCG/ML, 500MCG/ML,  
60MCG/0.3ML, 60MCG/ML

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Anemia due to myelodysplastic syndrome.
<b>Exclusion Criteria</b>	uncontrolled hypertension. Pure red cell aplasia that begins after ESA treatment.
<b>Required Medical Information</b>	Pre-treatment hemoglobin level less than 10 g/dL AND Patient has adequate iron stores prior to initiation of therapy defined as ferritin more than 100 mcg/L or serum transferrin saturation greater than 20% AND other causes of anemia such as iron deficiency, folate deficiency or B12 deficiency, hemolysis, gastrointestinal bleeding, other active or occult bleeding, or underlying hematologic disease (such as sickle cell anemia, thalassemia, and porphyria) have been ruled out AND Diagnosis of one of the following: A) Anemia due to chronic kidney disease (CKD) with or without hemodialysis, OR B) Anemia in patients with non-myeloid malignancies where anemia is due to the effect of concomitant myelosuppressive chemotherapy and two additional months of chemotherapy is anticipated C) anemia due to myelodysplastic syndrome (MDS) and endogenous serum erythropoietin level is 500 mUnits/mL or less
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CKD - prescribed by a nephrologist or hematologist. Non-myeloid malignancies, MDS - prescribed by an oncologist/hematologist.
<b>Coverage Duration</b>	Initial: 4 months. Renewal: CKD-12 months, Non-myeloid malignancies - 4 months

<b>Other Criteria</b>	For renewal of CKD, for dialysis patients: Hb less than 11 g/dL or physician will decrease or interrupt dose. For non-dialysis patients: Hb less than 10 g/dL or physician will decrease or interrupt dose. For renewal of non-myeloid malignancies: Concurrent myelosuppressive chemotherapy and Hb is 12g/dL or less and there is measurable response after eight weeks (defined as an increase in Hb 1 g/dL or more or a reduction in red blood cell transfusion requirements). For renewal of MDS, Hb is 12g/dL or less and there is measurable response after eight weeks (defined as an increase in Hb 1.5 g/dL or more or a reduction in red blood cell transfusion requirements) OR Patient will have a concomitant trial of G-CSF AND, if patient had concomitant trial of granulocyte colony-stimulating factor (G-CSF), patient had a measurable response as defined above after 8 weeks of G-CSF therapy. Excluded for ESRD patients on dialysis.
-----------------------	---

## ARCALYST (S)

---

### Products Affected

- Arcalyst

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cryopyrin-associated period syndromes (CAPS), including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS).
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist, or other medical specialist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient experienced disease stability or improvement.

## AUBAGIO (BCBS RI)

---

### Products Affected

- Aubagio

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. First clinical episode with MRI features consistent with multiple sclerosis.
<b>Exclusion Criteria</b>	Severe hepatic impairment. Current treatment with leflunomide. Patients who are pregnant or women of childbearing potential not using reliable contraception.
<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis AND patient has had an inadequate response, intolerance or contraindication to Betaseron (interferon beta 1b), Avonex (interferon beta 1A) or Copaxone (glatiramer)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced no or a decrease in the number of multiple sclerosis flare-ups.



## AVASTIN (S)

### Products Affected

- Avastin INJ 100MG/4ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Gastrointestinal perforation. Wound dehiscence. Serious hemorrhage or recent hemoptysis.
<b>Required Medical Information</b>	Diagnosis of one of the following: A) First-line or second-line treatment of metastatic carcinoma of the colon or rectum in combination with IV 5-fluorouracil-based chemotherapy B) Second-line treatment of metastatic colorectal cancer in combination with fluoropyrimidine-irinotecan- or fluoropyrimidine-oxaliplatin-based chemotherapy after progression on a first-line Avastin-containing regimen. C) First-line treatment of unresectable, locally advanced, recurrent or metastatic non-squamous non-small cell lung cancer in combination with carboplatin and paclitaxel, D) glioblastoma with progressive disease following prior therapy and the medication will be used as a single agent, E) metastatic renal cell carcinoma in combination with interferon alfa.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# AVONEX (S)

---

## Products Affected

- Avonex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All-FDA approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)

# BARACLUDE (BCBS RI)

## Products Affected

- Baraclude

- Entecavir

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis B AND patient is HBsAg-positive for at least 6 months AND For HBeAg-positive patients, serum HBV DNA greater than 20,000 IU/mL (105 copies per mL) and for HBeAg-negative patients, serum HBV DNA greater than 2,000 IU/mL (104 copies/mL) AND Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) at least 2 times the upper limit of normal or histologically active disease (i.e. necroinflammation on biopsy) AND patient is receiving anti-retroviral therapy if the patient has HIV co-infection
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient must be HBeAg negative and have not had HBsAg clearance OR HBeAg positive and have detectable HBV DNA and have not been anti-Hbe for at least 6 months.

## BELEODAQ (S)

---

### Products Affected

- Beleodaq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of peripheral T-cell lymphoma AND patient has tried and had an inadequate response, intolerance or contraindication to at least one prior therapy (e.g., conventional chemotherapy)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## BENZODIAZEPINES (S)

### Products Affected

- Alprazolam ORAL TABS
- Alprazolam Er
- Alprazolam Intensol
- Alprazolam Odt
- Alprazolam Xr
- Chlordiazepoxide Hcl
- Clonazepam ORAL TABS
- Clonazepam Odt
- Clorazepate Dipotassium
- Diazepam CONC
- Diazepam ORAL TABS
- Diazepam SOLN
- Diazepam Intensol
- Estazolam
- Flurazepam Hcl
- Lorazepam CONC
- Lorazepam ORAL TABS
- Lorazepam Intensol
- Onfi
- Oxazepam
- Temazepam
- Triazolam

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	For alprazolam only: concomitant use with ketoconazole or itraconazole.
<b>Required Medical Information</b>	Verify the medication is being used for an FDA-approved diagnosis or compendial supported indication.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

## BETHKIS (BCBS RI)

---

### Products Affected

- Bethkis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs AND has the patient tried and failed generic tobramycin inhalation solution
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations). For inhalation solution only (does not apply to TOBI Podhaler): Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## BONIVA IV (BCBS RI)

### Products Affected

- Boniva INJ

- Ibandronate Sodium INJ

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient is a postmenopausal female with osteoporosis AND Patient has a documented trial and failure of an oral bisphosphonate, where failure is defined as new fractures in compliant patient on therapy for at least 6 months, failure to produce a clinically significant change in biochemical markers of bone turnover, or a significant loss of bone mineral density on follow-up scans after 12-24 months of therapy OR documented contraindication or intolerance to oral bisphosphonate therapy or is unable to comply with appropriate administration recommendations for oral bisphosphonate therapy.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient had an objective response to therapy. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

## BOSULIF (BCBS RI)

---

### Products Affected

- Bosulif

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Philadelphia chromosome-positive chronic myelogenous leukemia AND resistance, relapse, inadequate response to prior therapy with a tyrosine kinase inhibitor (TKI) AND if patient had mutation testing, patient does not have T315I or V299L mutation or intolerant to prior TKI therapy
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



# BOTOX (BCBS RI)

---

## Products Affected

- Botox

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Achalasia. Lower limb spasticity due to stroke. Cervicogenic headache. Dysphagia. Salivation due to Parkinson's disease. Tourette's syndrome. Hemifacial spasm. Trigeminal neuralgia. Oromandibular dystonia. Tardive dyskinesia. Temporomandibular joint dysfunction.
<b>Exclusion Criteria</b>	Infection at the proposed injection site. Cosmetic use (e.g., wrinkles). For OAB and Urinary incontinence, patient has urinary tract infection or patient is not routinely performing clean intermittent self-catheterization (CIC) or is not willing/able to perform CIC if post-void residual urine volume is more than 200 mL.

<b>Required Medical Information</b>	Diagnosis of: A) strabismus, B) blepharospasm associated with dystonia, C) Urinary incontinence associated with neurologic condition and inadequate response to at least one antimuscarinic agent, unless contraindicated or intolerant to antimuscarinics (e.g., narrow angle glaucoma), D) Chronic migraine and Botox will be used as prophylaxis and experiences headaches on 15 or more days per month lasting four hours or longer and inadequate response with at least two first-line therapies from two different therapeutic classes (i.e. antiepileptics, beta-blockers, triptans, and tricyclic antidepressants), E) Cervical dystonia (including spasmodic torticollis) F) Overactive bladder and has symptoms (e.g., urge urinary incontinence, urgency, and frequency) and inadequate response to at least one antimuscarinic agent, unless contraindicated or intolerant to anti-muscarinics, G) Axillary hyperhidrosis refractory to topical aluminum chloride and condition significantly interferes with patient's daily activities, H) Upper limb spasticity and Botox will be used to improve muscle tone at wrist, elbow or finger flexors, I) lower limb spasticity due to stroke, J) achalasia and patient is not suitable for surgery, K) Cervicogenic headache and inadequate response to at least two therapies from two different therapeutic classes (e.g., NSAIDs, muscle relaxants, tricyclic antidepressants), L) Dysphagia/esophageal dysmotility, M) excessive salivation due to advanced Parkinson's disease, N) Gilles de la Tourette's syndrome and inadequate response to conventional agents (e.g., fluphenazine, haloperidol, pimozide), O) hemifacial spasm, P) refractory idiopathic trigeminal neuralgia and inadequate response to other therapies (e.g., carbamazepine, lamotrigine, or baclofen), Q) isolated oromandibular dystonia, R) tardive dyskinesia and patient failed other therapies (e.g., clozapine, risperidone, quetiapine) S) temporomandibular joint dysfunction
<b>Age Restrictions</b>	12 years of age or older - strabismus or blepharospasm, 18 years of age or older - axillary hyperhidrosis
<b>Prescriber Restrictions</b>	Urinary incontinence - prescribed by or in consultation with an urologist
<b>Coverage Duration</b>	12 months

<b>Other Criteria</b>	<p>For renewal of cervical dystonia, patient experienced improvement in at least one of the following: pain, severity of abnormal head position, or effects on patient's daily activities. For renewal of blepharospasm, patient experienced an improvement in symptoms (e.g., ability to open eyes, improvement in blinking/spasms of the eye). For renewal of hemifacial spasm, excessive salivation due to Parkinson's disease, oromandibular dystonia, tardive dyskinesia, and hyperhidrosis, the patient experienced an improvement in symptoms. For renewal of strabismus, patient has improvement in vision or eye alignment. For renewal of migraine, patient experienced a decrease in the frequency of headaches. For renewal of OAB and urinary incontinence due to a neurologic condition, patient experienced an improvement in symptoms (e.g., urge urinary incontinence, urgency, and frequency). For renewal of upper limb spasticity, patient experienced an improvement in flexor muscle tone. For renewal of achalasia and dysphagia/esophageal dysmotility, improvement in dysphagia, chest pain and regurgitation. For renewal of cervicogenic headache, improvement in neck range of motion or pain. For renewal of Gilles de la Tourette's syndrome, improvement in blinking, blepharospasm, or dystonic tics. For renewal of trigeminal neuralgia, improvement in pain. For renewal of Temporomandibular joint dysfunction, improvement in pain, functioning, or tenderness. For renewal of lower limb spasticity, improvement in lower limb muscle tone, gait pattern, or walking distance.</p>
-----------------------	--

## BUTRANS (S)

---

### Products Affected

- Butrans

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Significant respiratory depression or severe bronchial asthma. Known or suspected paralytic ileus.
<b>Required Medical Information</b>	Patient is not in hospice care AND Patient has a diagnosis of severe pain requiring continuous, around-the-clock opioid analgesic for an extended period of time (at least 2 weeks) AND patient tried and failed, is unable to tolerate, two generic extended-release opioid product and/or opioid combination product, unless the patient has documented swallowing difficulties
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

## CAPRELSA (BCBS RI)

---

### Products Affected

- Caprelsa

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Congenital long QT syndrome, Prenancy
<b>Required Medical Information</b>	Diagnosis of symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider and patient must be certified in the Caprelsa REMS program
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# CAYSTON (S)

---

## Products Affected

- Cayston

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs
<b>Age Restrictions</b>	7 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations)

## CELEBREX (S)

### Products Affected

- Celebrex

- Celecoxib ORAL CAPS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Hypersensitivity to celecoxib or sulfonamide drugs. History of asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Treatment of peri-operative pain in the setting of coronary artery bypass graft surgery. Active gastrointestinal bleeding
<b>Required Medical Information</b>	Patient has failed previous treatment or has an intolerance to at least one of the following: NSAIDs or salicylate OR patient is currently receiving treatment with any of the following: anticoagulants/antiplatelet drugs (i.e. warfarin, heparin, LMWH, Pradaxa, Plavix, etc.), antiulcer agents (i.e. proton pump inhibitors, histamine-2-receptor antagonists, or misoprostol), chronic use of oral corticosteroids, or methotrexate OR patient has a history of peptic ulcer disease or history of gastrointestinal bleed
<b>Age Restrictions</b>	PA applies to patients younger than 65 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## CERDELGA (S)

---

### Products Affected

- Cerdelga

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Extensive or intermediate metabolizers taking a strong or moderate CYP2D6 inhibitor concomitantly with a strong or moderate CYP3A inhibitor. Intermediate or poor metabolizers taking a strong CYP3A inhibitor.
<b>Required Medical Information</b>	The patient has a diagnosis of Gaucher disease type 1 AND patient is an extensive metabolizer, intermediate metabolizer, or poor metabolizer of CYP2D6 as detected by an FDA-cleared test
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient's condition has not progressed as defined by ALL of the following: hemoglobin level decreased by more than 1.5 g/dL from baseline, platelet count decreased more than 25% from baseline, spleen volume increased more than 25% from baseline, and liver volume increased more than 20% from baseline



## CEREZYME (BCBS RI)

---

### Products Affected

- Cerezyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of type 1 Gaucher disease. Gaucher disease confirmed by enzymatic testing AND 1 or more complication such as anemia, thrombocytopenia, bone disease, and hepatomegaly, and splenomegaly
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## CESAMET (S)

---

### Products Affected

- Cesamet

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chemotherapy-induced nausea and vomiting AND patient has tried and failed at least one conventional antiemetic treatments (e.g., aprepitant/fosaprepitant, t-hydroxytryptamine-3 serotonin receptor antagonists such as ondansetron) AND patient has tried and failed dronabinol
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	Part B if related to cancer treatment and is a full replacement for IV antiemetic within 48 hrs of cancer treatment. Part D if related to cancer treatment after the 48-hour period, or for any other medically accepted diagnosis.

## CIALIS 2.5 MG, 5 MG (S)

---

### Products Affected

- Cialis ORAL TABS 2.5MG, 5MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Concurrent use of nitrates. Diagnosis of erectile dysfunction without signs and symptoms of BPH.
<b>Required Medical Information</b>	Diagnosis of benign prostatic hyperplasia (BPH) and patient has experienced intolerance to or treatment failure with an alpha-blocker (e.g., doxazosin, prazosin, tamsulosin) or a 5-alpha reductase inhibitor (e.g., dutasteride, finasteride)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## CIMZIA (S)

### Products Affected

- Cimzia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis)
<b>Required Medical Information</b>	Diagnosis of one of the following: A) moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs for at least 3 consecutive months B) moderate to severe Crohn's disease and patient had an inadequate response to, is intolerant to, or is contraindicated to conventional therapy with one or more of the following: corticosteroids (i.e. prednisone, methylprednisolone) or non-biologic DMARDs (i.e. azathioprine, methotrexate, mercaptopurine, etc.) C) psoriatic arthritis and patient had an inadequate response, intolerance to, or contraindication to methotrexate D) ankylosing spondylitis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-steroidal anti-inflammatory drugs
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial 16 weeks (CD), 12 weeks (others). Renewal 12 months.
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. For renewal, patient has obtained a clinical response to therapy (e.g., for CD, symptomatic remission. For RA, improvement in tender/swollen joint count, improvement in ACR scoring. For PsA, improvement in number of swollen/tender joints, pain, stiffness. For AS, improvement in AS symptoms, such as stiffness and back pain) or patient's condition has stabilized.

## COMETRIQ (S)

---

### Products Affected

- Cometriq

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Gastrointestinal perforation. Fistula. Severe hemorrhage.
<b>Required Medical Information</b>	Diagnosis of progressive, metastatic medullary thyroid cancer
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# COPAXONE (S)

---

## Products Affected

- Copaxone

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsing-remitting multiple sclerosis OR diagnosis of first clinical episode with MRI features consistent with multiple sclerosis
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has no or slowed disease progression

## CYSTARAN (S)

---

### Products Affected

- Cystaran

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of cystinosis AND Patient has corneal cystine crystal accumulation
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## DALIRESP (S)

---

### Products Affected

- Daliresp

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Moderate to severe liver impairment (Child-Pugh B or C)
<b>Required Medical Information</b>	Diagnosis of severe chronic obstructive pulmonary disease (COPD) (defined as FEV1 less than or equal to 50% of predicted and FEV1/forced vital capacity [FVC] less than 0.7) associated with chronic bronchitis AND history of COPD exacerbations which requires the use of systemic corticosteroids, antibiotics, or hospital admission AND Medication will be used with a long-acting inhaled bronchodilator (i.e. long-acting anticholinergic, or long-acting beta agonist in combination with inhaled corticosteroid)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## DICLOFENAC GEL (S)

---

### Products Affected

- Voltaren GEL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	Known hypersensitivity to diclofenac. Previously experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Use for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery.
<b>Required Medical Information</b>	Patient has a diagnosis of osteoarthritis of the knees or hands AND patient had experienced treatment failure with at least 2 prescription strength oral NSAIDs or patient has a documented swallowing disorder OR has a history of peptic ulcer disease/gastrointestinal bleeding OR patient is more than 65 years of age with one additional risk factor for gastrointestinal adverse event (e.g., use of anticoagulants or chronic corticosteroids)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# DIFFERIN (S)

## Products Affected

- Adapalene

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Acne rosacea. Actinic keratoses (solar keratoses). Flat warts (verruca plana/juveniles). Hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans [Flegel's disease], palmoplantar hyperatosis, keratosis follicularis [Darier's disease]). Ichthyosis (i.e. congenital, lamellar, vulgaris, or X-linked)
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of mild to moderate acne vulgaris (including comedonal, cystic, and nodular) OR Diagnosis of acne rosacea and patient has experienced treatment failure of or intolerance to sulfacetamide, metronidazole, or azelaic acid OR patient is receiving treatment for actinic keratosis (or solar keratoses) with multiple lesions on the face OR Diagnosis of flat warts (verruca plana/jeveniles) OR Diagnosis of hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans [Flegel's disease], palmoplantar hyperatosis, keratosis follicularis [Darier's disease] OR Diagnosis of ichthyosis (i.e. congenital, lamellar, vulgaris, or X-linked)
<b>Age Restrictions</b>	PA applies to patients older than 26 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, medication has been effective in treating the patient's condition.

## EGRIFTA (S)

---

### Products Affected

- Egrifta INJ 2MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnant. Active malignancy (newly diagnosed or recurrent).
<b>Required Medical Information</b>	Diagnosis of excess abdominal fat secondary to HIV infection with lipodystrophy AND Patient has been receiving antiretroviral therapy AND waist circumference greater than 37.4 inches (men) or greater than 37 inches (women) AND waist-to-hip ratio greater than 0.94 (men) or greater than 0.88 (women)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	For renewal, documentation that the patient has experienced a reduction from baseline in visceral adipose tissue as measured by waist circumference.

## ELELYSO (BCBS RI)

---

### Products Affected

- Eleyso

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of type 1 Gaucher disease confirmed by enzymatic testing AND 1 or more complication such as anemia, thrombocytopenia, bone disease, hepatomegaly, and splenomegaly AND will not be used in combination with Zavesca
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ELIGARD (S)

---

### Products Affected

- Eligard

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced or metastatic prostate cancer
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## EMSAM (S)

### Products Affected

- Emsam

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pheochromocytoma. Patient is taking or will take any of the following: SSRIs, SNRIs, tricyclic antidepressants (TCAs), bupropion, buspirone, meperidine, tramadol, methadone, pentazocine, dextromethorphan, St. John's wort, mirtazapine, cyclobenzaprine, oral selegiline, other MAOIs, oxcarbazepine, carbamazepine, and/or sympathomimetic amines
<b>Required Medical Information</b>	Diagnosis of major depressive disorder AND Patient had adequate trial with at least 2 generic oral antidepressants from differing classes (at least one should be from the following list: selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, mirtazapine, or bupropion unless contraindicated), unless unable to take any oral medication AND Patient had an adequate washout period (for patients previously on agents requiring a washout period) AND Patients exceeding doses over 6mg/24 hours will be on a tyramine restricted diet (i.e. avoid aged/spoiled/fermented meat and cheese, tap beer, fava beans, or any foods with high amounts of tyramine)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, the patient has improved or stabilized on Emsam.

# ENBREL (BCBS RI)

## Products Affected

- Enbrel

- Enbrel Sureclick

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis). Currently taking cyclophosphamide, other TNF antagonist, Anakinra, or Orencia
<b>Required Medical Information</b>	Diagnosis of one of the following : A) moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs for at least 3 consecutive months B) moderate to severe polyarticular juvenile idiopathic arthritis and patient had an inadequate response, intolerance or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months C) psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate D) ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to one or more NSAIDs E) moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had an inadequate response, intolerance or contraindication to conventional therapy with at least one of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (i.e. methotrexate, cyclosporine, acitretin, sulfasalazine) for at least 3 consecutive months.
<b>Age Restrictions</b>	2 years of age or older for JIA. 18 years of age or older for all other indications
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial 3 months (plaque psoriasis), 12 months (others). Renewal 12 months.

<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. Dosing as per FDA approved labeling. For renewal, patient has stable disease or has improved while on therapy (e.g., for pJIA, reduction in disease flares, improvement in ACR scoring. For RA, improvement in tender/swollen joint count, improvement in ACR scoring. For PsA, improvement in number of swollen/tender joints, pain, stiffness. For AS, improvement in AS symptoms, such as stiffness and back pain).
-----------------------	---



## EPIDUO (S)

### Products Affected

- Epiduo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Acne rosacea. Actinic keratoses (solar keratoses). Flat warts (verucca plana/juveniles). Hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans [Flegel's disease], palmoplantar hyperatosis, keratosis follicularis [Darier's disease]). Ichthyosis (i.e. congenital, lamellar, vulgaris, or X-linked)
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of mild to moderate acne vulgaris (including comedonal, cystic, and nodular) OR Diagnosis of acne rosacea and patient has experienced treatment failure of or intolerance to sulfacetamide, metronidazole, or azelaic acid OR patient is receiving treatment for actinic keratosis (or solar keratoses) with multiple lesions on the face OR Diagnosis of flat warts (verucca plana/jeveniles) OR Diagnosis of hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans [Flegel's disease], palmoplantar hyperatosis, keratosis follicularis [Darier's disease] OR Diagnosis of ichthyosis (i.e. congenital, lamellar, vulgaris, or X-linked)
<b>Age Restrictions</b>	PA applies to patients older than 26 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, medication has been effective in treating the patient's condition.

# EPOETIN ALFA (S)

## Products Affected

- Epogen

- Procrit

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Anemia secondary to hepatitis C. Anemia secondary to myelodysplastic syndrome.
<b>Exclusion Criteria</b>	uncontrolled hypertension. Pure red cell aplasia that begins after ESA treatment.
<b>Required Medical Information</b>	Pre-treatment hemoglobin level less than 10 g/dL (unless, for perioperative blood loss Hb should be greater than 10 but less than or equal to 13 g/dL) AND Patient has adequate iron stores prior to initiation of therapy defined as ferritin more than 100 mcg/L or serum transferrin saturation greater than 20% AND other causes of anemia such as iron deficiency, folate deficiency or B12 deficiency, hemolysis, gastrointestinal bleeding, other active or occult bleeding, or underlying hematologic disease (such as sickle cell anemia, thalassemia, and porphyria) have been ruled out AND Diagnosis of one of the following: A) Anemia due to chronic kidney disease (CKD) with or without hemodialysis, OR B) Anemia in a patient with non-myeloid malignancies where anemia is due to the effect of concomitant myelosuppressive chemotherapy and two additional months of chemotherapy is anticipated, C) Treatment of anemia in a patient at high risk for perioperative blood loss from elective, noncardiac, nonvascular surgery to reduce the need for allogeneic blood transfusion, D) Anemia in zidovudine-treated HIV infection with serum erythropoietin levels 500 mUnits/mL or less and zidovudine doses 4,200 mg/week or less E) Anemia secondary to hepatitis C therapy and patient is receiving ribavirin and interferon/peginterferon F) Anemia due to myelodysplastic syndrome and endogenous serum erythropoietin level is 500 mUnits/mL or less.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	CKD - prescribed by a nephrologist or hematologist. Non-myeloid malignancies , MDS - prescribed by an oncologist/hematologist. Surgery - Prescribed by a surgeon. HIV - Prescribed by an infectious disease specialist. HCV - ID specialist or gastroenterologist
<b>Coverage Duration</b>	Initial: 4 mos (others). Renewal: CKD-12 mos, Others -4 mos. Sx-3 mos

<b>Other Criteria</b>	For renewal of CKD, for dialysis patients: Hb less than 11 g/dL or physician will decrease or interrupt dose and for non-dialysis patients: Hb less than 10 g/dL or physician will decrease or interrupt dose. For renewal of non-myeloid malignancies: Concurrent myelosuppressive chemotherapy and Hb is 12g/dL or less and there is measurable response after eight weeks (defined as an increase in Hb 1 g/dL or more or a reduction in red blood cell transfusion requirements). For renewal of zidovudine-treated HIV, Hb is 12g/dL or less AND Zidovudine dose remains 4,200 mg/week or less and there is a measurable response after eight weeks (defined as an increase in Hb or a reduction in RBC transfusion requirements or documented dose escalation [up to max of 300 units/kg/dose]). For renewal of MDS, Hb is 12 g/dL or less and patient had HB rise of 1.5g/dL or a decrease in RBC transfusion requirements after at least 8 weeks of treatment OR patient will have a concomitant trial with G-CSF AND if patient had concomitant trial with G-CSF, patient had a measureable response to therapy after at least 8 weeks. For renewal of HCV, Hb 12 or less and concurrent ribavirin/interferon or pegylated interferon/ribavirin therapy. ESRD patients on dialysis excluded.
-----------------------	---

## ERIVEDGE (S)

---

### Products Affected

- Erivedge

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of metastatic basal cell carcinoma OR Diagnosis of locally advanced basal cell carcinoma that has recurred following surgery or when the patient is not a candidate for surgery and radiation
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ERWINAZE (S)

---

### Products Affected

- Erwinaze

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Member has experienced any of the following with prior asparaginase therapy: serious hypersensitivity reactions, including anaphylaxis, serious pancreatitis, serious thrombosis, serious hemorrhagic events
<b>Required Medical Information</b>	Diagnosis of acute lymphoblastic leukemia AND Patient has a hypersensitivity to E. coli-derived asparaginase
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ESBRIET (S)

---

### Products Affected

- Esbriet

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The patient has a diagnosis of idiopathic pulmonary fibrosis confirmed by a high resolution CT scan or biopsy AND the patient does not have evidence or suspicion of an alternative interstitial lung disease diagnosis AND liver function tests have been performed prior to start of therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient experienced stabilization from baseline or a less than 10 percent decline in force vital capacity AND the patient has not experienced AST or ALT elevations greater than 5 times the upper limit of normal or greater than 3 times the upper limit of normal with signs or symptoms of severe liver damage.

# EXALGO (S)

## Products Affected

- Hydromorphone Hcl Er

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	Significant respiratory depression. Acute or severe bronchial asthma. Known or suspected paralytic ileus. Gastrointestinal obstruction.
Required Medical Information	Diagnosis of severe pain requiring continuous, around-the-clock opioid analgesic for an extended period of time AND patient has tried and failed or unable to tolerate at least two generic extended-release opioid products, such as: oxymorphone ER, morphine ER, fentanyl, methadone, tramadol ER AND Patient is opioid tolerant, taking at least 60 mg oral morphine per day, 25 µg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, or an equianalgesic dose of another opioid for a week or longer
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Not covered for members enrolled in a Medicare approved hospice program

## EXJADE (S)

### Products Affected

- Exjade

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Myelodysplastic syndrome in a patient with Low or Intermediate-1 disease or is a potential transplant patient and who has received more than 20 red blood cell transfusions.
<b>Exclusion Criteria</b>	Creatinine clearance less than 40 mL/minute. Platelet count less than 50 x 10 <sup>9</sup> /L. Poor performance status. Severe (Child-Pugh class C) hepatic impairment. High-risk myelodysplastic syndromes. Advanced malignancies. Gastrointestinal ulceration or hemorrhage.
<b>Required Medical Information</b>	Patient has a diagnosis of one of the following: A) chronic iron overload due to blood transfusions and patient has a baseline ferritin level more than 1,000 mcg/L and the patient has required the transfusion of at least 100 mL/kg packed red blood cells OR B) chronic iron overload due to non-transfusion-dependent thalassemia (NTDT) and liver iron concentration (LIC) is 5 mg of iron per gram of liver dry weight (mg Fe/g dw) or higher and serum ferritin level is greater than 300 mcg/L OR C) Myelodysplastic syndrome (MDS) AND The patient has Low or Intermediate-1 disease or is a potential transplant patient and patient has received more than 20 red blood cell transfusions
<b>Age Restrictions</b>	2 years of age or older for chronic iron overload due to transfusions. 10 years of age or older for chronic iron overload due to NTDT
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	NTDT - 6 months. Transfusion-dependent anemia, MDS - 12 months.
<b>Other Criteria</b>	For renewal of chronic iron overload due to blood transfusions and MDS, the experienced a reduction in serum ferritin level or LIC. For renewal of chronic iron overload due to NTDT, patient has LIC 3 mg Fe/g dw or higher and patient experienced a reduction in serum ferritin level or LIC.



# FABRAZYME (S)

---

## Products Affected

- Fabrazyme INJ 35MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Fabry disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## FENTANYL (S)

### Products Affected

- Abstral
- Fentora
- Lazanda
- Subsys

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Management of acute or post-operative pain, including headache/migraine, dental pain, or use in the emergency room. Opioid non-tolerant patients.
<b>Required Medical Information</b>	Patient meets the following: A) Diagnosis of cancer and use is for breakthrough cancer pain, B) patient is opioid tolerant and taking at least 60 mg morphine/day, at least 25 mcg transdermal fentanyl/hour, at least 30 mg of oxycodone daily, at least 8 mg oral hydromorphone daily or an equianalgesic dose of another opioid for a week or longer, C) at least one other formulary short-acting strong narcotic analgesic alternatives (other than fentanyl) have been ineffective, not tolerated, or contraindicated, D) prescriber and patient are registered in the Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy Access program, E) for brand requests, generic transmucosal fentanyl citrate has been ineffective or not tolerated.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

# FERRIPROX (S)

---

## Products Affected

- Ferriprox

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of transfusional iron overload due to thalassemia syndromes AND patient has failed prior chelation therapy with Desferal or Exjade (failure is defined as a serum ferritin level greater than 2,500 mcg/L) or patient has a contraindication or intolerance to Desferal or Exjade AND Patient has an absolute neutrophil count greater than $1.5 \times 10^9/L$ .
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced at least a 20% reduction in serum ferritin levels and has an absolute neutrophil count greater than $0.5 \times 10^9/L$

## FIRMAGON (S)

---

### Products Affected

- Firmagon

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced or metastatic prostate cancer
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## FLECTOR (S)

---

### Products Affected

- Flector

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Previously experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Use for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery. Application to non-intact or damaged skin
<b>Required Medical Information</b>	Patient is experiencing acute localized pain due to minor strains, sprains and contusions AND patient had experienced treatment failure with at least 2 prescription strength oral NSAIDs or patient has a documented swallowing disorder OR has a history of peptic ulcer disease/gastrointestinal bleeding OR patient is more than 65 years of age with one additional risk factor for gastrointestinal adverse event (e.g., use of anticoagulants or chronic corticosteroids)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	N/A

# FORTEO (BCBS RI)

## Products Affected

- Forteo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Underlying hypercalcemia disorder such as primary hyperparathyroidism
<b>Required Medical Information</b>	Patient has a diagnosis of one of the following: a) osteoporosis in a postmenopausal female, b) primary or hypogonadal osteoporosis in a male, or c) osteoporosis associated with sustained systemic glucocorticoid therapy AND patient is considered to be at high-risk for fracture by meeting one or more of the following: A) history of osteoporotic fracture, B) multiple risk factors for fracture (including older age, female gender, prior osteoporotic fracture, low body mass index, rheumatoid arthritis, smoker, alcohol intake more than 3 drinks/day, parental history of hip fracture, oral glucocorticoid therapy or patient ever took prednisone at a dose of 5 mg or higher), or C) documented trial and failure of bisphosphonate, or D) documented contraindication or intolerance to bisphosphonate therapy. Patient has not received more than 2 years of therapy with Forteo.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve doses based on FDA labeling

# GATTEX (S)

---

## Products Affected

- Gattex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active gastrointestinal malignancy (gastrointestinal tract, hepatobiliary, pancreatic), colorectal cancer, or small bowel cancer
<b>Required Medical Information</b>	Diagnosis of short bowel syndrome AND patient is receiving specialized nutritional support (i.e. parenteral nutrition)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has a reduced need for parenteral support (20% reduction) after at least 6 months of therapy.

# GAZYVA (S)

---

## Products Affected

- Gazyva

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic lymphocytic leukemia (CLL) AND patient is treatment naïve AND Gazyva will be used in combination with chlorambucil AND patient has been screened for hepatitis B virus infection and does not have HBV reactivation and diagnosis of progressive multifocal leukoencephalopathy will be considered in any patient presenting with new onset or changes to pre-existing neurologic manifestations and discontinued if PML develops
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



# GILENYA (BCBS RI)

## Products Affected

- Gilenya

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. First clinical episode with MRI features consistent with multiple sclerosis.
<b>Exclusion Criteria</b>	Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure. History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker. Baseline QTc interval greater than or equal to 500 ms. Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol, bretylium, disopyramide, dofetilide, ibutilide). Or concomitant therapy with antineoplastics, immunosuppressants, and immunomodulating therapy.
<b>Required Medical Information</b>	Diagnosis of a relapsing form of multiple sclerosis or diagnosis of first clinical episode with MRI features consistent with MS AND Patient will be observed for signs and symptoms of bradycardia in a controlled setting for at least 6 hours after the first dose. Immunity to varicella: history of chickenpox, vaccinated against varicella zoster virus and positive antibody testing AND patients who are not currently stable on therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	For renewal, the patient has experienced a reduction in the number of MS relapses, improvement in object symptoms scores, or has not had MRI progression.

# GILOTRIF (S)

---

## Products Affected

- Gilotrif

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of metastatic non-small cell lung cancer AND patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions of exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND the medication will be used first-line.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# GLEEVEC (S)

## Products Affected

- Gleevec

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following in an adult: A) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B) Ph+ acute lymphoblastic leukemia (ALL), C) Gastrointestinal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E) hypereosinophilic syndrome or chronic eosinophilic leukemia, F) myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, G) aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutation or with c-KIT mutational status unknown. Diagnosis of one of the following in a pediatric patient: A) Ph+ CML that is newly diagnosed in the chronic phase B) newly diagnosed Ph+ ALL
<b>Age Restrictions</b>	18 years of age or younger - newly diagnosed CML in the chronic phase or newly diagnosed Ph+ ALL. 18 years of age or older for other indications.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## GRALISE (BCBS RI)

---

### Products Affected

- Gralise

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of post-herpetic neuralgia and patient has tried and failed a dose of at least 1800 mg of generic gabapentin or patient has experienced intolerance to generic gabapentin
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has improvement in pain severity.

# GROWTH HORMONE (S)

## Products Affected

- Genotropin
- Genotropin Miniquick
- Humatrope INJ 12MG, 24MG, 6MG
- Humatrope Combo Pack
- Norditropin Flexpro
- Norditropin Nordiflex Pen INJ 30MG/3ML
- Nutropin INJ 10MG
- Nutropin Aq
- Nutropin Aq Pen INJ 20MG/2ML
- Omnitrope INJ 5MG/1.5ML
- Saizen
- Saizen Click.easy
- Tev-tropin

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Growth hormone deficiency (GHD). Small for gestational age (SGA). Chronic renal insufficiency (CRI). Short stature homeobox-containing gene (SHOX) deficiency. Noonan syndrome. Prader-Willi Syndrome (PWS). Turner Syndrome. Adult- or childhood-onset GHD.
<b>Exclusion Criteria</b>	Child with closed epiphyses. Acute critical illness due to complications following open heart or abdominal surgery, multiple accidental trauma or acute respiratory failure. Active malignancy. Active proliferative or severe non-proliferative diabetic retinopathy. For PWS only: severe obesity, history of upper airway obstruction or sleep apnea, or severe respiratory impairment.

<b>Required Medical Information</b>	<p>Diagnosis of pediatric indication: A) GHD and one stim test with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B) SGA and birth wt or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height [ht] 2 or more SDS below mean for age and gender), C) CRI and nutritional status optimized, metabolic abnormalities corrected, and not had renal transplant D) SHOX deficiency or Noonan syndrome E) PWS confirmed by genetic testing AND ht below 3rd percentile or growth velocity (GV) measured over 1 year more than 2 SD below mean for age and sex, F) Turner Syndrome confirmed by chromosome analysis and ht below 5th percentile for age and sex. For GHD, CRI, SHOX deficiency, and Noonan syndrome, one of the following: ht more than 3 SDS below mean for age and gender, or ht more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of adult indication: A) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy) AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or CV complications AND Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH).</p>
<b>Age Restrictions</b>	SGA more than 2 years of age
<b>Prescriber Restrictions</b>	Pediatric endocrinologist for pediatric indications. Endocrinologist for adult indications.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal of pediatric indications, final adult height has not been reached as determined by the fifth percentile of adult height and growth velocity is more than 2 cm/year. For renewal of adult indications, patient has experienced an improvement or normalization of IGF-1 levels (not a requirement in patients with panhypopituitarism).

## H.P. ACTHAR GEL (BCBS RI)

### Products Affected

- H.p. Acthar

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Infantile spasms, MS, psoriatic arthritis, rheumatoid arthritis, juvenile idiopathic arthritis, ankylosing spondylitis, SLE, systemic dermatomyositis, severe erythema multiforme, Stevens-Johnson Syndrome, serum sickness, severe acute & chronic allergic and inflammatory processes involving the eye and its adnexa such as: keratitis, iritis, iridocyclitis, diffuse posterior uveitis and choroiditis, optic neuritis, chorioretinitis, anterior segment inflammation, symptomatic sarcoidosis, to induce a diuresis or a remisison of proteinuria in the nephrotic syndrome without uremia of the idiopathic type or that due to lupus
<b>Age Restrictions</b>	2 years of age or younger for diagnosis of IS, 18 years of age or older in MS
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	IS-plan year, Collagen diseases - 6 months. Others- 1 month
<b>Other Criteria</b>	For rheumatic disorders: Failure of 12 week course of corticosteroid and concomitant therapy with a 3 month course of biologic DMARD. For MS: Failure of 3 to 5 day course of corticosteroid and concomitant use of a 3 month course of an immunomodulator

# HARVONI (BCBS RI)

## Products Affected

- Harvoni

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Mixed genotypes AND Patients who have received Olysio/Sovaldi treatment
<b>Required Medical Information</b>	Diagnosis of Hepatitis C infection genotype 1 only AND medication will be used as monotherapy OR Harvoni will be used in combination with ribavirin in one of the following settings: Post liver transplant, sofosbuvir/RBV/Peg- or SOF/RBV-treatment experienced patients AND Patient is not currently receiving Hep C treatment for GT1 AND will not be used with the following medications: rifampin, St. Johns Wort, carbamazepine, Tipranavir (Aptivus), phenobarbital, phenytoin, oxcarbazepine, rifabutin, rifapentine AND daily dose of any acid reducing medication does not exceed: Prevacid 15mg, omeprazole 20mg, Protonix 20mg, Aciphex 20mg, Zantac 150mg, Pepcid 40mg
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a gastroenterologist, hepatologist or ID only
<b>Coverage Duration</b>	8 weeks, 12 weeks, or 24 weeks depending patient specific parameters outlined in Other Criteria
<b>Other Criteria</b>	8 weeks: treatment naïve patients (never taken any Hep C medication ) without cirrhosis whose initial HCV RNA levels are less than 6 million IU/ml 12 weeks: treatment naïve patients (never taken any Hep C medication) without cirrhosis whose initial HCV RNA levels are greater than 6 million IU/ml OR treatment experienced patients with a non-sofosbuvir regimen without cirrhosis 24 weeks: treatment experienced patients with a non-sofosbuvir regimen with cirrhosis 12 weeks: when used with ribavirin for patients who are 6 months post liver transplant or for patients who are treatment experienced with a sofosbuvir plus RBV with or without PEG



## HEPSERA (S)

---

### Products Affected

- Adefovir Dipivoxil
- Hepsera

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis B AND patient is HBsAg-positive for at least 6 months AND For HBeAg-positive patients, serum HBV DNA greater than 20,000 IU/mL (105 copies per mL) and for HBeAg-negative patients, serum HBV DNA greater than 2,000 IU/mL (104 copies/mL) AND Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) at least 2 times the upper limit of normal or histologically active disease (i.e. necroinflammation on biopsy)
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient must be HBeAg negative and have not had HBsAg clearance OR HBeAg positive and have detectable HBV DNA and have not been anti-Hbe for at least 6 months

## HEXALEN (S)

---

### Products Affected

- Hexalen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of persistent or recurrent ovarian cancer AND the medication will be used as palliative treatment AND the medication will be used as a single agent AND the medication will be used following first-line therapy with a cisplatin and/or alkylating agent-based combination.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## HORIZANT (S)

---

### Products Affected

- Horizant

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of moderate to severe primary restless leg syndrome and patient has experienced treatment failure or is intolerant to ropinirole or pramipexole OR Patient has a diagnosis of post-herpetic neuralgia and patient tried and failed a dose of at least 1800 mg of generic gabapentin or patient has experienced intolerance to generic gabapentin
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient had a measurable response to therapy (e.g. decrease in symptoms onset or severity, improved sleep, decrease in symptom intensity for RLS or decreased pain severity for PHN)

# HRM - ANALGESICS

## Products Affected

- Meperidine Hcl INJ 25MG/ML, 50MG/ML
- Meperidine Hcl ORAL SOLN
- Meperitab
- Pentazocine/acetaminophen
- Pentazocine/naloxone Hcl
- Talwin

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
Age Restrictions	PA applies to patients 65 years or older
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Mild pain: acetaminophen, codeine. Moderate to severe pain: short-term NSAIDs, tramadol, tramadol/APAP, morphine sulfate, hydrocodone/APAP, oxycodone, oxycodone/APAP, fentanyl. Not covered for members enrolled in a Medicare approved hospice program

# HRM - ANTIDEPRESSANTS

## Products Affected

- Amitriptyline Hcl ORAL TABS
- Clordiazepoxide/amitriptyline
- Clomipramine Hcl ORAL CAPS
- Doxepin Hcl CONC
- Doxepin Hcl ORAL CAPS
- Imipramine Hcl ORAL TABS
- Imipramine Pamoate
- Perphenazine/amitriptyline
- Surmontil
- Trimipramine Maleate ORAL CAPS

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Applies to New Starts only.TCA: nortriptyline, desipramine, low-dose doxepin, trazodone. Depression: SSRI, SNRI, mirtazapine, bupropion. duloxetine, gabapentin.

# HRM - ANTIEMETIC DRUGS

## Products Affected

- Diphenhydramine Hcl CAPS 50MG
- Phenadoz
- Phenergan RECTAL SUPP
- Promethazine Hcl INJ
- Promethazine Hcl ORAL TABS
- Promethazine Hcl RECTAL SUPP
- Promethazine Hcl SYRP
- Promethegan
- Trimethobenzamide Hcl CAPS
- Trimethobenzamide Hcl INJ

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program.

## HRM - ANTIHISTAMINES

### Products Affected

- Arbinoxa
- Carbinoxamine Maleate SOLN
- Carbinoxamine Maleate TABS
- Clemastine Fumarate SYRP
- Clemastine Fumarate TABS 2.68MG
- Cyproheptadine Hcl SYRP
- Cyproheptadine Hcl TABS
- Diphenhydramine Hcl INJ
- Hydroxyzine Hcl INJ
- Hydroxyzine Hcl ORAL SOLN
- Hydroxyzine Hcl ORAL TABS
- Hydroxyzine Pamoate ORAL CAPS
- Promethazine Vc Plain

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## HRM - ANTIHYPERTENSIVE AGENTS

### Products Affected

- Guanfacine Hcl

- Reserpine TABS 0.25MG

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Low dose thiazide or a second generation calcium channel blocker OR ACE inhibitor, ARB, beta-blocker or combination product based on specific chronic conditions



## HRM - ANTI-INFECTIVE (BCRI)

### Products Affected

- Macrodantin CAPS 25MG
- Nitrofurantoin SUSP
- Nitrofurantoin Macrocrystals CAPS 50MG
- Nitrofurantoin Monohydrate

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	N/A
Required Medical Information	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) when used longer than 90 cumulative days and wishes to proceed with the originally prescribed medication AND intended duration of therapy will be verified
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) when used longer than 90 days and wishes to proceed with the originally prescribed medication AND intended duration of therapy will be verified.

## HRM - ANTIPARKINSON AGENTS

---

### Products Affected

- Benztropine Mesylate INJ
- Benztropine Mesylate ORAL TABS
- Trihexyphenidyl Hcl

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## HRM - ANXIOLYTICS

---

### Products Affected

- Meprobamate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program.

# HRM - CALCIUM CHANNEL BLOCKERS, DIHYDROPYRIDINE

## Products Affected

- Nifedipine ORAL CAPS

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	extended-release nifedipine, nicardipine, amlodipine

## HRM - DEMENTIA AGENTS

---

### Products Affected

- Ergoloid Mesylates TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	donepezil, galantamine, rivastigmine, memantine

# HRM - ORAL AND TRANSDERMAL ESTROGENS AND PROGESTINS

## Products Affected

- Alora
- Cenestin
- Climara Pro
- Combipatch
- Divigel TRANSDERMAL GEL  
0.25MG/0.25GM, 0.5MG/0.5GM
- Enjuvia
- Estradiol ORAL TABS
- Estradiol TRANSDERMAL PTTW
- Estradiol TRANSDERMAL PTWK
- Estradiol/norethindrone Acetate
- Estropipate ORAL TABS
- Jinteli
- Menest
- Menostar
- Mimvey
- Mimvey Lo
- Ortho-est
- Prefest
- Premarin ORAL TABS
- Prempro

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## HRM - PLATELET INHIBITORS

---

### Products Affected

- Ticlopidine Hcl

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	clopidogrel, dipyridamole with aspirin

## HRM - SEDATIVE HYPNOTIC AGENTS

---

### Products Affected

- Edluar
- Zolpidem Tartrate
- Zolpidem Tartrate Er
- Zaleplon

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) when used longer than 90 days and wishes to proceed with the originally prescribed medication AND intended duration of therapy will be verified.
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## HRM - SKELETAL MUSCLE RELAXANTS

---

### Products Affected

- Amrix
- Carisoprodol TABS 350MG
- Carisoprodol/aspirin
- Carisoprodol/aspirin/codeine
- Chlorzoxazone TABS
- Cyclobenzaprine Hcl ORAL TABS
- Methocarbamol ORAL TABS
- Orphenadrine Citrate INJ
- Orphenadrine Citrate Er

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## HRM - SULFONYLUREAS

### Products Affected

- Chlorpropamide
- Glyburide ORAL TABS
- Glyburide Micronized
- Glyburide/metformin Hcl

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	glimepiride, glipizide

## HRM - VASODILATORS

---

### Products Affected

- Dipyridamole ORAL TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The drug is being prescribed for an FDA-approved indication AND the patient has tried and failed at least one non-HRM alternative (listed in OTHER CRITERIA) AND the prescribing physician has been made aware that the requested drug is considered a high risk medication for elderly patients (age 65 years and older) and wishes to proceed with the originally prescribed medication .
<b>Age Restrictions</b>	PA applies to patients 65 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Requires trial of at least one Non-HRM alternative: clopidogrel, Aggrenox

# HUMIRA (S)

## Products Affected

- Humira
- Humira Pediatric Crohns Disease Starter Pack
- Humira Pen-crohns Diseasestarter

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis)
<b>Required Medical Information</b>	Diagnosis of one of the following: A) moderate to severe rheumatoid arthritis and inadequate response, intolerance, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) (e.g., hydroxychloroquine [HCQ], sulfasalazine, methotrexate [MTX], leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months B) moderate to severe polyarticular juvenile idiopathic arthritis and inadequate response, intolerance or contraindication to one or more non-biologic DMARDs (e.g., HCQ, sulfasalazine, MTX, leflunomide, azathioprine, cyclosporine) for at least 3 consecutive months C) psoriatic arthritis and inadequate response, intolerance, or contraindication to MTX D) ankylosing spondylitis and inadequate response, intolerance or contraindication to one or more NSAIDs E) moderate to severe chronic plaque psoriasis (affecting more than 5% of body surface area or crucial body areas such as the hands, feet, face, or genitals) and inadequate response, intolerance or contraindication to at least one of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments (e.g., MTX, cyclosporine, acitretin, sulfasalazine) for at least 3 consecutive months F) moderate to severe Crohn's disease and inadequate response, intolerance, or contraindication to two or more of the following: corticosteroids (e.g., prednisone, methylprednisolone) or non-biologic DMARDs (e.g., azathioprine, MTX, mercaptopurine) G) moderate to severe ulcerative colitis and inadequate response, intolerance or contraindication to two or more of the following: corticosteroids (e.g., prednisone, methylprednisolone), 5-ASA (i.e. mesalamine, sulfasalazine, balsalazide, olsalazine) or non-biologic DMARDs (azathioprine, MTX, mercaptopurine).
<b>Age Restrictions</b>	4 years of age or older for JIA. 18 years of age or older for all other indications

<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 16 weeks (CD), 12 weeks (UC), 12 months (others). Renewal - 12 months.
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated as per guidelines. For renewal, patient has stable disease or has improved while on therapy (For pJIA, reduction in disease flares, improvement in ACR scoring. For RA, improvement in tender/swollen joint count, improvement in ACR scoring. For PsA, improvement in number of swollen/tender joints, pain, stiffness. For AS, improvement in AS symptoms, such as stiffness and back pain. For CD, symptomatic remission. For UC, clinical remission, reduction in steroid use.)

# HYQVIA (S)

---

## Products Affected

- Hyqvia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	History of hypersensitivity (including anaphylaxis or severe systemic reaction) to immune globulin or any component of the preparation. Hyperprolinemia.
<b>Required Medical Information</b>	Medication will be given by the subcutaneous route and patient has a diagnosis of one of the following types of primary immunodeficiencies: hypogammaglobulinemia, severe combined immunodeficiency, X-linked immunodeficiency with hyperimmunoglobulin M, congenital agammaglobulinemia, common variable immunodeficiency, Wiskott-Aldrich syndrome, Hyperimmunoglobulinemia E syndrome, selective antibody deficiency
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Part B: Patient has a diagnosis of primary immune deficiency and the medication will be provided in the home. Part D: Medication will be given in the home and the patient has a diagnosis other than PID OR the medication will not be given in the home.

## HYSINGLA (BCBS RI)

---

### Products Affected

- Hysingla Er

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Significant respiratory depression. Acute or severe bronchial asthma. Hypercarbia. Known or suspected paralytic ileus.
<b>Required Medical Information</b>	Diagnosis of severe pain requiring continuous around the clock long-term opioid treatment AND Patient has tried and failed or unable to tolerate at least two generic extended-release opioid products, such as: oxycodone ER, morphine ER, fentanyl, tramadol ER AND Patient is opioid tolerant for 1 week or longer and taking at least: 60mg morphine/day, 25mcg transdermal fentanyl /hour, 30mg of oxycodone daily, 8mg of hydromorphone daily AND Current opioid will be discontinued with the inception of Hysingla ER
<b>Age Restrictions</b>	18 years or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

# ICLUSIG (S)

---

## Products Affected

- Iclusig

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic myelogenous leukemia (CML) and patient has tried and failed, resistance, relapse or contraindication to TWO other FDA-approved tyrosine kinase inhibitors OR Diagnosis of CML and the patient has a known T315I mutation OR Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia and the patient has tried and failed, resistance, relapse or contraindication to TWO other FDA-approved tyrosine kinase inhibitors.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## ILARIS (S)

---

### Products Affected

- Ilaris

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cryopyrin-associated period syndromes (CAPS), including Familial Cold Auto-inflammatory Syndrome (FCAS) and/or Muckle-Wells Syndrome (MWS) OR Diagnosis of active systemic juvenile idiopathic arthritis (sJIA) AND patient has tried and had an inadequate response, contraindication or intolerance to corticosteroids (e.g., prednisone, methylprednisolone) or methotrexate
<b>Age Restrictions</b>	CAPS - 4 years of age or older. sJIA - 2 years of age or older
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist, or other medical specialist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient experienced disease stability or improvement.

# IMBRUVICA (S)

---

## Products Affected

- Imbruvica

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of mantle cell lymphoma AND patient has relapsed or is refractory to at least one prior therapy for the treatment of MCL or Diagnosis of chronic lymphocytic leukemia and patient has relapsed or is refractory to at least one prior therapy for the treatment of CLL.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## INCRELEX (S)

---

### Products Affected

- Increlex

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Closed epiphyses. Active or suspected malignancy.
<b>Required Medical Information</b>	Diagnosis of growth failure in a child with severe primary IGF-1 deficiency, defined as height standard deviation score (SDS) less than or equal to -3.0 AND basal IGF-1 SDS less than or equal to -3.0 AND normal or elevated growth hormone OR Diagnosis of growth hormone gene deletion with development of neutralizing antibodies to growth hormone AND other causes of IGF-1 deficiency (e.g., hypothyroidism, nutritional deficiencies, pituitary disorders, etc.) have been ruled out or corrected prior to initiating therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Pediatric endocrinologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient had a minimum growth rate of at least 2 cm/year.

## INFERGEN (S)

---

### Products Affected

- Infergen

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Uncontrolled depression. Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon and ribavirin.
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis C with compensated liver disease AND patient will be receiving combination therapy with ribavirin, unless ribavirin is contraindicated AND Infergen will not be used as part of triple therapy with a protease inhibitor AND Patient has a clinical reason for not using peginterferon
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	48 weeks
<b>Other Criteria</b>	N/A

# INHALED TOBRAMYCIN (BCBS RI)

---

## Products Affected

- Tobi Podhaler

- Tobramycin NEBU

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis AND patient has evidence of P. aeruginosa in the lungs AND has the patient tried and failed generic tobramycin inhalation solution
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations). For inhalation solution only (does not apply to TOBI Podhaler): Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## INLYTA (S)

---

### Products Affected

- Inlyta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced renal cell carcinoma AND patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# INTRON A (S)

## Products Affected

- Intron A

- Intron-a INJ 6000000UNIT/ML
- Intron-a W/diluent

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Uncontrolled depression. Solid organ transplant other than liver. Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon and ribavirin.
<b>Required Medical Information</b>	Diagnosis of hairy cell leukemia OR Diagnosis of Condylomata acuminata OR Diagnosis of AIDS-related Kaposi's sarcoma OR Clinically aggressive follicular lymphoma and the medication will be used concurrently with anthracycline-containing chemotherapy or is not a candidate for anthracycline-containing chemotherapy OR Malignant melanoma and the request for coverage is within 56 days of surgery and the patient is at high risk of disease recurrence OR Diagnosis of chronic hepatitis B with compensated liver disease and patient has evidence of hepatitis B viral replication and patient has been serum hepatitis B surface antigen-positive for at least 6 months OR Diagnosis of chronic hepatitis C with compensated liver disease and is receiving combination therapy with ribavirin, unless ribavirin is contraindicated, and the medication will not be used as part of triple therapy with a protease inhibitor and patient has a clinical reason for not using peginterferon
<b>Age Restrictions</b>	1 year of age or older for HBV. 3 years of age or older for HCV. 18 years of age or older for other indications.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Condylomata: 3 mos. HBV e antigen pos: 16 wks, e antigen neg: 48 wks. KS: 16 wks. Others: 12 mos
<b>Other Criteria</b>	N/A

# IVIG (S)

## Products Affected

- Bivigam
- Carimune Nanofiltered INJ 6GM
- Flebogamma Dif INJ 10%, 10%, 10%
- Gammagard Liquid
- Gammaked
- Gamunex-c
- Hizentra
- Octagam INJ 2GM/20ML
- Privigen INJ 20GM/200ML

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	History of hypersensitivity (including anaphylaxis or severe systemic reaction) to immune globulin or any component of the preparation. Privigen only: hyperprolinemia. Octagam only: allergy to corn.
<b>Required Medical Information</b>	Medication will be given intravenously and the patient has one of the following: A) idiopathic thrombocytopenic purpura (ITP) after trial of corticosteroids unless the platelet count is less than 30,000 cells/mm <sup>3</sup> , B) Kawasaki syndrome, C) hypogammaglobulinemia (Ig level less than 500 mg/dL) and recurrent bacterial infection associated with B-cell chronic lymphocytic leukemia, D) chronic inflammatory demyelinating polyneuropathy (CIDP), E) multifocal motor neuropathy (MMN). OR Medication will be given by the subcutaneous route or intravenous route and patient has a diagnosis of one of the following types of primary immunodeficiency (PI): hypogammaglobulinemia, severe combined immunodeficiency, X-linked immunodeficiency with hyperimmunoglobulin M, congenital agammaglobulinemia, common variable immunodeficiency, Wiskott-Aldrich syndrome, Hyperimmunoglobulinemia E syndrome, selective antibody deficiency. For IV administration only: Patient will not take the medication at the minimum concentration available and the minimum infusion rate practicable if they have one of the following conditions: Pre-existing renal insufficiency, diabetes mellitus, volume depletion, sepsis, paraproteinemia, age over 65 years, receiving known nephrotoxic drugs.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	ITP, Kawasaki - 3 months, Other - 12 months.



<b>Other Criteria</b>	Part B: Patient has a diagnosis of primary immune deficiency and the medication will be provided in the home. Part D: Medication will be given in the home and the patient has a diagnosis other than PID OR the medication will not be given in the home.
-----------------------	--

# JAKAFI (S)

## Products Affected

- Jakafi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of myelofibrosis (primary, post-pocythemia vera or post-essential thrombocythemia) AND patient has two or more of the following: age older than 65 years, white blood cell count greater than 25 x 10 <sup>9</sup> /L, hemoglobin less than 10 g/dL, peripheral blasts more than 1%, constitutional symptoms (e.g., night sweats, fevers, unintentional weight loss, debilitating fatigue)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	For renewal, the patient experienced one of the following: At least 35% reduction in spleen volume from baseline as measured by CT or MRI or a 50% reduction in spleen size from baseline based on palpation OR 2 g/dL or greater increase in hemoglobin level (in transfusion-independent) or becoming transfusion independent (for transfusion dependent) OR Improvement in symptoms (i.e. abdominal discomfort, pain under left ribs, early satiety, night sweats, itching, bone or muscle pain) without progressive splenomegaly or worsening of anemia (i.e. newly transfusion dependent or hemoglobin reduction by 2 g/dL that persists for at least 12 weeks), thrombocytopenia (more than 2-grade decline but above 25,000 x 10 <sup>9</sup> /L) or neutropenia (more than 2-grade decline but above 0.5 x 10 <sup>9</sup> /L)

# JUXTAPID (BCBS RI)

## Products Affected

- Juxtapid

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests. Pregnancy. Concomitant use with strong or moderate CYP3A4 inhibitors.
<b>Required Medical Information</b>	Diagnosis of homozygous familial hypercholesterolemia as evidenced by one of the following: A) genetic confirmation of 2 mutant alleles at the LDL receptor, ApoB, PCSK9, or autosomal recessive hypercholesterolemia (ARH) adaptor protein gene locus OR B) untreated/pre-treatment LDL greater than 500 mg/dL with at least one of the following: cutaneous or tendonous xanthoma before age 10 years, history of early vascular disease (men younger than 55 years, women younger than 60 years) on both sides fo the family if parenteral LDL levels are unknown, elevated LDL cholesterol levels before lipid-lowering therapy consistent with heterozygous FH in both parents where LDL levels are known: LDL cholesterol more than 250 mg/dL in a patient 30 years of age or older, LDL cholesterol greater than 220 mg/dL for patients 20 to 29 years of age, LDL cholesterol greater than 190 mg/dL in patients younger than 20 years AND Medication will be used as adjunct to a low-fat diet and other lipid-lowering treatments AND Patient has tried and had an inadequate response to the maximum tolerated dose of a high potency statin (e.g., atorvastatin, rosuvastatin), unless all statin are contraindicated
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider and patient must be registered in the Juxtapid REMS program
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	For renewal, patient has responded to therapy with a decrease in LDL levels from baseline AND patient does not have contraindications to therapy.

# KADCYLA (BCBS RI)

---

## Products Affected

- Kadcyla INJ 100MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of HER2-positive metastatic breast cancer and the member has been previously treated with trastuzumab and a taxane
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Prescriber has assessed the patient's hepatic function and left ventricular ejection fraction prior to initiation of therapy. Female patients of child-bearing potential had pregnancy status verified prior to the initiation of Kadcyla and have been advised of the risk of fetal harm and the need for contraception. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

# KALYDECO (S)

## Products Affected

- Kalydeco

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis AND Patient has one of the following mutations: G551D, G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N, or S549R on at least one allele in the cystic fibrosis transmembrane conductance regulator gene documented by an FDA-cleared cystic fibrosis-mutation test and followed by verification with bi-directional sequencing when recommended by the mutation test instructions
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced benefit from therapy (i.e. improvement in pulmonary lung function [FEV1], decreased number of pulmonary exacerbations)

# KEYTRUDA (S)

---

## Products Affected

- Keytruda INJ 50MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The patient has a diagnosis of metastatic melanoma AND the patient has experienced disease progression following therapy with Yervoy AND if the patient is positive for a BRAF V600E or V600K mutation, the patient has experienced disease progression following therapy with one of the following: Zelboraf, Tafinlar, Mekinist
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# KINERET (BCBS RI)

## Products Affected

- Kineret

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis) Concomitant use with TNF or Orencia
<b>Required Medical Information</b>	Diagnosis of moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months OR Diagnosis of cryopyrin-associated periodic syndrome (CAPS) with neonatal-onset multisystem inflammatory disease (NOMID)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	For CAPS, diagnosed by, or upon consultation with or recommendation of, an immunologist, allergist, dermatologist, rheumatologist, neurologist or other medical specialist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient has been tested for TB in the past year and latent TB has been ruled out or is being treated per guidelines. Dosing as per the FDA labeling for rheumatoid arthritis. For renewal, patient has stable disease or improved on treatment (For RA, improvement in tender/swollen joint count, improvement in ACR scoring. For CAPS, symptom improvement, improvement in serum markers of inflammation)

# KORLYM (S)

## Products Affected

- Korlym

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnancy. Patient requires concomitant treatment with long-term corticosteroids (e.g., immunosuppression for organ transplant). History of unexplained vaginal bleeding. Endometrial hyperplasia with atypia or endometrial carcinoma. Concomitantly taking simvastatin, lovastatin, or a CYP3A substrate with a narrow therapeutic index (e.g., cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus)
<b>Required Medical Information</b>	Diagnosis of endogenous Cushing's syndrome AND diagnosis of type 2 diabetes mellitus or glucose intolerance AND Patient has hyperglycemia secondary to hypercortisolism AND patient has failed or is not a candidate for surgery
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient had a positive response to therapy with Korlym



# KUVAN (S)

## Products Affected

- Kuvan

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of phenylketonuria (PKU) and patient is and will be maintained on a phenylalanine-restricted diet
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial: 2 months. Renewal: 12 months.
<b>Other Criteria</b>	For initial approval, Patient will have phenylalanine levels measured one week after starting therapy and periodically for up to two months of therapy to determine response. For renewal, patient has been determined to be a responder to therapy (i.e. phenylalanine levels have decreased by at least 30% from baseline) and phenylalanine levels will be measured periodically during therapy.

# KYNAMRO (S)

## Products Affected

- Kynamro

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests.
<b>Required Medical Information</b>	Diagnosis of homozygous familial hypercholesterolemia as evidenced by one of the following: A) genetic confirmation of 2 mutant alleles at the LDL receptor, ApoB, PCSK9, or autosomal recessive hypercholesterolemia (ARH) adaptor protein gene locus OR B) untreated/pre-treatment LDL greater than 500 mg/dL with at least one of the following: cutaneous or tendonous xanthoma before age 10 years, history of early vascular disease (men younger than 55 years, women younger than 60 years) on both sides of the family if parenteral LDL levels are unknown, elevated LDL cholesterol levels before lipid-lowering therapy consistent with heterozygous FH in both parents AND Medication will be used as adjunct to a low-fat diet and other lipid-lowering treatments AND Patient has tried and had an inadequate response to the maximum tolerated dose of a high potency statin (e.g., atorvastatin, rosuvastatin), unless all statin are contraindicated
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	For renewal, patient has responded to therapy with a decrease in LDL levels from baseline AND patient does not have contraindications to therapy.

## LEMTRADA (BCBS RI)

---

### Products Affected

- Lemtrada

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	HIV infection. Active or latent TB. Active infections. Concurrent antineoplastic/immunosuppressive therapy. History of PML.
<b>Required Medical Information</b>	Diagnosis of relapsing multiple sclerosis AND Clinical failure for efficacy (defined as relapse or increased active MRI lesions) to an interferon beta product and at least one agent in a different MS class AND Clinical failure will be confirmed with evidence of medication adherence to treatment by pharmacist via claims data AND Lemtrada will be used as monotherapy.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Total duration of therapy: 24 months lifetime
<b>Other Criteria</b>	Dosing: 12mg daily for 5 days (total of 60mg), 12 months later: 12mg daily for 3 days (total of 36mg). Premedicate with corticosteroids for initial 3 days of therapy (1 gm of methylprednisolone).

## LETAIRIS (BCBS RI)

---

### Products Affected

- Letairis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class II or III AND pregnancy must be excluded prior to the start of therapy and will be prevented thereafter with two forms of reliable contraception in female patients of reproductive potential.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Written by or in consultation with a pulmonologist or cardiologist
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	N/A

# LEUKINE (S)

## Products Affected

- Leukine

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Excessive leukemic myeloid blasts (10% or higher) in bone marrow/peripheral blood. Simultaneous use with cytotoxic chemotherapy or radiotherapy.
<b>Required Medical Information</b>	<p>A) malignant melanoma, B) Adjunct therapy (tx) for severe febrile neutropenia (FN) and receiving myelosuppressive tx for non-myeloid malignancy and one: Received prophylactic CSF (not Neulasta) OR Did not receive prophylactic CSF and used as adjunct to antibiotics in high-risk pt with any one: 65 years or older, Pneumonia, Hypotension and multiorgan dysfunction (sepsis syndrome), Invasive fungal infection or clinically-documented infection, Hospitalized when fever developed, Prior FN, Severe (ANC less than 100/mcL) or anticipated prolonged (more than 10 days) neutropenia, C) Autologous peripheral-blood progenitor cell transplant for collection by leukapheresis, D) Myeloablative chemotx for non-myeloid malignancy followed by BMT, E) Acute myeloid leukemia after completing induction/consolidation chemotx, F) Acute lymphoblastic leukemia after completing first few days of chemotx of initial induction or first post-remission course, G) Myelodysplastic syndrome with severe neutropenia and recurrent infection, H) Receiving radiation tx, not chemotx, and expect prolonged tx delays due to neutropenia, I) Neutropenia due to HIV infection and antiretroviral tx, J) Aplastic anemia, K) Primary prophylaxis of FN and one: at least 20% FN risk based on chemotx OR 10% up to 20% risk with one of the following: 65 years or older, Poor performance status, Poor nutritional status, Previous FN, Extensive prior tx with large radiation ports, Cytopenias due to bone marrow involvement by tumor, Receiving combined chemoradiotx, Open wounds or active infections, Other serious comorbidity (e.g., renal or liver dysfunction) OR less than 10% risk and intent is curative or adjuvant and at risk for serious medical consequences, including death L) Myelosuppressive chemotx for non-myeloid malignancy. M) Secondary prophylaxis of FN in pt with neutropenic complication from prior chemotx cycle (where primary prophylaxis was not given) N) Allogeneic or autologous bone marrow transplant</p>
<b>Age Restrictions</b>	N/A

<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# LIDODERM (S)

---

## Products Affected

- Lidocaine PTCH

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of post-herpetic neuralgia.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has improvement in pain severity.

# LINZESS (S)

## Products Affected

- Linzess

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Mechanical gastrointestinal obstruction.
<b>Required Medical Information</b>	Diagnosis of irritable bowel syndrome-constipation for at least 12 non-consecutive weeks and patient has tried and failed increasing fluid and fiber intake and patient has tried and failed or has an intolerance to osmotic laxatives, stimulant laxatives or probiotics OR Diagnosis of chronic idiopathic constipation for at least 3 months and patient has tried and failed increasing fluid and fiber intake and patient has tried and failed or has an intolerance to osmotic laxatives, stimulant laxatives or stool softeners.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 4 months. Renewal - 12 months
<b>Other Criteria</b>	For renewal, the patient has experienced an increase in the number of bowel movements.



# LUMIZYME (S)

---

## Products Affected

- Lumizyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of late (non-infantile) onset Pompe disease (GAA) deficiency and the patient does not have evidence of cardiac hypertrophy
<b>Age Restrictions</b>	8 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# LUPANETA (S)

## Products Affected

- Lupaneta Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Undiagnosed abnormal uterine bleeding. Breastfeeding. Known, suspected or history of breast cancer or other hormone-sensitive cancer. Thrombotic or thromboembolic disorders. Liver tumors or liver disease.
<b>Required Medical Information</b>	Diagnosis of endometriosis And patient is undergoing an initial treatment course and has had an inadequate pain control response, intolerance or contraindication to one of the following: Danazol, Combination [estrogen/progesterone] oral contraceptives, Progestins OR Patient is undergoing a recurrent treatment course and is experiencing a reappearance of symptoms after an initial course of leuprolide therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	In patients of child-bearing potential, pregnancy has been excluded and patient will use non-hormonal contraception during and for 12 weeks after therapy.

# LUPRON DEPOT (S)

## Products Affected

- Leuprolide Acetate INJ
- Lupron Depot INJ 22.5MG, 3.75MG, 30MG, 45MG, 7.5MG

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnancy in patients with child-bearing potential. Breastfeeding. Undiagnosed abnormal vaginal bleeding.
<b>Required Medical Information</b>	Diagnosis of one of the following: A) advanced or metastatic prostate cancer (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), B) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only) and for initial, patient has had an inadequate pain control response or patient has an intolerance or contraindication to one of the following: Danazol OR Combination [estrogen/progesterone] Oral Contraceptives OR Progestins and for retreatment course, Patient is experiencing recurrence of symptoms after an initial course of therapy with leuprolide acetate and Norethindrone acetate 5 mg daily will be co-administered, or C) Anemia due to uterine Leiomyomata (Fibroids) (3.75 mg 1-month & 11.25 mg 3-month depots only) and Patient is preoperative and Patient has tried and had an inadequate response to monotherapy with iron and Patient will be receiving concomitant iron therapy while on leuprolide.
<b>Age Restrictions</b>	Uterine fibroids - 18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Endometriosis - 6 months, Uterine fibroids - 3 months, Prostate cancer - 12 months.
<b>Other Criteria</b>	For endometriosis and uterine fibroids, patient will be using nonhormonal contraception during and for 12 weeks after therapy.

## LUPRON DEPOT- PED (S)

### Products Affected

- Lupron Depot INJ 11.25MG
- Lupron Depot-ped INJ 11.25MG, 11.25MG, 15MG

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of central precocious puberty and patient had early onset of secondary sexual characteristics (male: earlier than 9 years of age. female: earlier than 8 years of age) and advanced bone age of at least one year compared with chronological age and has undergone gonadotropin-releasing hormone agonist (GnRHa) testing with peak luteinizing hormone (LH) level above pre-pubertal range or random LH level in pubertal range and Patient had the following diagnostic evaluations to rule out tumors, when suspected: diagnostic imaging of the brain (MRI or CT scan), Pelvic/testicular/adrenal ultrasound (if steroid levels suggest suspicion), Human chorionic gonadotropin levels (in all boys), Adrenal steroids to rule out congenital adrenal hyperplasia
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pediatric endocrinologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal of CPP, LH levels have been suppressed to pre-pubertal levels and consideration for discontinuation of therapy when the patient is 11 years of age for girls and 12 years of age for boys.

# MARINOL (S)

## Products Affected

- Dronabinol

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chemotherapy-induced nausea and vomiting AND patient has tried and failed conventional antiemetic treatments (e.g., aprepitant/fosaprepitant, dexamethasone, t-hydroxytriptamine-3 serotonin receptor antagonists, butyrophenones, phenothiazines, metoclopramide) OR Patient has a diagnosis of anorexia associated with weight loss due to AIDS.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	Part B if related to cancer treatment and is a full replacement for IV antiemetic within 48 hrs of cancer treatment. Part D if related to cancer treatment after the 48-hour period, or for any other medically accepted diagnosis.

# MEKINIST (S)

## Products Affected

- Mekinist

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of unresectable or metastatic melanoma and medication is used as a single agent and patient has a positive BRAF V600E or V600K mutation as detected by an FDA-approved test (THxID-BRAF Kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility, and the patient has not received prior BRAF-inhibitor therapy OR medication will be used in combination with Tafenlar in a patient with BRAF V600E or V600K mutations, as detected by an FDA-approved test (THxID-BRAF kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## MODAFINIL (S)

### Products Affected

- Modafinil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: A) excessive sleepiness associated with obstructive sleep apnea (OSA)/hypopnea syndrome confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and documentation of residual excessive sleepiness B) excessive sleepiness associated with narcolepsy confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and patient has tried and failed, is unable to tolerate, or has contraindication(s) to at least one other central nervous system stimulant (e.g., methylphenidate, mixed amphetamine salts, dextroamphetamine) OR Diagnosis of excessive sleepiness associated with shift work disorder confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and sleep disturbance causes measurable functional impairment in social, occupational, or other important areas of functioning that has persisted for at least three months.
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	OSA/hypopnea syndrome - 6 months (initial), 12 months (renewal). Other diagnoses - 12 months.
<b>Other Criteria</b>	N/A

# MOZOBIL (S)

---

## Products Affected

- Mozobil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient is to undergo autologous stem cell transplantation for the treatment of non-Hodgkin's lymphoma or multiple myeloma AND Patient will concomitantly receive a daily dose of a granulocyte colony-stimulating factor (G-CSF) for 4 days prior to the first evening dose of Mozobil and on each day prior to apheresis while using Mozobil.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	4 days
<b>Other Criteria</b>	N/A



# MYALEPT (S)

---

## Products Affected

- Myalept

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of congenital or acquired generalized lipodystrophy AND patient has one or more of the following metabolic abnormalities: insulin resistance (defined as requiring more than 200 units per day), hypertriglyceridemia, or diabetes AND patient is refractory to current standards of care for lipid and diabetic management AND the prescriber is registered in the MYALEPT REMS program
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with an endocrinologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, the patient has met one of the following: sustained reduction in hemoglobin A1C level from baseline or sustained reduction in triglyceride levels from baseline.

## MYOZYME (S)

---

### Products Affected

- Myozyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of infantile-onset Pompe disease (GAA) deficiency.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# NAGLAZYME (S)

---

## Products Affected

- Naglazyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Mucopolysaccharidosis VI (MPS VI or Maroteaux-Lamy syndrome).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## NEULASTA (S)

### Products Affected

- Neulasta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	For use as primary prophylaxis of febrile neutropenia (FN) in one of the following patients: A) Patient has a 20% or higher risk of FN based on chemotherapy regimen OR B) Patient has 10% to less than 20% risk of developing FN based on chemotherapy regimen AND at least one of the following risk factors are present: 65 years or older, Poor performance status, Poor nutritional status, Previous episodes of febrile neutropenia, Extensive prior treatment including large radiation ports, Cytopenias due to bone marrow involvement by tumor, Administration of combined chemoradiotherapy, Presence of open wounds or active infections, Other serious comorbidities (including renal or liver dysfunction notably elevated bilirubin), or C) patient has less than 10% risk of developing FN based on chemotherapy regimen AND the intent of treatment is curative or adjuvant and patient is at risk for serious medical consequences of FN, including death and patient is receiving myelosuppressive chemotherapy regimen for a non-myeloid malignancy, or D) For use as secondary prophylaxis of FN in a patient who had a neutropenic complication from a prior cycle of chemotherapy (for which primary prophylaxis was not received)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## NEUMEGA (S)

---

### Products Affected

- Neumega

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Patient has undergone myeloablative chemotherapy.
<b>Required Medical Information</b>	Diagnosis of non-myeloid malignancy and patient has undergone myelosuppressive chemotherapy (placing them at high risk for thrombocytopenia)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# NEUPOGEN (S)

## Products Affected

- Neupogen

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	N/A
Required Medical Information	<p>A) congenital, cyclic, or idiopathic neutropenia, B) Adjunct therapy (tx) for severe febrile neutropenia (FN) and receiving myelosuppressive tx for non-myeloid malignancy with any one: Received prophylactic CSF (not Neulasta) OR Did not receive prophylactic CSF and Use as adjunct to antibiotics in high-risk pt and any one: 65 years or older, Pneumonia, Hypotension and multiorgan dysfunction (sepsis syndrome), Invasive fungal infection or clinically-documented infection, Hospitalized when developed fever, Prior FN, Severe (ANC less than 100/mcL) or anticipated prolonged (more than 10 days) neutropenia, C) Autologous peripheral-blood progenitor cell transplant for collection by leukapheresis, D) Myeloablative chemotx for non-myeloid malignancy followed by BMT, E) Acute myeloid leukemia after completing induction/consolidation chemotx, F) Acute lymphoblastic leukemia after completing first few days of chemotx of initial induction or first post-remission course, G) Myelodysplastic syndrome with severe neutropenia and recurrent infection, H) Receiving radiation tx, not chemotx, and expect prolonged tx delays due to neutropenia, I) Neutropenia due to HIV infection and antiretroviral tx, J) Aplastic anemia, K) Primary prophylaxis of FN in one of the following: at least 20% FN risk based on chemotx OR 10% up to 20% risk with one of the following: 65 years or older, Poor performance status, Poor nutritional status, Previous FN, Extensive prior tx with large radiation ports, Cytopenias due to bone marrow involvement by tumor, Combined chemoradiotx, Open wounds or active infections, Other serious comorbidities (e.g., renal or liver dysfunction) OR less than 10% risk and intent is curative or adjuvant and risk for serious medical consequences, including death L) Receiving myelosuppressive chemotx for non-myeloid malignancy, M) Secondary prophylaxis of FN in pt with neutropenic complication from prior chemotx cycle (where primary prophylaxis was not received)</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A

<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## NEXAVAR (S)

---

### Products Affected

- Nexavar

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	squamous cell lung cancer being treated with carboplatin and paclitaxel.
<b>Required Medical Information</b>	Diagnosis of one of the following: A) unresectable hepatocellular carcinoma, B) Advanced renal cell carcinoma C) locally recurrent or metastatic, progressive, differentiated thyroid carcinoma refractory to radioactive iodine treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



# NORTHERA (S)

## Products Affected

- Northera

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of neurogenic orthostatic hypertension (NOH) AND NOH is due to one of the following: primary autonomic failure (Parkinson's disease, multiple system atrophy, and pure autonomic failure), dopamine beta-hydroxylase deficiency, non-diabetic autonomic neuropathy AND Patient has symptoms of NOH: Orthostatic dizziness, Lightheadedness, "feeling that you are about to black out" AND Patient has tried and had an inadequate response, contraindication or intolerance to midodrine
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	For renewal, Patient does not have persistent or sustained supine hypertension (SBP more than 180 mmHg or DBP more than 110 mmHg), Patient does not have persistent or sustained standing or sitting hypertension (SBP more than 180 mmHg or DBP more than 110 mmHg), and Patient had improvement in symptoms of NOH. Sustained mean elevated blood pressure that persists for longer than 5 minutes after change in position. Persistent means elevated BP that occurs on more than one occasion on separate physician office visits

## NOXAFIL (S)

### Products Affected

- Noxafil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Concomitant treatment with sirolimus, CYP 3A4 substrates (pimozide, quinidine), HMG-CoA Reductase inhibitors primarily metabolized through CYP 3A4, or ergot alkaloids
<b>Required Medical Information</b>	Diagnosis of oropharyngeal candidiasis and patient tried and failed itraconazole and/or fluconazole OR Medication will be used as prophylaxis of invasive Aspergillus and Candida infections and the patient is at high risk of developing these infections due to being severely immunocompromised, such as hematopoietic stem cell transplant (HSCT) recipients with graft-versus-host disease (GVHD) or those with hematologic malignancies with prolonged neutropenia from chemotherapy.
<b>Age Restrictions</b>	13 years of age or older for prophylaxis of invasive Aspergillus or Candidate infection
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 weeks
<b>Other Criteria</b>	N/A

## NPLATE (S)

### Products Affected

- Nplate INJ 250MCG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsed/refractory chronic immune (idiopathic) thrombocytopenic purpura (ITP) for greater than 6 months AND Baseline platelet count is less than 50,000/mcL AND Degree of thrombocytopenia and clinical condition increase the risk of bleeding AND Patient had an insufficient response, intolerance, contraindication to corticosteroids or immune globulin or inadequate response or contraindication to splenectomy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, after at least 4 weeks of therapy at the maximum weekly dose (10 mcg/kg) the platelet count increased to a sufficient level to avoid clinically important bleeding.

# NUVIGIL (S)

## Products Affected

- Nuvigil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: A) excessive sleepiness associated with obstructive sleep apnea (OSA)/hypopnea syndrome confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and documentation of residual excessive sleepiness B) excessive sleepiness associated with narcolepsy confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and patient has tried and failed, is unable to tolerate, or has contraindication(s) to at least one other central nervous system stimulant (e.g., methylphenidate, mixed amphetamine salts, dextroamphetamine) OR Diagnosis of excessive sleepiness associated with shift work disorder confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and sleep disturbance causes measurable functional impairment in social, occupational, or other important areas of functioning that has persisted for at least three months
<b>Age Restrictions</b>	17 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	OSA/hypopnea syndrome - 6 months (initial), 12 months (renewal). Other diagnoses - 12 months.
<b>Other Criteria</b>	N/A

## OFEV (S)

### Products Affected

- Ofev

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The patient has a diagnosis of idiopathic pulmonary fibrosis confirmed by a high resolution CT scan or biopsy AND the patient does not have evidence or suspicion of an alternative interstitial lung disease diagnosis AND liver function tests have been performed prior to start of therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescribed by or in consultation with a pulmonologist
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient experienced stabilization from baseline or a less than 10 percent decline in force vital capacity AND the patient has not experienced AST or ALT elevations greater than 5 times the upper limit of normal or greater than 3 times the upper limit of normal with signs or symptoms of severe liver damage.

# OLYSIO (BCBS RI)

## Products Affected

- Olysio

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has not previously failed a treatment regimen with a hepatitis C protease inhibitor AND Diagnosis of chronic hepatitis C genotype 1 virus infection with compensated liver disease AND A) medication will be used with ribavirin and peginterferon alfa AND patients with genotype 1a HCV infection have been screened for NS3 Q80K polymorphism and the patient does not have the polymorphism or B) medication will be used with Sovaldi with or without ribavirin AND the patient is treatment-naïve AND the patient is ineligible (e.g., labeled contraindication to peginterferon or patient has a concomitant condition that precludes the use of peginterferon) for an interferon-based regimen AND patient has not received prior treatment that included a protease inhibitor.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Medication is prescribed or in consultation with gastroenterologist, hepatologist or infection disease specialist
<b>Coverage Duration</b>	Initial -12 weeks. Renewal - 12 weeks
<b>Other Criteria</b>	N/A

# OPSUMIT (S)

---

## Products Affected

- Opsumit

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnancy.
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO group I with New York Heart Association Functional Class II or III AND diagnosis was confirmed by right heart catheterization AND female patients are enrolled in the OPSUMIT REMS program
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	For renewal, medication was effective (i.e. improved 6 minute walk distance, oxygen saturation, etc.)

# ORENCIA (BCBS RI)

## Products Affected

- Orencia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis)
<b>Required Medical Information</b>	Diagnosis of moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months AND has tried, failed or has a contraindication to Enbrel or Humira OR Diagnosis of moderate to severe polyarticular juvenile idiopathic arthritis and patient had an inadequate response, intolerance or contraindication to one or more non-biologic DMARDs for at least 3 consecutive months and patient had an inadequate response to one or more tumor necrosis factor inhibitors Enbrel or Humira
<b>Age Restrictions</b>	6 years of age or older for JIA. 18 years of age or older for rheumatoid arthritis
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. Dosing as per FDA approved labeling. For renewal, patient has stable disease or has improved while on therapy (e.g., for pJIA, reduction in disease flares, improvement in ACR scoring. For RA, improvement in tender/swollen joint count, improvement in ACR scoring.)



# ORENITRAM (S)

---

## Products Affected

- Orenitram

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	severe hepatic impairment (Child Pugh Class C)
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class III or III that was confirmed by right heart catheterization
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	N/A

## OTEZLA (S)

---

### Products Affected

- Otezla

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of psoriatic arthritis AND patient has tried and had an inadequate response, contraindication, or intolerance to methotrexate
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has stable disease or has improved on therapy, such as improvement in number of swollen/tender joints, pain, or stiffness.

# OXANDRIN (S)

## Products Affected

- Oxandrolone ORAL TABS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Breast or prostate cancer in men. Breast cancer in women with hypercalcemia. Pregnancy. Nephrosis or nephrotic phase of nephritis. Hypercalcemia.
<b>Required Medical Information</b>	Patient is receiving treatment as an adjunct therapy to promote weight gain and has one of the following: Extensive surgery, Chronic infections, Severe trauma, Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons and Patient has had an inadequate response, intolerance, or contraindication to nutritional supplements and a nutritional consult was performed OR Oxandrin (oxandrolone) will be used to counterbalance protein catabolism associated with chronic corticosteroid administration OR Patient has bone pain associated with osteoporosis.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Osteoporosis bone pain: 1 month. Other diagnoses: 3 months
<b>Other Criteria</b>	For renewal, patient has experienced an objective improvement (i.e. weight gain, increase in lead body mass, or reduction in muscle pain/weakness)

# OXYCONTIN (S)

## Products Affected

- Oxycodone Hcl Er
- Oxycontin

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	Significant respiratory depression. Acute or severe bronchial asthma. Known or suspected paralytic ileus. GI obstruction
<b>Required Medical Information</b>	Diagnosis of severe pain requiring continuous, around-the-clock opioid analgesic for an extended period of time AND patient has tried and failed or unable to tolerate at least two generic extended-release opioid products, such as: oxymorphone ER, morphine ER, fentanyl, methadone, tramadol ER AND Patient is opioid tolerant (for 60-80mg only), taking at least 60 mg oral morphine per day, 25 µg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, or an equianalgesic dose of another opioid for a week or longer
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

## PEGINTRON (S)

### Products Affected

- Pegintron
- Peg-intron
- Peg-intron Redipen
- Peg-intron Redipen Pak 4

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Uncontrolled depression. Autoimmune hepatitis or other autoimmune condition known to be exacerbated by interferon and ribavirin.
<b>Required Medical Information</b>	Diagnosis of hepatitis C with compensated liver disease and patient will be receiving one of the following: A) Combination therapy with an HCV protease inhibitor and ribavirin and Patient has HCV genotype 1 OR B) Combination therapy with ribavirin only and if patient has HCV genotype 1, the patient has a contraindication to HCV protease inhibitors OR C) Monotherapy when ribavirin is contraindicated OR D) Combination with Sovaldi and ribavirin for genotype 1 or 4 infection.
<b>Age Restrictions</b>	18 yrs or older if triple therapy (except with Sovaldi or Olysio), else 3 yrs or older.
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Sovaldi, Olysio- 12 wks. Others-16 wks (initial) Renewal- based on FDA label.
<b>Other Criteria</b>	For renewal, approval is based on the requirements outlined in the FDA-approved labeling, including viral load, combination therapy, and response to prior therapy.

## PENNSAID (S)

---

### Products Affected

- Pennsaid SOLN 2%

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	Previously experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Use for the treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery.
<b>Required Medical Information</b>	Patient has a diagnosis of osteoarthritis of the knees AND patient had experienced treatment failure with at least 2 prescription strength oral NSAIDs or patient has a documented swallowing disorder OR has a history of peptic ulcer disease/gastrointestinal bleeding OR patient is more than 65 years of age with one additional risk factor for gastrointestinal adverse event (e.g., use of anticoagulants or chronic corticosteroids)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## PERJETA (BCBS RI)

---

### Products Affected

- Perjeta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of HER2-positive metastatic breast cancer AND medication is being used in combination with trastuzumab and docetaxel AND Patient has not received prior anti-HER2 therapy (e.g., trastuzumab) or chemotherapy for metastatic disease AND pregnancy status will be verified prior to initiation of therapy AND females of reproductive potential will be advised of the risks of embryo-fetal death and birth defects, and the need for effective contraception during and after pertuzumab treatment.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

## PLEGRIDY (S)

---

### Products Affected

- Plegridy

- Plegridy Starter Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)



## POMALYST (S)

---

### Products Affected

- Pomalyst

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of multiple myeloma and the patient has received two prior therapies, including Revlimid and Velcade unless the patient has a contraindication or intolerance to Revlimid or Velcade and the patient has demonstrated disease progression on or within 60 days of completion of last therapy AND the prescriber is registered and patient is enrolled in the Pomalyst REMS program
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 month
<b>Other Criteria</b>	N/A

## PROCYSBI (S)

---

### Products Affected

- Procysbi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of nephropathic cystinosis AND Patient has tried and failed, or had an intolerance to, therapy with Cystagon (immediate-release cysteamine bitartrate)
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# PROLIA (BCBS RI)

## Products Affected

- Prolia

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	N/A
Required Medical Information	<p>Patient is at high risk for fracture defined as one of the following: Personal history of low-trauma fractures as an adult, History of osteoporotic fracture in a first degree relative, Concurrent use of systemic corticosteroids (avg dose more than 5 mg of prednisone per day), Concurrent cigarette smoking, Low body weight less than 127 pounds, Low bone mineral density (T-score of -2.5 or lower) AND Diagnosis of one of the following: Patient is female and is receiving adjuvant aromatase inhibitor therapy for breast cancer, Patient is male and is receiving androgen deprivation therapy for non-metastatic prostate cancer, Patient is a male or postmenopausal female with a diagnosis of osteoporosis AND Patient has a documented trial and failure with a bisphosphonate (failure is defined as new fractures in compliant patients) or contraindication or intolerance to bisphosphonate therapy or is unable to comply with appropriate administration recommendations for oral or injectable bisphosphonate therapy AND Patient is concomitantly taking calcium 1000 mg daily and at least 400 IU vitamin D daily AND Patient will have pre-existing hypocalcemia and vitamin D deficiency corrected prior to administration of the medication</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	Calcium, mineral, and vitamin D levels will be monitored periodically during treatment. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

# PROMACTA (S)

## Products Affected

- Promacta

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: A) Relapsed/refractory chronic immune (idiopathic) thrombocytopenic purpura (ITP) for greater than 6 months AND Baseline platelet count is less than 50,000/mcL AND Degree of thrombocytopenia and clinical condition increase the risk of bleeding AND Patient had an insufficient response, intolerance, contraindication to corticosteroids or immune globulin or inadequate response or contraindication to splenectomy, B)Chronic hepatitis C and patient has thrombocytopenia defined as platelets less than 90,000/mcL for initiation (pre-treatment) of interferon therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	ITP - 12 months. Hep C - 9 weeks (initial), 24 weeks (renewal)
<b>Other Criteria</b>	For renewal of ITP, after at least 4 weeks of therapy at the maximum weekly dose (10 mcg/kg) the platelet count increased to a sufficient level to avoid clinically important bleeding. For renewal of Hepatitis C, platelets less than 75,000/mcL for maintenance of optimal interferon-based therapy.

# PULMOZYME (S)

---

## Products Affected

- Pulmozyme

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient is benefiting from treatment (i.e. improvement in lung function [FEV1], decreased number of pulmonary exacerbations). Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## QUALAQUIN (S)

---

### Products Affected

- Quinine Sulfate CAPS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Prolongation of QT interval. Glucose-6-phosphate dehydrogenase deficiency. Myasthenia gravis. Known hypersensitivity to mefloquine or quinidine. Optic neuritis.
<b>Required Medical Information</b>	Patient is being treated for uncomplicated Plasmodium falciparum malaria
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	One month
<b>Other Criteria</b>	Dosing will be approved per the FDA labeling.

## RAVICTI (BCBS RI)

---

### Products Affected

- Ravicti

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	acute hyperammonemia. N-acetylglutamate synthase (NAGS) deficiency
<b>Required Medical Information</b>	Diagnosis of urea cycle disorder involving deficiencies of carbamoyl phosphate synthetase (CPS), ornithine transcarbamoylase (OTC), or argininosuccinic acid synthetase (AAS) confirmed via enzymatic, biochemical, or genetic testing AND protein-restricted diet alone or amino acid supplements alone has been ineffective AND patient has tried and had an inadequate response, is intolerant, or has a contraindication to Buphenyl for at least 3 months AND patient will maintain a protein-restricted diet while on therapy.
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## REBIF (S)

---

### Products Affected

- Rebif

- Rebif Titration Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. First clinical episode with MRI features consistent with multiple sclerosis.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsing form of multiple sclerosis OR diagnosis of first clinical episode and MRI features consistent with multiple sclerosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has experienced an objective response to therapy (i.e. no or slowed progression of disease)



## RECLAST (BCBS RI)

### Products Affected

- Zoledronic Acid INJ 5MG/100ML

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indication not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Creatinine clearance less than 35 mL/min or evidence of acute renal impairment. Hypocalcemia (corrected calcium level less than 8.0 mg/dL). Patient is currently receiving Zometa.
<b>Required Medical Information</b>	Supplementation with calcium and vitamin D AND Diagnosis of one of the following: A) Postmenopausal female with osteoporosis diagnosed by BMD (T-score -2.5 or below) or by history of fracture, B) Patient will receive medication for prevention of osteoporosis AND patient is a postmenopausal female with one of the following risk factors for fracture: Low BMD (T-score -1.5 or below), Low body weight (less than 57.6 kg or 127 pounds), Smoking, Alcohol use (3 or more drinks/day), Family history of osteoporosis, Early menopause, Age 65 years or older, Equivalent dose of 7.5 mg prednisone or more/day for a minimum of 3 months, C) Male with osteoporosis diagnosed by BMD (T-score -2.5 or below) or by history of fracture, D) Patient is male and will receive medication for prevention of osteoporosis AND patient requires an equivalent dose of 7.5 mg prednisone or more/day for a minimum of 3 months, E) Premenopausal female with a history of fracture AND Patient requires the use of glucocorticoids for at least 3 months or requires an equivalent dose of 7.5 mg prednisone or more/day, F) Premenopausal female who requires an equivalent dose of 7.5 mg prednisone or more/day for a minimum of 12 months G) Diagnosis of Paget's disease of bone with any one of the following: Serum alkaline phosphatase two times or more higher than the upper limit of the age-specific normal reference range, Symptomatic disease (i.e. bone pain, headache with skull involvement, back pain due to radiculopathy or arthropathy, fissure fractures), At risk for complications from the disease (e.g., those with active disease near neurovascular structures or major joints) AND For osteoporosis indications, Patient has tried and had inadequate response to oral bisphosphonates, unless patient has a contraindication/intolerance to oral bisphosphonate.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A

<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve doses based on FDA labeling. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.

# RELISTOR (S)

## Products Affected

- Relistor INJ 12MG/0.6ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Known or suspected mechanical gastrointestinal obstruction. On renewal, patient does not have severe or persistent diarrhea.
<b>Required Medical Information</b>	Diagnosis of opioid-induced constipation AND Patient has used opioid medication for a minimum of 2 weeks AND Patient is experiencing fewer than 3 bowel movements in a week or no bowel movement for longer than 2 days AND Patient is diagnosed with an advanced illness (e.g., incurable cancer, end-stage chronic obstructive pulmonary disease/emphysema, cardiovascular disease/heart failure, Alzheimer's disease/dementia, HIV/AIDS, etc.) AND Patient is receiving palliative care AND Patient has tried and had an insufficient response to laxative (e.g., lubiprostone) therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	4 months
<b>Other Criteria</b>	For renewal, the patient has responded to therapy (i.e. increase in bowel movements)

# REMICADE (BCBS RI)

## Products Affected

- Remicade

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis). Moderate to severe heart failure in patients receiving doses greater than 5 mg/kg.
<b>Required Medical Information</b>	Diagnosis of moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months and patient will be on concomitant methotrexate OR Diagnosis of ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to one or more NSAIDs OR Diagnosis of severe chronic plaque psoriasis (affecting more than 10% of body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) and patient had an inadequate response, intolerance or contraindication to conventional therapy with at least one of the following: phototherapy (including but not limited to Ultraviolet A with a psoralen [PUVA] and/or retinoids [RePUVA] for at least one continuous month or one or more oral systemic treatments for at least 3 consecutive months OR Diagnosis of moderate to severe Crohn's disease and patient had an inadequate response, intolerance, or contraindication to conventional therapy with two or more of the following: corticosteroids or non-biologic DMARDs OR Diagnosis of fistulizing Crohn's disease OR Diagnosis of moderate to severe ulcerative colitis and patient had an inadequate response, intolerance or contraindication to conventional therapy with two or more of the following: corticosteroids, 5-ASA (i.e. mesalamine, sulfasalazine, balsalazide, olsalazine) or non-biologic DMARDs OR Diagnosis of psoriatic arthritis and patient had an inadequate response, intolerance or contraindication to methotrexate.
<b>Age Restrictions</b>	6 years of age or older for UC or Crohn's disease (non-fistulizing). 18 years of age or older for all other indications, including fistulizing Crohn's disease
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months

<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. Approvable under Part D only if patient is in a long term care facility OR the medication is not being administered with an infusion pump.
-----------------------	---

## REMODULIN (S)

---

### Products Affected

- Remodulin

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class II-IV that was confirmed by right heart catheterization
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## REVATIO (S)

---

### Products Affected

- Revatio INJ
- Revatio SUSR
- Sildenafil

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Receiving nitrate therapy (includes intermittent use)
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class II or III that was confirmed by right heart catheterization
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	N/A

# REVLIMID (S)

## Products Affected

- Revlimid

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of multiple myeloma and patient has received at least one prior therapy and medication will be used in combination with dexamethasone OR diagnosis of transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes associated with a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities OR Diagnosis of mantle cell lymphoma and patient's disease has relapsed or progressed after trying at least two prior therapies including Velcade AND patient is enrolled in the Revlimid REMS Program AND patient is not using the medication for the treatment of chronic lymphocytic leukemia
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## RUCONEST (BCBS RI)

---

### Products Affected

- Ruconest

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	N/A
<b>Other Criteria</b>	N/A

## SABRIL (S)

### Products Affected

- Sabril

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: A) infantile spasms B) complex partial seizures and patient had an inadequate response to at least one generic first-line agents (carbamazepine, lamotrigine, levetiracetam, oxcarbazepine, valproic acid, divalproex sodium) and at least one adjunctive agent (carbamazepine, clobazam, gabapentin, lamotrigine, levetiracetam, oxcarbazepine, valproic acid, divalproex sodium, topiramate) AND patient and prescriber are enrolled in the SHARE restricted distribution program.
<b>Age Restrictions</b>	seizures - 10 years of age or older. Infantile spasms - at least one month to 2 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# SANDOSTATIN (S)

## Products Affected

- Octreotide Acetate

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses OR Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes OR Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

## SANDOSTATIN LAR (S)

---

### Products Affected

- Sandostatin Lar Depot

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of acromegaly and patient had an inadequate response or cannot be treated with surgical resection, pituitary irradiation, and/or bromocriptine mesylate at maximally tolerated doses OR Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea and flushing episodes OR Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea AND Patient received at least two weeks of Sandostatin Injection and has tolerate therapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal of acromegaly, IGF-1 level has normalized or improved. For renewal of metastatic carcinoid tumor, patient has improvement in diarrhea and flushing episodes. For renewal of vasoactive intestinal peptide tumor, improvement in diarrhea episodes.

## SAVELLA (S)

---

### Products Affected

- Savella

- Savella Titration Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Use of monoamine oxidase inhibitors concomitantly or within 14 days. Uncontrolled narrow-angle glaucoma.
<b>Required Medical Information</b>	Diagnosis of fibromyalgia AND patient had a previous trial (of at least 30 days) with or has a contraindication, intolerance, or allergy to one of the following agents used for the treatment of fibromyalgia: tricyclic antidepressant, SNRI, SSRI, gabapentin.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal patient had an improvement in pain, physical functioning, etc.

## SEROSTIM (BCBS RI)

---

### Products Affected

- Serostim

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Acute critical illness due to complications following open heart or abdominal surgery, multiple accidental trauma or acute respiratory failure. Active malignancy. Active proliferative or severe non-proliferative diabetic retinopathy. Active malignancy or undergoing cancer treatment
<b>Required Medical Information</b>	Diagnosis of AIDS-wasting syndrome or cachexia (defined as unintentional weight loss of at least 10% of baseline weight) AND Treatment failure with or intolerance to dronabinol or megestrol AND Patient is currently receiving treatment with antiretrovirals
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 weeks
<b>Other Criteria</b>	For renewal, patient has experienced an increase in body weight and/or improvement in lead body mass AND wasting is still evident

## SIGNIFOR (S)

---

### Products Affected

- Signifor

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of (pituitary) Cushing's disease AND pituitary surgery is not an option or has not been curative
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient had a clinically meaningful reduction in 24-hour urinary free cortisol levels and/or improvement in signs or symptoms of the disease

# SIMPONI (S)

## Products Affected

- Simponi

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis)
<b>Required Medical Information</b>	Diagnosis of one of the following: A) moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months and patient will be on concomitant methotrexate B) psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate C) ankylosing spondylitis and patient had an inadequate response, intolerance or contraindication to one or more NSAIDs D) moderately to severely active ulcerative colitis and patient has had inadequate responses to, is intolerant to, or is contraindicated to conventional therapy with two or more of the following: Corticosteroids (i.e. prednisone, methylprednisolone), 5-ASAs (i.e. mesalamine, sulfasalazine, balsalazide, olsalazine), or Non-biologic DMARDs (i.e. azathioprine, methotrexate, mercaptopurine)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial 10 weeks (UC), 12 months others. Renewal 12 months.
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. For renewal, patient has stable disease or has improved while on therapy (e.g., for RA, improvement in tender/swollen joint count, improvement in ACR scoring. For PsA, improvement in number of swollen/tender joints, pain, stiffness. For AS, improvement in AS symptoms, such as stiffness and back pain. For UC, clinical remission, reduction in steroid use.)



## SIMPONI ARIA (S)

---

### Products Affected

- Simponi Aria

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis)
<b>Required Medical Information</b>	Diagnosis of moderate to severe rheumatoid arthritis and patient had an inadequate response to, intolerance to, or contraindication to one or more non-biologic disease modifying anti-rheumatic drugs (DMARDs) for at least 3 consecutive months and patient will be on concomitant methotrexate
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months.
<b>Other Criteria</b>	Patient has been tested for TB and latent TB has been ruled out or is being treated. For renewal, patient has stable disease or has improved while on therapy (e.g., for RA, improvement in tender/swollen joint count, improvement in ACR scoring.)

# SIMVASTATIN (S)

## Products Affected

- Simvastatin TABS 80MG
- Vytorin TABS 10MG; 80MG

PA Criteria	Criteria Details
Covered Uses	All FDA-approved indications not otherwise excluded from Part D.
Exclusion Criteria	Active liver disease. Pregnancy. Nursing. Patient is taking or initiating therapy with any of the following: verapamil, diltiazem, amiodarone, dronedarone, amlodipine, ranolazine, strong CYP3A4 inhibitors (i.e., itraconazole, ketoconazole, posaconazole, protease inhibitors, erythromycin, clarithromycin, telithromycin, and nefazodone), gemfibrozil, cyclosporine, and danazol.
Required Medical Information	Patient has been taking simvastatin 80 mg chronically (12 months or more) without evidence of muscle toxicity AND, if patient is of Chinese descent, they are not concurrently receiving lipid-modifying doses (at least 1 gram/day) of niacin-containing products
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	12 months
Other Criteria	N/A

## SOMATULINE DEPOT (S)

---

### Products Affected

- Somatuline Depot

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of acromegaly AND Inadequate response to surgery and/or radiation therapy or patient is not a candidate for surgery and/or radiotherapy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient's IGF-1 levels for age and gender has normalized or improved.

## SOMAVERT (S)

---

### Products Affected

- Somavert

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of acromegaly AND Inadequate response to surgery and/or radiation therapy and/or other medical therapies (such as dopamine agonists) AND patient had an inadequate response or intolerance to generic octreotide OR patient is not a candidate for any of those treatment options.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient has experienced a decrease or normalization of insulin-like growth factor-1 (IGF-1) levels.

## SOVALDI (BCBS RI)

### Products Affected

- Sovaldi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis C AND A) medication will be used with ribavirin and peginterferon alfa and has genotype 1 or 4 infection, B) medication will be used with ribavirin and patient meets one of the following: a) Genotype 2 infection, b) Genotype 3 infection and patient is ineligible for an interferon-based regimen (e.g., labeled contraindication to peginterferon or specific condition that precludes the use of peginterferon), c) Genotype 1 infection and patient is ineligible for an interferon-based regimen, or d) Hepatocellular carcinoma and meets Milan criteria for liver transplant (i.e. a single HCC nodule with a maximum size of 5 cm or as many as 3 nodules with the largest not exceeding 3 cm and no macrovascular invasion) OR C) Medication will be used with simeprevir with or without ribavirin and the patient has genotype 1 infection and patient is treatment-naive and ineligible for an interferon-based regimen.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Medication is prescribed or in consultation with gastroenterologist, hepatologist or infection disease specialist
<b>Coverage Duration</b>	12 wk: G1 triple tx, G2, G4, w/ Olysio. 24 wk: G1 dual tx, G3. 16 wk: G2 non-responder. 48 wk: HCC
<b>Other Criteria</b>	N/A

## SPRYCEL (S)

### Products Affected

- Sprycel

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. GIST.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) in the chronic phase and patient is found to be Ph+ or BCR-ABL positive as detected by bone marrow cytogenetics or FISH OR Ph+ CML with resistance, relapse or inadequate response to prior TKI therapy (e.g., Gleevec, Tasigna, Iclusig, Bosulif) and if patient has mutation testing, patient does not have T315I mutation OR Ph+ CML with intolerance to prior therapy OR Diagnosis of Ph+ acute lymphoblastic leukemia OR Gastrointestinal stromal tumors (GIST) after disease progression on Gleevec (imatinib) or Sutent (sunitinib)
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# STELARA (BCBS RI)

---

## Products Affected

- Stelara

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis [TB]) BCG during and 1 year prior to treatment
<b>Required Medical Information</b>	Diagnosis of one of the following: A) moderate to severe plaque psoriasis (affecting more than 5% of the body surface area or affecting crucial body areas such as the hands, feet, face, or genitals) AND patient tried and had an inadequate response, is intolerant of, or is contraindicated to conventional therapy with at least one of the following: phototherapy (including but not limited to, psoralen with ultraviolet-a [PUVA] and/or retinoids [rePUVA]) for at least one continuous month or oral systemic treatment (e.g., methotrexate, cyclosporine, acitretin) for at least 3 consecutive months B) psoriatic arthritis and patient had an inadequate response, intolerance, or contraindication to methotrexate.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient has been tested for latent TB infection and latent TB has been ruled out or is being treated per guidelines. Dosing as per FDA approved labeling. For renewal, patient has stable disease or has improved while on therapy (e.g., for PsA, improvement in number of swollen/tender joints, pain, stiffness).

## STIVARGA (BCBS RI)

---

### Products Affected

- Stivarga

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of: A) metastatic colorectal cancer AND patient has previously treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based therapy, an anti-vascular endothelial growth factor (VEGF) therapy, and, if KRAS wild type, an anti-epidermal growth factor receptor (EGFR) therapy or B) gastrointestinal stromal tumors that is locally advanced, unresectable or metastatic and patient has tried and had an inadequate response, contraindication or intolerance to Gleevec or Sutent.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Hepatic function will be monitored prior to and during treatment and, if patient has elevated liver function tests of hepatocellular necrosis, therapy will be interrupted and then reduced or discontinued.



# SUBOXONE (S)

## Products Affected

- Buprenorphine Hcl/naloxone Hcl

- Suboxone SUBLINGUAL FILM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of opioid dependence AND Prescription is a part of an overall treatment program (e.g., self-help groups, counseling, provide ongoing care, vocational training) AND Patient is not receiving any other opioids written by a different prescriber AND patient is not pregnant
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	Prescriber is certified through SAMHSA (Substance Abuse and Mental Health Services Administration) to prescribe Suboxone and provide registration number
<b>Coverage Duration</b>	Initial - 3 months. Renewal - 9 months
<b>Other Criteria</b>	For renewal, patient meets all initial criteria and prescriber is evaluating random urine drug screens and assessing the patient's progress (e.g., relapse, progress/accomplishment of treatment goals) and patient is not pregnant.

## SUBUTEX (S)

---

### Products Affected

- Buprenorphine Hcl SUBLINGUAL SUBL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of opioid dependence AND Prescription is a part of an overall treatment program (e.g., self-help groups, counseling, provide ongoing care, vocational training) AND Patient is not receiving any other opioids since starting therapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber is certified through SAMHSA (Substance Abuse and Mental Health Services Administration) to prescribe Suboxone and provide registration number
<b>Coverage Duration</b>	Initial - 3 months (pregnant) or 1 month (not pregnant). Renewal - 9 months
<b>Other Criteria</b>	For renewal, patient meets all initial criteria and prescriber is evaluating random urine drug screens and assessing the patient's progress (e.g., relapse, progress/accomplishment of treatment goals)

## SUTENT (S)

---

### Products Affected

- Sutent

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of gastrointestinal stromal tumors after disease progression on or intolerance to Gleevec OR Diagnosis of progressive, well-differentiated pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## SYLATRON (S)

---

### Products Affected

- Sylatron

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	autoimmune hepatitis. Hepatic decompensation (Child-Pugh score greater than 6 [Class B or C])
<b>Required Medical Information</b>	Patient is being treated adjuvantly for melanoma with microscopic or gross nodal involvement within 84 days of definitive surgical resection including complete lymphadenectomy.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## SYLVANT (S)

---

### Products Affected

- Sylvant INJ 100MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Patient has a diagnosis of multicentric Castleman's disease AND patient is HIV negative AND patient is human herpes virus-8 (HHV-8) negative
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has not experienced treatment failure defined as disease progression based on increase in symptoms, radiologic progression, or deterioration in performance status

# SYNAGIS (S)

## Products Affected

- Synagis

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	<p>Patient will use palivizumab for immunoprophylaxis of respiratory syncytial virus (RSV) during the peak months of infection in the patient's geographic region AND Patient meets one of the following criteria: A) Infants born at 28 weeks, six days gestation or earlier and who are younger than 12 months of age at the start of the RSV season OR B) Infants born at 29 to 31 weeks, six days gestation and who are younger than six months of age at the start of the RSV season OR C) Infants born at 32 to 34 weeks, six days gestation and who are younger than three months of age at the start of RSV season with at least one of the following risk factors may be dosed until 90 days of age: Child care attendance or Sibling younger than five years of age living in the same household (who is not a multiple birth younger than one year of age) OR D) Infants and children younger than one year of age at the start of RSV season with either congenital abnormalities of the airway or neuromuscular disease that compromises handling of respiratory secretions OR E) Infants and children younger than two years of age with hemodynamically significant congenital heart disease and who have at least one of the following criteria: Receiving medication to control congestive heart failure, Has moderate to severe pulmonary hypertension, or Has cyanotic heart disease OR F) Infants and children younger than two years of age who have received medical therapy (oxygen, bronchodilator, diuretic, or corticosteroid therapy) for chronic lung disease within six months of the start of the RSV season</p>
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Approve 5 doses based on patient body weight.

## SYNRIBO (S)

---

### Products Affected

- Synribo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of chronic myelogenous leukemia AND patient has tried and failed or has a contraindication or intolerance to 2 tyrosine kinase inhibitors
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## TACLONEX (BCBS RI)

---

### Products Affected

- Calcipotriene/betamethasone Dipropionate

- Taclonex SUSP

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of stable psoriasis vulgaris (plaque psoriasis) AND Patient tried adequate therapy (at least two weeks) with at least one of the following agents: medium to high potency topical steroid (unless contraindicated/intolerant without concurrent vitamin D analog use) or vitamin D analogs (unless contraindicated/intolerant without concurrent steroid use) or tazarotene (unless contraindicated/intolerant to its use)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Suspension: 8 weeks, Ointment: 4 weeks
<b>Other Criteria</b>	For renewal, has the patient responded to therapy (e.g., patient's symptoms have improved)



# TAFINLAR (S)

## Products Affected

- Tafinlar

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of unresectable or metastatic melanoma and medication will be used as a single agent in a patient with a positive BRAF V600E mutation as detected by an FDA-approved test (THxID-BRAF Kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility OR medication will be used in combination with Mekinist in a patient with BRAF V600E or V600K mutation, as detected by an FDA-approved test (THxID-BRAF kit) or Clinical Laboratory Improvement Amendments (CLIA)-approved facility.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# TARCEVA (S)

## Products Affected

- Tarceva

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of locally advanced, unresectable, or metastatic pancreatic cancer and Tarceva will be used in combination with gemcitabine OR Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer with one of the following: A) failure with at least one prior chemotherapy regimen and Tarceva will be used as monotherapy, or B) no evidence of disease progression after four cycles of first-line platinum-based chemotherapy and Tarceva will be used as maintenance treatment and Tarceva will be used as monotherapy, or C) Patient has known active epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# TARGRETIN (S)

---

## Products Affected

- Targretin

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Female patients of child-bearing potential have a documented negative pregnancy test one week prior to the initiation of therapy. For renewal, Patient has not had disease progression while on therapy and female patients of child-bearing potential are not pregnant and are continuing to use adequate birth-control measures during therapy.

# TASIGNA (S)

## Products Affected

- Tasigna

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. GIST.
<b>Exclusion Criteria</b>	Long QT syndrome. Uncorrected hypokalemia. Uncorrected hypomagnesemia. Concomitant use with a drug known to prolong the QT interval or strong cytochrome P450 3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of one of the following: A) newly diagnosed Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML) in the chronic phase and patient is found to be Ph+ or BCR-ABL positive as detected by bone marrow cytogenetics or FISH OR Diagnosis of Ph+ CML with resistance, relapse or inadequate response to prior therapy and if the patient had mutation testing, patient does not have the T315I mutation OR Ph+ CML with intolerance to prior therapy B) Gastrointestinal stromal tumors after disease progression on Gleevec or Sutent
<b>Age Restrictions</b>	18 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# TAZORAC (S)

## Products Affected

- Tazorac

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indication not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnancy.
<b>Required Medical Information</b>	Diagnosis of acne vulgaris and patient has tried an adequate trial (at least two weeks) with at least one other topical acne product (e.g., benzoyl peroxide, salicylic acid, clindamycin, erythromycin, adapalene, azelaic acid, and/or tretinoin) OR Diagnosis of stable moderate to severe plaque psoriasis and 20% or less body surface area involvement and patient has a contraindication or tried adequate trial (at least 2 weeks) with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs) AND females of child-bearing potential are using adequate birth control measures during therapy.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# TECFIDERA (S)

---

## Products Affected

- Tecfidera

- Tecfidera Starter Pack

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of relapsing forms of multiple sclerosis (relapsing-remitting MS or progressive-relapsing MS, or secondary-progressive MS) OR patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, Patient had an objective response to therapy (ie no or slowed progression of disease)

## THALOMID (S)

---

### Products Affected

- Thalomid

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of multiple myeloma that is newly diagnosed and is receiving concurrent dexamethasone OR Diagnosis of severe erythema nodosum leprosum with cutaneous manifestations and the medication will not be used as monotherapy if the member has moderate to severe neuritis
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Prescriber is registered and the member is enrolled in the Thalomid REMS program
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## TOPICAL RETINOIDS (S)

### Products Affected

- Retin-a Micro
- Tretinoin EXTERNAL CREA
- Tretinoin EXTERNAL GEL
- Tretinoin Microsphere
- Tretinoin Microsphere Pump
- Tretin-x CREA 0.075%
- Tretin-x EXTERNAL KIT 0; 0; 0; 0; 0; 0.025%, 0; 0; 0; 0; 0; 0.05%, 0; 0; 0; 0; 0.1%
- Ziana

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Acne rosacea. Actinic keratoses (solar keratoses). Flat warts (verruca plana/juveniles). Hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans [Flegel's disease]). Ichthyosis.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of one of the following: A) mild to moderate acne vulgaris (including comedonal, cystic, and nodular) B) acne rosacea and patient has experienced treatment failure or intolerance to sulfacetamide/sulfur, metronidazole or azelaic acid C) actinic keratosis (or solar keratoses) with multiple lesions on the face C) flat warts (verruca plana/juveniles), hyperkeratosis disorder (i.e. epidermolytic hyperkeratosis, lenticularis perstans (Flegel's disease), palmoplantar hyperatosis, keratosis follicularis (Darier's disease)) F) Ichthyosis (i.e. congenital, lamellar, vulgaris, or X-linked)
<b>Age Restrictions</b>	PA applies to patients older than 26 years of age
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, medication has been effective in treating the patient's condition.



## TRACLEER (S)

---

### Products Affected

- Tracleer

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Receiving concomitant cyclosporine A or glyburide therapy.
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class II-IV that was confirmed by right heart catheterization AND patient has been enrolled into the TAP Restricted Distribution Program
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	N/A

## TRELSTAR (S)

---

### Products Affected

- Trelstar Depot Mixject
- Trelstar La Mixject
- Trelstar Mixject

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced or metastatic prostate cancer
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# TYKERB (S)

---

## Products Affected

- Tykerb

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND a) the medication will be used in combination with Xeloda in a patient with advanced or metastatic disease and the patient has received prior therapy including an anthracycline, a taxane, and trastuzumab or b) The medication will be used in combination with Femara for the treatment of a postmenopausal woman with hormone receptor-positive metastatic disease for whom hormonal therapy is indicated.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient's hepatic function has been assessed prior to starting therapy.

# TYSABRI (S)

## Products Affected

- Tysabri

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	History of progressive multifocal leukoencephalopathy.
<b>Required Medical Information</b>	Diagnosis of relapsing form of multiple sclerosis and medication will be used as monotherapy and patient had an inadequate response, intolerance, or contraindication to one of the following: An interferon beta product, Copaxone, Gilenya, Aubagio, or Tecfidera OR Diagnosis of moderate to severe active Crohn's disease and medication will not be used in combination with immunosuppressants or inhibitors of tumor necrosis factor-alfa and patient had an inadequate response, intolerance, or contraindication to any of the following: Humira, Remicade, or Cimzia.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient and physician are registered in the TOUCH prescribing program.

# TYVASO (S)

---

## Products Affected

- Tyvaso
- Tyvaso Refill
- Tyvaso Starter

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class III that was confirmed by right heart catheterization
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## TYZEKA (S)

### Products Affected

- Tyzeka

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Co-administration with pegylated interferon alfa-2a
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis B AND patient is HBsAg-positive for at least 6 months AND For HBeAg-positive patients, serum HBV DNA greater than 20,000 IU/mL (105 copies per mL) and for HBeAg-negative patients, serum HBV DNA greater than 2,000 IU/mL (104 copies/mL) AND Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) at least 2 times the upper limit of normal or histologically active disease (i.e. necroinflammation on biopsy)
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient must be HBeAg negative and have not had HBsAg clearance OR HBeAg positive and have detectable HBV DNA and have not been anti-Hbe for at least 6 months

# UNIVERSAL CRITERIA

---

## Products Affected

- Blincyto
- Entyvio
- Lynparza
- Opdivo

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	N/A
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	N/A
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	N/A
<b>Other Criteria</b>	N/A

## VALCHLOR (S)

---

### Products Affected

- Valchlor

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of mycosis fungoides-type cutaneous T-cell lymphoma AND patient has early stage disease (defined as Stage 1A or 1B) AND patient has received prior skin-directed therapy (e.g., very high potency class I topical corticosteroids for at least 3 months (i.e. clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate), phototherapy, topical nitrogen mustard, or a topical retinoid (e.g., bexarotene)).
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



## VELETRI (S)

---

### Products Affected

- Epoprostenol Sodium

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Congestive heart failure due to severe left ventricular systolic dysfunction. Pulmonary edema during dose initiation.
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension WHO Group I with New York Heart Association (NYHA) functional class III or IV that was confirmed by right heart catheterization
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	6 months - initial. 12 months - renewal
<b>Other Criteria</b>	Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## VENTAVIS (BCBS RI)

---

### Products Affected

- Ventavis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Congestive heart failure due to severe left ventricular systolic dysfunction.
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension that was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) AND Patient has WHO Group I PAH AND Patient has New York Heart Association (NYHA) Functional Class III or IV.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Initial - 6 months. Renewal - 12 months
<b>Other Criteria</b>	Part D if patient in long term care (defined by customer location code on claim) otherwise Part B.

## VICTRELIS(BCBS RI)

---

### Products Affected

- Victrelis

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Pregnant. Unwilling to comply with required contraception methods. Co-administration with alfuzosin, carbamazepine, cisapride, dihydroergotamine, drospirinone, ergonovine, ergotamine, lovastatin, methylergonovine, midazolam (oral), phenobarbital, phenytoin, pimozone, rifampin, sildenafil (Revatio), simvastatin, St. John's wort, tadalafil (Adcirca), triazolam.
<b>Required Medical Information</b>	Diagnosis of chronic hepatitis C genotype 1 with compensated liver disease AND medication will be used with ribavirin and peginterferon alfa AND Has not previously failed a treatment regimen with a hepatitis C protease inhibitor.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Written by or in consultation with a gastroenterologist, hepatologist or ID specialist
<b>Coverage Duration</b>	Initial - 12 weeks. Renewal - duration based on FDA approved labeling
<b>Other Criteria</b>	For renewal, approval is based on the requirements outlined in the FDA-approved labeling, including viral load, presence of cirrhosis, and response to prior therapy and confirm that the patient is continuing to receive concurrent therapy with ribavirin and peginterferon alfa.

## VIDAZA (S)

---

### Products Affected

- Azacitidine

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Advanced malignant hepatic tumor.
<b>Required Medical Information</b>	Patient has a diagnosis of one of the following myelodysplastic syndrome subtypes: refractory anemia, refractory anemia with ringed sideroblasts (if accompanied by neutropenia or thrombocytopenia or requiring transfusions), refractory anemia with excess blasts, refractory anemia with excess blasts in transformation, or chronic myelomonocytic leukemia.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## VIEKIRA (S)

### Products Affected

- Viekira Pak

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh C). Known hypersensitivity to ritonavir (e.g., toxic epidermal necrolysis or Stevens-Johnson syndrome). Concomitant use with drugs that are highly dependent on cytochrome P450 enzyme (CYP)3A for clearance, strong inducers of CYP3A and CYP2C8, and strong inhibitors of CYP2C8. Concomitant use with any of the following medications: Alfuzosin, Carbamazepine, phenytoin, phenobarbital, Gemfibrozil, Rifampin, Ergotamine, dihydroergotamine, ergonovine, methylergonovine, Ethinyl estradiol-containing medications, such as combined oral contraceptives, St. John's Wort, Lovastatin, simvastatin, Pimozide, Sildenafil (REVATIO), Triazolam, orally-administered midazolam
<b>Required Medical Information</b>	Patient has a diagnosis of chronic hepatitis C virus (HCV) genotype 1 infection AND Patient has tried and had an inadequate response, contraindication, or intolerance to HARVONI AND Patient does not have decompensated cirrhosis AND Patient will not use another HCV protease inhibitor (e.g., VICTRELIS, OLYSIO) or nucleotide analog polymerase inhibitor (i.e. SOVALDI, HARVONI) in combination with VIEKIRA AND The patient meets one of the following: Genotype 1a or 1b with compensated cirrhosis AND VIEKIRA will be used with ribavirin, Genotype 1a without cirrhosis AND VIEKIRA will be used with ribavirin, Liver transplant recipients without cirrhosis and Metavir fibrosis score 2 or less AND VIEKIRA will be used with ribavirin, or Genotype 1b infection without cirrhosis AND VIEKIRA will be used alone.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	hepatologist, gastroenterologist, or infectious disease specialist
<b>Coverage Duration</b>	12 weeks. 24 weeks for G1 with cirrhosis and liver transplant recipients.
<b>Other Criteria</b>	N/A

## VOTRIENT (S)

---

### Products Affected

- Votrient

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of advanced/metastatic renal cell carcinoma OR Diagnosis of advanced soft tissue sarcoma and patient received at least one prior chemotherapy (e.g., ifosfamide, doxorubicin, cisplatin, dacarbazine, docetaxel, oxaliplatin, etc.)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## VPRIV (BCBS RI)

---

### Products Affected

- Vpriv

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of type 1 Gaucher disease confirmed by enzymatic testing AND 1 or more complication such as anemia, thrombocytopenia, bone disease, hepatomegaly, and splenomegaly AND will not be used in combination with Zavesca
<b>Age Restrictions</b>	4 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## XALKORI (BCBS RI)

---

### Products Affected

- Xalkori

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of locally advanced or metastatic (stage III or IV) non-small cell lung cancer AND patient has non-squamous cell histology AND Patient has alkaline phosphatase (ALK)-positive disease as detected with an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility. Use as monotherapy
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A



# XELJANZ (S)

## Products Affected

- Xeljanz

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Active serious infection (including tuberculosis). Combined use with a biologic disease-modifying anti-rheumatic drugs or potent immunosuppressant (e.g., azathioprine or cyclosporine)
<b>Required Medical Information</b>	Diagnosis of moderately to severely active rheumatoid arthritis AND patient tried and had an inadequate response, intolerance or contraindication to methotrexate or other non-biologic disease modifying anti-rheumatic drugs for at least 3 consecutive months AND patient has a documented inadequate response to, or contraindication to, a biologic medication (e.g., Humira, Enbrel, Actemra, Cimzia, Orencia, Remicade, Simponi), unless there is a clinical reason to not use an injected product (e.g., fear of needles), patient has been tested for tuberculosis (TB) infection in the past year and latent TB has been ruled out or is being treated per guidelines.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient has stable disease or has improved while on therapy (e.g., improvement in tender/swollen joint count, improvement in ACR scoring)

## XENAZINE (S)

### Products Affected

- Xenazine

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Tourette's syndrome.
<b>Exclusion Criteria</b>	Actively suicidal. Untreated or inadequately treated depression. Impaired hepatic function. Concomitant use of monoamine oxidase inhibitors or within a minimum of 14 days after discontinuing a MAOI. Concomitant use of reserpine or within 20 days of discontinuing reserpine.
<b>Required Medical Information</b>	Diagnosis of chorea associated with Huntington's disease or tardive dyskinesia with failure of at least one previous therapy (e.g., amantadine, benzodiazepines, haloperidol, atypical antipsychotics, etc.) or Gilles de la Tourette's syndrome with failure or least one previous therapy (e.g., antipsychotic agents, clonidine) AND any medication possibly contributing to the underlying symptoms of chorea and/or tardive dyskinesia has been discontinued (e.g., anticonvulsants, antipsychotics, metoclopramide, amphetamines, etc.) unless cessation would be detrimental to the underlying condition AND patients who require doses greater than 50 mg/day will be genotyped for CYP2D6 to determine whether the patient is a poor, intermediate or extensive metabolizer.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, patient had lack of progression of disease or improvement in abnormal movements.

## XEOMIN (S)

---

### Products Affected

- Xeomin INJ 50UNIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Hypersensitivity to human albumin or sucrose. Presence of infection at the proposed injection site(s).
<b>Required Medical Information</b>	Diagnosis of blepharospasm and patient has been previously treated with Botox OR cervical dystonia (spasmodic torticollis) AND Xeomin will not be used for cosmetic uses (e.g., wrinkles)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal of cervical dystonia, patient experienced improvement in at least one of the following: pain, severity of abnormal head position, or effects on the patient's daily activities. For renewal of blepharospasm, patient experienced an improvement in symptoms (e.g., ability to open eyes, improvement in blinking/spasms of the eye).

## XGEVA (BCBS RI)

---

### Products Affected

- Xgeva

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Hypocalcemia (calcium less than 8.0 mg/dL).
<b>Required Medical Information</b>	Diagnosis of a solid tumor (e.g., breast cancer, castrate-resistant prostate cancer, thyroid carcinoma, kidney, or non-small cell lung cancer) and patient has bone metastases and Medication will be used for the prevention of skeletal-related events (e.g., spinal cord compression, hypercalcemia, bone pain or lesions requiring radiation or surgery) OR Patient has a diagnosis of giant cell tumor of bone and Tumor is unresectable or surgical resection is likely to result in severe morbidity AND patient will receive supplementation with calcium and vitamin D.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## XIFAXAN (S)

---

### Products Affected

- Xifaxan

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Allergy to rifamycin agents
<b>Required Medical Information</b>	Diagnosis of traveler's diarrhea and patient does not have fever or blood in the stool OR Diagnosis of hepatic encephalopathy and tried and failed lactulose therapy
<b>Age Restrictions</b>	Traveler's diarrhea - 12 years of age or older. hepatic encephalopathy - 18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	Traveler's diarrhea - 3 days. hepatic encephalopathy - 6 months
<b>Other Criteria</b>	N/A

# XOLAIR (S)

## Products Affected

- Xolair

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of moderate to severe persistent allergic asthma AND Evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. radioallergosorbent test) for a specific IgE or in vitro reactivity to a perennial aeroallergen AND Pretreatment serum IgE levels greater than or equal to 30 and less than or equal to 700 IU/mL AND Symptoms are not adequately controlled with high-dose inhaled corticosteroid (ICS) plus long-acting beta2-agonist (LABA) for at least 3 months AND patient has been adherent within a 12 month period, and is currently adherent, with asthma therapy.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	Asthma specialist (i.e., allergist, immunologist, or pulmonologist)
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	For renewal, patient has experienced an objective response to therapy, defined as one or more of the following: Reduction in number of asthma exacerbations from baseline (i.e. asthma exacerbation requiring treatment with systemic corticosteroids or doubling of ICS dose from baseline), Improvement in forced expiratory volume in 1 second (FEV1) from baseline, Decreased use of rescue medications from baseline.

# XTANDI (S)

---

## Products Affected

- Xtandi

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of metastatic castration-resistant prostate cancer AND patient had prior chemotherapy that included docetaxel AND the patient has tried and had an inadequate response, contraindication or intolerance to Zytiga.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

# XYREM (BCBS RI)

## Products Affected

- Xyrem

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Concomitant treatment with sedative hypnotic agents. Succinic semialdehyde dehydrogenase deficiency.
<b>Required Medical Information</b>	Diagnosis of one of the following: A) narcolepsy with excessive daytime sleepiness, cataplexy or both confirmed by sleep lab evaluation (e.g., multiple sleep latency test, polysomnography) and for patients with excessive daytime sleepiness, patient has had a previous trial with or has a contraindication, intolerance, or allergy to modafinil, armodafinil, methylphenidate, dextroamphetamine, or mixed amphetamine salts B) fibromyalgia syndrome and patient had a previous trial (of at least 30 days) with, or has a contraindication, intolerance, or allergy to two of the following: duloxetine, milnacipran, or pregabalin.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	Provider and patient must be enrolled in the Xyrem Success Program
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	For renewal, the patient had a positive response to the medication (increased sleep quality for patients with narcolepsy). Patient and physician are enrolled in the Xyrem Success Program.



# YERVOY (S)

## Products Affected

- Yervoy INJ 50MG/10ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of unresectable or metastatic melanoma AND If the request is for re-induction, the patient had no significant toxicity with the prior course of Yervoy AND the patient experienced progression after having stable disease for longer than three months or relapse after having a clinical response to therapy AND the prescriber is aware of the Yervoy REMS program.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	16 weeks
<b>Other Criteria</b>	N/A

## ZALTRAP (S)

---

### Products Affected

- Zaltrap INJ 100MG/4ML

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	severe hemorrhage, development of gastrointestinal perforation, compromised wound healing
<b>Required Medical Information</b>	Diagnosis of metastatic colon or rectal cancer AND will be used in combination with irinotecan or 5-fluorouracil, leucovorin, and irinotecan (FOLFIRI) AND disease is resistant to or has progressed following an oxaliplatin-containing regimen (e.g. 5-fluorouracil, leucovorin, and oxaliplatin [FOLFOX])
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Patient will be monitored for signs and symptoms of gastrointestinal bleeding and other severe bleeding. Therapy will be suspended for at least 4 weeks prior to elective surgery and not resumed for at least 4 weeks following major surgery and until the wound is fully healed.

## ZAVESCA (S)

---

### Products Affected

- Zavesca

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ZELBORAF (S)

---

### Products Affected

- Zelboraf

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility.
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ZOHYDRO (BCBS RI)

---

### Products Affected

- Zohydro Er

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D
<b>Exclusion Criteria</b>	Significant respiratory depression. Acute or severe bronchial asthma or hypercarbia. Known or suspected paralytic ileus.
<b>Required Medical Information</b>	Diagnosis of severe pain requiring continuous, around the clock opioid long term opioid treatment AND patient who are opioid tolerant receiving for 1 week or longer at least: 60mg oral morphine/day or 25mcg fentanyl patch/hour or 30mg oxycodone/day or 8mg hydromorphone/day or 25mg oxmorphone/day AND patient has tried and failed or unable to tolerate at least two generic extended-release opioid products, such as: oxymorphone ER, morphine ER, fentanyl, tramadol ER
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Provider must be enrolled in the REMS program
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Not covered for members enrolled in a Medicare approved hospice program

## ZOLINZA (S)

---

### Products Affected

- Zolinza

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of cutaneous T-cell lymphoma AND progressive, persistent or recurrent disease or patient is not a candidates for or following 2 systemic therapies (e.g., bexarotene, romidepsin, etc.)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ZOMETA (S)

### Products Affected

- Zoledronic Acid INJ 4MG/5ML

- Zometa INJ 4MG/100ML

PA Criteria	Criteria Details
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D. Osteopenia due to hormone therapy or androgen deprivation therapy. Osteopenia or osteoporosis due to monoclonal gammopathy of uncertain significance.
<b>Exclusion Criteria</b>	Current treatment with Reclast.
<b>Required Medical Information</b>	Diagnosis of hypercalcemia of malignancy and has a corrected calcium greater than or equal to 12 mg/dL OR Diagnosis of multiple myeloma and associated bone disease (e.g., osteolytic bone lesions, bone metastases, osteopenia, etc.) OR Diagnosis of a solid tumor (e.g., breast cancer, prostate cancer that has progressed after at least one hormonal therapy (i.e. antiandrogen [bicalutamide, flutamide, nilutamide], LHRH agonist [leuprolide, goserelin], LHRH antagonists [degarelix]), kidney cancer, non-small cell lung cancer, or thyroid cancer) and patient has bone metastases and medication will be used in conjunction with standard antineoplastic therapy and medication is used for the prevention of skeletal-related events (e.g. fracture, spinal cord compression, hypercalcemia, bone pain or lesions requiring radiation or surgery) AND Patient has tried and had an inadequate response or has a contraindication/intolerance to pamidronate OR patient has one of the following: osteopenia (T-score -1.0 to -2.5) secondary to androgen deprivation therapy and patient has a diagnosis of prostate cancer, osteopenia due to hormone therapy and patient has a diagnosis of breast cancer, or osteopenia or osteoporosis and patient has a diagnosis of monoclonal gammopathy of uncertain significant (MGUS)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	hypercalcemia of malignancy - 1 month. Others - 12 months

<b>Other Criteria</b>	Retreatment for hypercalcemia of malignancy will be considered a minimum of 7 days after initial treatment, if serum calcium does not return to normal or does not remain normal after initial treatment. For renewal of therapy for patients with bone metastases, skeletal-related events such as fractures have decreased or stabilized. For renewal of therapy for osteoporosis or osteopenia, improved or stabilized BMD, no new fractures, etc.
-----------------------	---



## ZORBTIVE (BCBS RI)

---

### Products Affected

- Zorbtive

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	Acute critical illness due to complications following open heart or abdominal surgery, multiple accidental trauma or acute respiratory failure. Active malignancy. Active proliferative or severe non-proliferative diabetic retinopathy. Active malignancy or undergoing cancer treatment.
<b>Required Medical Information</b>	Diagnosis of short bowel syndrome AND patient is receiving specialized nutritional support (i.e. parenteral nutrition)
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	4 weeks
<b>Other Criteria</b>	N/A

## ZORTRESS (S)

---

### Products Affected

- Zortress

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Medication is being used for: A) Prevention of kidney transplant organ rejection AND patient is at low-to-moderate immunologic risk AND member is prescribed concurrent therapy with reduced doses of cyclosporine and corticosteroids, or B) Prevention of liver transplant organ rejection AND 30 or more days have passed since the transplant procedure AND the member is prescribed concurrent therapy with reduced doses of tacrolimus and corticosteroids
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Prescriber is experienced in immunosuppressive therapy and management of transplant patients.
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	Part B if transplant covered by Medicare. otherwise Part D

## ZYGELIG (S)

### Products Affected

- Zydelig

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	The patient has one of the following diagnoses: A) chronic lymphocytic leukemia AND The medication will be used in combination with rituximab AND The patient has relapsed on at least one prior therapy (e.g., purine analogues [fludarabine, pentostatin, cladribine], alkylating agents [chlorambucil, cyclophosphamide], or monoclonal antibodies [rituximab]) AND the patient does not have any co-morbidities that prevents the use of cytotoxic chemotherapy (i.e. severe neutropenia or thrombocytopenia, creatinine clearance less than 60 mL/minute), B) follicular lymphoma AND the patient has relapsed on at least two prior systemic therapies (e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine]), or C) small lymphocytic lymphoma AND The patient has relapsed on at least two prior systemic therapies(e.g., rituximab, alkylating agents [cyclophosphamide, chlorambucil], anthracyclines [doxorubicin, daunorubicin], purine analogs [fludarabine])
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ZYKADIA (S)

---

### Products Affected

- Zykadia

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of metastatic non-small cell lung cancer AND patient has anaplastic lymphoma kinase (ALK)-positive disease as detected by an FDA-approved or Clinical Laboratory Improvement Amendments (CLIA)-approved facility AND patient had an inadequate response, progressed on, or had an intolerance or contraindication to Xalkori
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## ZYTIGA (S)

---

### Products Affected

- Zytiga

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Covered Uses</b>	All FDA-approved indications not otherwise excluded from Part D.
<b>Exclusion Criteria</b>	N/A
<b>Required Medical Information</b>	Diagnosis of metastatic castration-resistant prostate cancer AND Zytiga will be used in combination with prednisone
<b>Age Restrictions</b>	N/A
<b>Prescriber Restrictions</b>	N/A
<b>Coverage Duration</b>	12 months
<b>Other Criteria</b>	N/A

## PART B VERSUS PART D

---

### Products Affected

- Abelcet
- Abraxane
- Acetylcysteine INHALATION SOLN
- Acyclovir Sodium INJ 50MG/ML
- Adriamycin INJ 2MG/ML
- Adrucil INJ 2.5GM/50ML
- Albuterol Sulfate INHALATION NEBU
- Alimta INJ 500MG
- Aloxi
- Amifostine
- Amino Acids
- Aminophylline INJ
- Aminosyn INJ 148MEQ/L; 1280MG/100ML; 980MG/100ML; 1280MG/100ML; 300MG/100ML; 720MG/100ML; 940MG/100ML; 720MG/100ML; 400MG/100ML; 440MG/100ML; 5.4MEQ/L; 860MG/100ML; 420MG/100ML; 520MG/100ML; 160MG/100ML; 44MG/100ML; 800MG/100ML, 50.3MEQ/L; 695MG/100ML; 713MG/100ML; 490MG/100ML; 517MG/100ML; 350MG/100ML; 210MG/100ML; 462MG/100ML; 700MG/100ML; 735MG/100ML; 120MG/100ML; 209MG/100ML; 505MG/100ML; 371MG/100ML; 31.3MEQ/L; 280MG/100ML; 140MG/100ML; 189MG/100ML; 350MG/100ML, 51MEQ/L; 448MG/100ML; 343MG/100ML; 448MG/100ML; 105MG/100ML; 252MG/100ML; 329MG/100ML; 252MG/100ML; 140MG/100ML; 154MG/100ML; 300MG/100ML; 147MG/100ML; 182MG/100ML; 56MG/100ML; 31MG/100ML; 280MG/100ML, 90MEQ/L; 1100MG/100ML; 850MG/100ML; 35MEQ/L; 1100MG/100ML; 260MG/100ML; 620MG/100ML; 810MG/100ML; 624MG/100ML; 340MG/100ML; 380MG/100ML; 5.4MEQ/L; 750MG/100ML; 370MG/100ML; 460MG/100ML; 150MG/100ML; 44MG/100ML; 680MG/100ML
- Aminosyn 7%/electrolytes
- Aminosyn 8.5%/electrolytes

- Aminosyn II INJ 50.3MEQ/L; 695MG/100ML; 713MG/100ML; 490MG/100ML; 517MG/100ML; 350MG/100ML; 210MG/100ML; 462MG/100ML; 700MG/100ML; 735MG/100ML; 120MG/100ML; 209MG/100ML; 505MG/100ML; 371MG/100ML; 31.3MEQ/L; 280MG/100ML; 140MG/100ML; 189MG/100ML; 350MG/100ML, 61.1MEQ/L; 844MG/100ML; 865MG/100ML; 595MG/100ML; 627MG/100ML; 425MG/100ML; 255MG/100ML; 561MG/100ML; 850MG/100ML; 893MG/100ML; 146MG/100ML; 253MG/100ML; 614MG/100ML; 450MG/100ML; 33.3MEQ/L; 340MG/100ML; 170MG/100ML; 230MG/100ML; 425MG/100ML, 71.8MEQ/L; 993MG/100ML; 1018MG/100ML; 700MG/100ML; 738MG/100ML; 500MG/100ML; 300MG/100ML; 660MG/100ML; 1000MG/100ML; 1050MG/100ML; 172MG/100ML; 298MG/100ML; 722MG/100ML; 530MG/100ML; 45.3MEQ/L; 400MG/100ML; 200MG/100ML; 270MG/100ML; 500MG/100ML
- Aminosyn II 4.25/dextrose25% INJ 30.6MEQ/L; 422MG/100ML; 432MG/100ML; 298MG/100ML; 25%; 314MG/100ML; 212MG/100ML; 128MG/100ML; 280MG/100ML; 425MG/100ML; 446MG/100ML; 73MG/100ML; 126MG/100ML; 307MG/100ML; 225MG/100ML; 19MEQ/L; 170MG/100ML; 85MG/100ML; 115MG/100ML; 212MG/100ML
- Aminosyn II 8.5%/electrolytes
- Aminosyn-hbc
- Aminosyn-hf
- Aminosyn-pf
- Aminosyn-pf 7%
- Aminosyn-rf
- Amiodarone Hcl INJ 50MG/ML
- Amphotericin B INJ
- Ampicillin Sodium INJ 10GM, 125MG, 1GM, 2GM
- Ampicillin-sulbactam
- Anzemet ORAL TABS
- Aralast Np INJ 400MG
- Arranon
- Arzerra INJ 100MG/5ML
- Astagraf XL
- Atgam
- Azactam In Iso-osmotic Dextrose
- Azasan
- Azathioprine TABS
- Azathioprine Sodium
- Azithromycin INJ 500MG
- Bicnu
- Bleomycin Sulfate INJ 30UNIT
- Brovana
- Budesonide INHALATION SUSP 0.25MG/2ML, 0.5MG/2ML
- Busulfex
- Calcitriol INJ 1MCG/ML
- Calcitriol ORAL CAPS
- Calcitriol ORAL SOLN
- Carboplatin INJ 150MG/15ML
- Cefazolin Sodium INJ 10GM, 1GM, 1GM; 5%, 500MG
- Cefotaxime Sodium INJ 10GM, 1GM, 2GM, 500MG
- Cefoxitin Sodium INJ 10GM, 1GM, 2GM
- Ceftazidime INJ 1GM, 2GM, 6GM
- Ceftriaxone Sodium INJ
- Cefuroxime Sodium INJ 1.5GM, 7.5GM, 7.5GM, 750MG
- Cefuroxime/dextrose
- Cellcept SUSR
- Cellcept Intravenous
- Chlorothiazide Sodium
- Cidofovir
- Ciprofloxacin INJ 400MG/40ML
- Ciprofloxacin I.v.-in D5w
- Cisplatin INJ 100MG/100ML

- Clindamycin Phosphate Add-vantage
- Clindamycin Phosphate In D5w
- Clinimix 2.75%/dextrose 5%
- Clinimix 4.25%/dextrose 10%
- Clinimix 4.25%/dextrose 20%
- Clinimix 4.25%/dextrose 25%
- Clinimix 4.25%/dextrose 5%
- Clinimix 5%/dextrose 15%
- Clinimix 5%/dextrose 20%
- Clinimix 5%/dextrose 25%
- Clinimix E 4.25%/dextrose 10%
- Clinisol Sf 15%
- Clolar
- Colistimethate Sodium INJ
- Cosmegen
- Coumadin INJ
- Cromolyn Sodium NEBU
- Cubicin
- Cyclophosphamide ORAL TABS
- Cyclosporine INJ
- Cyclosporine ORAL CAPS
- Cyclosporine Modified
- Cytarabine INJ 500MG
- Cytarabine Aqueous
- Dacarbazine INJ 200MG
- Dactinomycin
- Daunorubicin Hcl INJ 5MG/ML
- Decitabine
- Dexrazoxane INJ 250MG
- Dextrose 10% Flex Container
- Dextrose 10%/nacl 0.2%
- Dextrose 10%/nacl 0.45%
- Dextrose 2.5%/sodium Chloride 0.45%
- Dextrose 5%
- Dextrose 5% /electrolyte #48 Viaflex
- Dextrose 5%/nacl 0.2%
- Dextrose 5%/nacl 0.225%
- Dextrose 5%/nacl 0.33%
- Dextrose 5%/nacl 0.45%
- Dextrose 5%/nacl 0.9%
- Dextrose 5%/potassium Chloride 0.15%
- Diltiazem Hcl INJ 100MG, 50MG/10ML
- Docefrez
- Docetaxel INJ 140MG/7ML, 20MG/ML, 80MG/4ML, 80MG/8ML
- Doxercalciferol ORAL CAPS
- Doxil
- Doxorubicin Hcl INJ 2MG/ML
- Doxorubicin Hcl Liposome
- Elitek INJ 1.5MG
- Ellence INJ 200MG/100ML
- Emend ORAL CAPS
- Engerix-b
- Epirubicin Hcl INJ 50MG/25ML
- Eraxis INJ 100MG
- Erbitux INJ 100MG/50ML
- Esomeprazole Sodium
- Etopophos
- Etoposide INJ
- Famotidine Premixed
- Fluconazole In Dextrose INJ 56MG/ML; 400MG/200ML
- Fludarabine Phosphate
- Fluorouracil INJ 2.5GM/50ML
- Freamine Hbc 6.9%
- Freamine III INJ 89MEQ/L; 710MG/100ML; 950MG/100ML; 3MEQ/L; 24MG/100ML; 1400MG/100ML; 280MG/100ML; 690MG/100ML; 910MG/100ML; 730MG/100ML; 530MG/100ML; 560MG/100ML; 10MMOLE/L; 120MG/100ML; 1120MG/100ML; 590MG/100ML; 10MEQ/L; 400MG/100ML; 150MG/100ML; 660MG/100ML
- Ganciclovir INJ
- Gemcitabine Hcl INJ 1GM
- Gengraf
- Granisetron Hcl INJ 0.1MG/ML, 1MG/ML
- Granisetron Hcl TABS
- Halaven
- Hecoria



- Heparin Sodium INJ  
10000UNIT/ML, 1000UNIT/ML,  
20000UNIT/ML, 2000UNIT/ML,  
2500UNIT/ML, 5000UNIT/ML
- Heparin Sodium/d5w
- Hepatamine
- Hepatasol
- Herceptin
- Idarubicin Hcl INJ 10MG/10ML
- Ifosfamide INJ 1GM
- Ifosfamide/mesna INJ 1GM; 1GM
- Imipenem/cilastatin
- Imovax Rabies (h.d.c.v.)
- Intralipid INJ 20GM/100ML
- Ionosol-b/dextrose 5%
- Ionosol-mb/dextrose 5%
- Ipratropium Bromide INHALATION  
SOLN 0.02%
- Ipratropium Bromide/albuterol  
Sulfate
- Irinotecan INJ 100MG/5ML
- Isolyte-p/dextrose 5%
- Isolyte-s
- Istodax
- Ixempra Kit INJ 45MG
- Jevtana
- Kcl 0.075%/d5w/nacl 0.45%
- Kcl 0.15%/d5w/ Nacl 0.3%
- Kcl 0.15%/d5w/lr
- Kcl 0.15%/d5w/nacl 0.2%
- Kcl 0.15%/d5w/nacl 0.225%
- Kcl 0.15%/d5w/nacl 0.9%
- Kcl 0.3%/d5w/nacl 0.45%
- Kcl 0.3%/d5w/nacl 0.9%
- Kepivance
- Labetalol Hcl INJ
- Lactated Ringers Viaflex
- Levalbuterol NEBU
- Levalbuterol Hcl INHALATION  
NEBU
- Levaquin INJ 5%; 750MG/150ML
- Levetiracetam INJ 500MG/5ML
- Levocarnitine INJ
- Levocarnitine ORAL SOLN
- Levocarnitine TABS
- Levofloxacin INJ
- Levofloxacin In D5w INJ 5%;  
500MG/100ML
- Linezolid
- Lioresal Intrathecal INJ 0.05MG/ML,  
10MG/20ML, 10MG/5ML
- Liothyronine Sodium INJ
- Lipodox
- Lipodox 50
- Liposyn III INJ 1.2GM/100ML;  
2.5GM/100ML; 10GM/100ML,  
2.5%; 30%
- Magnesium Sulfate In D5w INJ 5%;  
10MG/ML
- Melphalan Hydrochloride
- Meropenem
- Mesna
- Methotrexate TABS
- Metoprolol Tartrate INJ
- Metronidazole In Nacl 0.79%
- Mitomycin INJ 20MG
- Mitoxantrone Hcl
- Mustargen
- Mycophenolate Mofetil
- Mycophenolic Acid Dr
- Nafcillin Sodium
- Nallpen/dextrose INJ 0; 1GM/50ML
- Nebupent
- Nephramine
- Neutrexin
- Nitroglycerin INJ
- Normosol-r
- Normosol-r In D5w
- Nulojix
- Nutrilipid
- Ondansetron Hcl INJ 4MG/2ML
- Ondansetron Hcl ORAL SOLN
- Ondansetron Hcl ORAL TABS
- Ondansetron Odt
- Oxaliplatin INJ 100MG/20ML
- Paclitaxel INJ 300MG/50ML
- Pantoprazole Sodium INJ
- Paricalcitol INJ 5MCG/ML

- Paricalcitol ORAL CAPS
- Penicillin G Potassium In Iso-osmotic Dextrose
- Pentam 300
- Pentostatin
- Perforomist
- Piperacillin Sodium/ Tazobactam Sodium
- Piperacillin Sodium/tazobactam Sodium INJ 3GM; 0.375GM, 4GM; 0.5GM
- Piperacillin/tazobactam INJ 2GM; 0.25GM, 4GM; 0.5GM
- Plasma-lyte A
- Plasma-lyte-148
- Plasma-lyte-56/d5w
- Potassium Chloride INJ 10MEQ/100ML, 20MEQ/100ML, 2MEQ/ML, 30MEQ/100ML, 40MEQ/100ML
- Potassium Chloride 0.15% /nacl 0.45% Viaflex
- Potassium Chloride 0.15% D5w/nacl 0.33%
- Potassium Chloride 0.15% D5w/nacl 0.45%
- Potassium Chloride 0.15% D5w/nacl 0.45% Viaflex
- Potassium Chloride 0.15% W/nacl 0.9% Viaflex
- Potassium Chloride 0.15%/nacl 0.9%
- Potassium Chloride 0.22% D5w/nacl 0.45%
- Potassium Chloride 0.3%/ Nacl 0.9%
- Potassium Chloride 0.3%/d5w
- Premasol
- Prograf INJ
- Prolastin
- Prolastin-c
- Proleukin
- Propranolol Hcl INJ
- Pulmicort SUSP 1MG/2ML
- Rabavert
- Rapamune ORAL TABS 1MG, 2MG
- Rapamune SOLN
- Recombivax Hb
- Rifampin INJ
- Ringers Injection
- Rituxan
- Sandimmune SOLN
- Simulect INJ 20MG
- Sirolimus ORAL TABS
- Sodium Chloride INJ 0.9%, 2.5MEQ/ML, 3%, 5%
- Sodium Chloride 0.45% Viaflex
- Sodium Edecrin
- Sulfamethoxazole/trimethoprim INJ
- Tacrolimus ORAL CAPS
- Taxotere INJ 80MG/2ML
- Tazicef
- Teflaro
- Tetanus Toxoid Adsorbed
- Thiotepa INJ
- Thymoglobulin
- Tobramycin Sulfate INJ 10MG/ML, 80MG/2ML
- Tobramycin Sulfate/sodium Chloride INJ 0.9%; 0.8MG/ML
- Toposar
- Topotecan Hcl INJ 4MG
- Torisel
- Toremide INJ 20MG/2ML
- Tranexamic Acid INJ
- Travasol
- Treanda INJ 100MG, 180MG/2ML, 45MG/0.5ML
- Trexall
- Trisenox
- Trophamine INJ 97MEQ/L; 0.54GM/100ML; 1.2GM/100ML; 0.32GM/100ML; 0; 0; 0.5GM/100ML; 0.36GM/100ML; 0.48GM/100ML; 0.82GM/100ML; 1.4GM/100ML; 1.2GM/100ML; 0.34GM/100ML; 0.48GM/100ML; 0.68GM/100ML; 0.38GM/100ML; 5MEQ/L; 0.025GM/100ML; 0.42GM/100ML; 0.2GM/100ML; 0.24GM/100ML; 0.78GM/100ML
- Twinrix

- Tygacil
- Valproate Sodium INJ
- Vancomycin Hcl INJ 1000MG, 10GM, 500MG
- Vectibix INJ 100MG/5ML
- Velcade
- Verapamil Hcl INJ
- Vibativ INJ 250MG
- Vinblastine Sulfate INJ 1MG/ML
- Vincasar Pfs
- Vincristine Sulfate INJ
- Vinorelbine Tartrate INJ 50MG/5ML
- Virazole
- Zanosar
- Zemplar INJ
- Zosyn INJ 5%; 2GM/50ML; 0.25GM/50ML, 5%; 3GM/50ML; 0.375GM/50ML, 5%; 4GM/100ML; 0.5GM/100ML
- Zyvox INJ

## Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

# INDEX

## A

Abelcet .....	238	Amino Acids .....	238
Abraxane .....	238	Aminophylline .....	238
Abstral .....	58	Aminosyn .....	238
Acetylcysteine .....	238	Aminosyn 7%/electrolytes .....	238
Acitretin .....	1	Aminosyn 8.5%/electrolytes .....	238
Acitretin (s) .....	1	Aminosyn II .....	239
Actemra .....	2	Aminosyn II 4.25/dextrose25% .....	239
Actemra (bcbs Ri) .....	2	Aminosyn II 8.5%/electrolytes .....	239
Actiq (bcbs Ri) .....	3	Aminosyn-hbc .....	239
Acyclovir Sodium .....	238	Aminosyn-hf .....	239
Adagen .....	4	Aminosyn-pf .....	239
Adagen (s) .....	4	Aminosyn-pf 7% .....	239
Adapalene .....	42	Aminosyn-rf .....	239
Adcirca .....	5	Amiodarone Hcl .....	239
Adcirca (s) .....	5	Amitiza .....	10
Adefovir Dipivoxil .....	73	Amitiza (s) .....	10
Adempas .....	6	Amitriptyline Hcl .....	77
Adempas (s) .....	6	Amphotericin B .....	239
Adriamycin .....	238	Ampicillin Sodium .....	239
Adrucil .....	238	Ampicillin-sulbactam .....	239
Afinitor .....	7	Ampyra .....	11
Afinitor (s) .....	7	Ampyra (bcbs Ri) .....	11
Afinitor Disperz .....	8	Amrix .....	89
Afinitor Disperz (s) .....	8	Anadrol-50 .....	12
Albuterol Sulfate .....	238	Anadrol-50 (s) .....	12
Aldurazyme .....	9	Anzemet .....	239
Aldurazyme (s) .....	9	Aralast Np .....	239
Alimta .....	238	Aranesp (s) .....	13
Alora .....	86	Aranesp Albumin Free .....	13
Aloxi .....	238	Arbinoxa .....	79
Alprazolam .....	21	Arcalyst .....	15
Alprazolam Er .....	21	Arcalyst (s) .....	15
Alprazolam Intensol .....	21	Arranon .....	239
Alprazolam Odt .....	21	Arzerra .....	239
Alprazolam Xr .....	21	Astagraf XL .....	239
Amifostine .....	238	Atgam .....	239
		Aubagio .....	16
		Aubagio (bcbs Ri) .....	16

Avastin .....	17	Carimune Nanofiltered.....	104
Avastin (s) .....	17	Carisoprodol.....	89
Avonex .....	18	Carisoprodol/aspirin.....	89
Avonex (s).....	18	Carisoprodol/aspirin/codeine .....	89
Azacitidine .....	212	Cayston .....	30
Azactam In Iso-osmotic Dextrose.....	239	Cayston (s) .....	30
Azasan .....	239	Cefazolin Sodium.....	239
Azathioprine .....	239	Cefotaxime Sodium .....	239
Azathioprine Sodium.....	239	Cefoxitin Sodium .....	239
Azithromycin.....	239	Ceftazidime .....	239
<b>B</b>			
Baraclude.....	19	Ceftriaxone Sodium .....	239
Baraclude (bcbs Ri).....	19	Cefuroxime Sodium .....	239
Beleodaq.....	20	Cefuroxime/dextrose.....	239
Beleodaq (s).....	20	Celebrex .....	31
Benzodiazepines (s).....	21	Celebrex (s).....	31
Benzotropine Mesylate.....	82	Celecoxib .....	31
Bethkis.....	22	Cellcept .....	239
Bethkis (bcbs Ri).....	22	Cellcept Intravenous .....	239
Bicnu .....	239	Cenestin.....	86
Bivigam .....	104	Cerdelga .....	32
Bleomycin Sulfate .....	239	Cerdelga (s).....	32
Blinicyto .....	207	Cerezyme .....	33
Boniva .....	23	Cerezyme (bcbs Ri).....	33
Boniva IV (bcbs Ri) .....	23	Cesamet.....	34
Bosulif .....	24	Cesamet (s).....	34
Bosulif (bcbs Ri) .....	24	Chlordiazepoxide Hcl .....	21
Botox .....	25	Chlordiazepoxide/amitriptyline .....	77
Botox (bcbs Ri) .....	25	Chlorothiazide Sodium .....	239
Brovana .....	239	Chlorpropamide .....	90
Budesonide .....	239	Chlorzoxazone .....	89
Buprenorphine Hcl .....	186	Cialis .....	35
Buprenorphine Hcl/naloxone Hcl.....	185	Cialis 2.5 Mg, 5 Mg (s).....	35
Busulfex .....	239	Cidofovir.....	239
Butrans .....	28	Cimzia.....	36
Butrans (s) .....	28	Cimzia (s).....	36
<b>C</b>			
Calcipotriene/betamethasone Dipropionate .....	192	Ciprofloxacin .....	239
Calcitriol.....	239	Ciprofloxacin I.v.-in D5w.....	239
Caprelsa.....	29	Cisplatin .....	239
Caprelsa (bcbs Ri).....	29	Clemastine Fumarate .....	79
Carbinoxamine Maleate .....	79	Climara Pro .....	86
Carboplatin .....	239	Clindamycin Phosphate Add-vantage.....	240
		Clindamycin Phosphate In D5w .....	240
		Clinimix 2.75%/dextrose 5% .....	240
		Clinimix 4.25%/dextrose 10% .....	240

Clinimix 4.25%/dextrose 20% .....	240
Clinimix 4.25%/dextrose 25% .....	240
Clinimix 4.25%/dextrose 5% .....	240
Clinimix 5%/dextrose 15% .....	240
Clinimix 5%/dextrose 20% .....	240
Clinimix 5%/dextrose 25% .....	240
Clinimix E 4.25%/dextrose 10% .....	240
Clinisol Sf 15% .....	240
Clolar .....	240
Clomipramine Hcl .....	77
Clonazepam .....	21
Clonazepam Odt .....	21
Clorazepate Dipotassium .....	21
Colistimethate Sodium .....	240
Combipatch .....	86
Cometriq .....	37
Cometriq (s) .....	37
Copaxone .....	38
Copaxone (s) .....	38
Cosmegen .....	240
Coumadin .....	240
Cromolyn Sodium .....	240
Cubicin .....	240
Cyclobenzaprine Hcl .....	89
Cyclophosphamide .....	240
Cyclosporine .....	240
Cyclosporine Modified .....	240
Cyproheptadine Hcl .....	79
Cystaran .....	39
Cystaran (s) .....	39
Cytarabine .....	240
Cytarabine Aqueous .....	240

## **D**

Dacarbazine .....	240
Dactinomycin .....	240
Daliresp .....	40
Daliresp (s) .....	40
Daunorubicin Hcl .....	240
Decitabine .....	240
Dexrazoxane .....	240
Dextrose 10% Flex Container .....	240
Dextrose 10%/nacl 0.2% .....	240
Dextrose 10%/nacl 0.45% .....	240
Dextrose 2.5%/sodium Chloride 0.45% .....	240

Dextrose 5% .....	240
Dextrose 5% /electrolyte #48 Viaflex .....	240
Dextrose 5%/nacl 0.2% .....	240
Dextrose 5%/nacl 0.225% .....	240
Dextrose 5%/nacl 0.33% .....	240
Dextrose 5%/nacl 0.45% .....	240
Dextrose 5%/nacl 0.9% .....	240
Dextrose 5%/potassium Chloride 0.15% .....	240
Diazepam .....	21
Diazepam Intensol .....	21
Diclofenac Gel (s) .....	41
Differin (s) .....	42
Diltiazem Hcl .....	240
Diphenhydramine Hcl .....	78, 79
Dipyridamole .....	91
Divigel .....	86
Docefrez .....	240
Docetaxel .....	240
Doxepin Hcl .....	77
Doxercalciferol .....	240
Doxil .....	240
Doxorubicin Hcl .....	240
Doxorubicin Hcl Liposome .....	240
Dronabinol .....	125

## **E**

Edluar .....	88
Egrifta .....	43
Egrifta (s) .....	43
Elelyso .....	44
Elelyso (bcbs Ri) .....	44
Eligard .....	45
Eligard (s) .....	45
Elitek .....	240
Ellence .....	240
Emend .....	240
Emsam .....	46
Emsam (s) .....	46
Enbrel .....	47
Enbrel (bcbs Ri) .....	47
Enbrel Sureclick .....	47
Engerix-b .....	240
Enjuvia .....	86
Entecavir .....	19
Entyvio .....	207

Epiduo .....	49
Epiduo (s) .....	49
Epirubicin Hcl .....	240
Epoetin Alfa (s) .....	50
Epogen.....	50
Epoprostenol Sodium .....	209
Eraxis.....	240
Erbix.....	240
Ergoloid Mesylates.....	85
Erivedge .....	52
Erivedge (s) .....	52
Erwinaze.....	53
Erwinaze (s).....	53
Esbriet.....	54
Esbriet (s) .....	54
Esomeprazole Sodium.....	240
Estazolam .....	21
Estradiol .....	86
Estradiol/norethindrone Acetate.....	86
Estropipate.....	86
Etopophos.....	240
Etoposide.....	240
Exalgo (s) .....	55
Exjade.....	56
Exjade (s).....	56

## **F**

Fabrazyme .....	57
Fabrazyme (s).....	57
Famotidine Premixed .....	240
Fentanyl (s).....	58
Fentanyl Citrate Oral Transmucosal .....	3
Fentora.....	58
Ferriprox.....	59
Ferriprox (s).....	59
Firmagon .....	60
Firmagon (s) .....	60
Flebogamma Dif.....	104
Flector.....	61
Flector (s) .....	61
Fluconazole In Dextrose.....	240
Fludarabine Phosphate .....	240
Fluorouracil .....	240
Flurazepam Hcl .....	21
Forteo .....	62

Forteo (bcbs Ri) .....	62
Freamine Hbc 6.9% .....	240
Freamine III .....	240

## **G**

Gammagard Liquid .....	104
Gammaked .....	104
Gamunex-c.....	104
Ganciclovir.....	240
Gattex .....	63
Gattex (s).....	63
Gazyva .....	64
Gazyva (s).....	64
Gemcitabine Hcl .....	240
Gengraf .....	240
Genotropin .....	69
Genotropin Miniquick.....	69
Gilenya.....	65
Gilenya (bcbs Ri) .....	65
Gilotrif.....	66
Gilotrif (s) .....	66
Gleevec .....	67
Gleevec (s) .....	67
Glyburide .....	90
Glyburide Micronized.....	90
Glyburide/metformin Hcl.....	90
Gralise.....	68
Gralise (bcbs Ri) .....	68
Granisetron Hcl.....	240
Growth Hormone (s).....	69
Guanfacine Hcl .....	80

## **H**

H.p. Acthar.....	71
H.p. Acthar Gel (bcbs Ri) .....	71
Halaven .....	240
Harvoni .....	72
Harvoni (bcbs Ri).....	72
Hecoria.....	240
Heparin Sodium .....	241
Heparin Sodium/d5w .....	241
Hepatamine .....	241
Hepatasol.....	241
Hepsera .....	73
Hepsera (s) .....	73

Herceptin.....	241	Ilaris .....	97
Hexalen.....	74	Ilaris (s).....	97
Hexalen (s) .....	74	Imbruvica .....	98
Hizentra.....	104	Imbruvica (s).....	98
Horizant.....	75	Imipenem/cilastatin.....	241
Horizant (s).....	75	Imipramine Hcl .....	77
Hrm - Analgesics.....	76	Imipramine Pamoate .....	77
Hrm - Antidepressants.....	77	Imovax Rabies (h.d.c.v.).....	241
Hrm - Antiemetic Drugs.....	78	Increlex .....	99
Hrm - Antihistamines.....	79	Increlex (s) .....	99
Hrm - Antihypertensive Agents .....	80	Infergen .....	100
Hrm - Anti-infective (bcrl).....	81	Infergen (s).....	100
Hrm - Antiparkinson Agents.....	82	Inhaled Tobramycin (bcbs Ri) .....	101
Hrm - Anxiolytics.....	83	Inlyta .....	102
Hrm - Calcium Channel Blockers, Dihydropyridine .....	84	Inlyta (s).....	102
Hrm - Dementia Agents .....	85	Intralipid.....	241
Hrm - Oral And Transdermal Estrogens And Progestins .....	86	Intron A.....	103
Hrm - Platelet Inhibitors.....	87	Intron A (s).....	103
Hrm - Sedative Hypnotic Agents .....	88	Intron-a.....	103
Hrm - Skeletal Muscle Relaxants.....	89	Intron-a W/diluent.....	103
Hrm - Sulfonylureas .....	90	Ionosol-b/dextrose 5% .....	241
Hrm - Vasodilators .....	91	Ionosol-mb/dextrose 5% .....	241
Humatrope.....	69	Ipratropium Bromide .....	241
Humatrope Combo Pack .....	69	Ipratropium Bromide/albuterol Sulfate.....	241
Humira.....	92	Irinotecan .....	241
Humira (s) .....	92	Isolyte-p/dextrose 5% .....	241
Humira Pediatric Crohns Disease Starter Pack...	92	Isolyte-s.....	241
Humira Pen-crohns Diseasestarter .....	92	Istodax.....	241
Hydromorphone Hcl Er.....	55	Ivig (s).....	104
Hydroxyzine Hcl .....	79	Ixempra Kit.....	241
Hydroxyzine Pamoate .....	79		
Hyqvia .....	94	<b>J</b>	
Hyqvia (s).....	94	Jakafi.....	106
Hysingla (bcbs Ri).....	95	Jakafi (s).....	106
Hysingla Er.....	95	Jevtana.....	241
		Jinteli.....	86
<b>I</b>		Juxtapid.....	107
Ibandronate Sodium .....	23	Juxtapid (bcbs Ri) .....	107
Iclusig .....	96		
Iclusig (s).....	96	<b>K</b>	
Idarubicin Hcl.....	241	Kadcyla .....	108
Ifosfamide.....	241	Kadcyla (bcbs Ri) .....	108
Ifosfamide/mesna .....	241	Kalydeco .....	109
		Kalydeco (s).....	109
		Kcl 0.075%/d5w/nacl 0.45% .....	241



Kcl 0.15%/d5w/ Nacl 0.3%	241
Kcl 0.15%/d5w/lr	241
Kcl 0.15%/d5w/nacl 0.2%	241
Kcl 0.15%/d5w/nacl 0.225%	241
Kcl 0.15%/d5w/nacl 0.9%	241
Kcl 0.3%/d5w/nacl 0.45%	241
Kcl 0.3%/d5w/nacl 0.9%	241
Kepivance	241
Keytruda	110
Keytruda (s)	110
Kineret	111
Kineret (bcbs Ri)	111
Korlym	112
Korlym (s)	112
Kuvan	113
Kuvan (s)	113
Kynamro	114
Kynamro (s)	114

## L

Labetalol Hcl	241
Lactated Ringers Viaflex	241
Lazanda	58
Lemtrada	115
Lemtrada (bcbs Ri)	115
Letairis	116
Letairis (bcbs Ri)	116
Leukine	117
Leukine (s)	117
Leuprolide Acetate	123
Levalbuterol	241
Levalbuterol Hcl	241
Levaquin	241
Levetiracetam	241
Levocarnitine	241
Levofloxacin	241
Levofloxacin In D5w	241
Lidocaine	119
Lidoderm (s)	119
Linezolid	241
Linzess	120
Linzess (s)	120
Lioresal Intrathecal	241
Liothyronine Sodium	241
Lipodox	241

Lipodox 50	241
Liposyn III	241
Lorazepam	21
Lorazepam Intensol	21
Lumizyme	121
Lumizyme (s)	121
Lupaneta (s)	122
Lupaneta Pack	122
Lupron Depot	123, 124
Lupron Depot (s)	123
Lupron Depot- Ped (s)	124
Lupron Depot-ped	124
Lynparza	207

## M

Macrochantin	81
Magnesium Sulfate In D5w	241
Marinol (s)	125
Mekinist	126
Mekinist (s)	126
Melphalan Hydrochloride	241
Menest	86
Menostar	86
Meperidine Hcl	76
Meperitab	76
Meprobamate	83
Meropenem	241
Mesna	241
Methocarbamol	89
Methotrexate	241
Metoprolol Tartrate	241
Metronidazole In Nacl 0.79%	241
Mimvey	86
Mimvey Lo	86
Mitomycin	241
Mitoxantrone Hcl	241
Modafinil	127
Modafinil (s)	127
Mozobil	128
Mozobil (s)	128
Mustargen	241
Myalept	129
Myalept (s)	129
Mycophenolate Mofetil	241
Mycophenolic Acid Dr	241

Myozyme.....	130
Myozyme (s) .....	130
<b>N</b>	
Nafcillin Sodium .....	241
Naglazyme.....	131
Naglazyme (s) .....	131
Nallpen/dextrose.....	241
Nebupent .....	241
Nephramine .....	241
Neulasta.....	132
Neulasta (s).....	132
Neumega.....	133
Neumega (s) .....	133
Neupogen .....	134
Neupogen (s) .....	134
Neutrexin.....	241
Nexavar .....	136
Nexavar (s) .....	136
Nifedipine.....	84
Nitrofurantoin.....	81
Nitrofurantoin Macrocrystals .....	81
Nitrofurantoin Monohydrate .....	81
Nitroglycerin .....	241
Norditropin Flexpro.....	69
Norditropin Nordiflex Pen .....	69
Normosol-r .....	241
Normosol-r In D5w .....	241
Northera.....	137
Northera (s) .....	137
Noxafil.....	138
Noxafil (s) .....	138
Nplate .....	139
Nplate (s).....	139
Nulojix.....	241
Nutrilipid.....	241
Nutropin .....	69
Nutropin Aq.....	69
Nutropin Aq Pen.....	69
Nuvigil.....	140
Nuvigil (s) .....	140
<b>O</b>	
Octagam .....	104
Octreotide Acetate.....	171

Ofev.....	141
Ofev (s) .....	141
Olysio.....	142
Olysio (bcbs Ri) .....	142
Omnitrope .....	69
Ondansetron Hcl .....	241
Ondansetron Odt.....	241
Onfi .....	21
Opdivo.....	207
Opsumit.....	143
Opsumit (s).....	143
Orencia.....	144
Orencia (bcbs Ri) .....	144
Orenitram .....	145
Orenitram (s).....	145
Orphenadrine Citrate.....	89
Orphenadrine Citrate Er.....	89
Ortho-est .....	86
Otezla .....	146
Otezla (s).....	146
Oxaliplatin.....	241
Oxandrin (s) .....	147
Oxandrolone.....	147
Oxazepam .....	21
Oxycodone Hcl Er.....	148
Oxycontin.....	148
Oxycontin (s) .....	148
<b>P</b>	
Paclitaxel.....	241
Pantoprazole Sodium .....	241
Paricalcitol .....	241, 242
Part B Versus Part D.....	238
Pegintron .....	149
Peg-intron.....	149
Pegintron (s).....	149
Peg-intron Redipen .....	149
Peg-intron Redipen Pak 4 .....	149
Penicillin G Potassium In Iso-osmotic Dextrose .....	242
Pennsaid .....	150
Pennsaid (s).....	150
Pentam 300.....	242
Pentazocine/acetaminophen.....	76
Pentazocine/naloxone Hcl.....	76

Pentostatin.....	242	Promacta .....	156
Perforomist.....	242	Promacta (s).....	156
Perjeta.....	151	Promethazine Hcl.....	78
Perjeta (bcbs Ri).....	151	Promethazine Vc Plain.....	79
Perphenazine/amitriptyline.....	77	Promethegan .....	78
Phenadoz .....	78	Propranolol Hcl.....	242
Phenergan .....	78	Pulmicort.....	242
Piperacillin Sodium/ Tazobactam Sodium.....	242	Pulmozyme .....	157
Piperacillin Sodium/tazobactam Sodium .....	242	Pulmozyme (s) .....	157
Piperacillin/tazobactam .....	242		
Plasma-lyte A .....	242	<b>Q</b>	
Plasma-lyte-148.....	242	Qualaquin (s).....	158
Plasma-lyte-56/d5w.....	242	Quinine Sulfate .....	158
Plegridy .....	152		
Plegridy (s) .....	152	<b>R</b>	
Plegridy Starter Pack.....	152	Rabavert .....	242
Pomalyst.....	153	Rapamune .....	242
Pomalyst (s).....	153	Ravicti.....	159
Potassium Chloride .....	242	Ravicti (bcbs Ri).....	159
Potassium Chloride 0.15% /nacl 0.45% Viaflex.....	242	Rebif.....	160
Potassium Chloride 0.15% D5w/nacl 0.33% ....	242	Rebif (s) .....	160
Potassium Chloride 0.15% D5w/nacl 0.45% ....	242	Rebif Titration Pack.....	160
Potassium Chloride 0.15% D5w/nacl 0.45%		Reclast (bcbs Ri).....	161
Viaflex.....	242	Recombivax Hb .....	242
Potassium Chloride 0.15% W/nacl 0.9% Viaflex		Relistor.....	163
.....	242	Relistor (s).....	163
Potassium Chloride 0.15%/nacl 0.9% .....	242	Remicade.....	164
Potassium Chloride 0.22% D5w/nacl 0.45% ....	242	Remicade (bcbs Ri).....	164
Potassium Chloride 0.3%/ Nacl 0.9% .....	242	Remodulin.....	166
Potassium Chloride 0.3%/d5w .....	242	Remodulin (s).....	166
Prefest.....	86	Reserpine.....	80
Premarin .....	86	Retin-a Micro .....	200
Premasol .....	242	Revatio .....	167
Prempro .....	86	Revatio (s).....	167
Privigen .....	104	Revlimid.....	168
Procrit .....	50	Revlimid (s) .....	168
Procysbi.....	154	Rifampin .....	242
Procysbi (s).....	154	Ringers Injection.....	242
Prograf.....	242	Rituxan.....	242
Prolastin.....	242	Ruconest.....	169
Prolastin-c.....	242	Ruconest (bcbs Ri).....	169
Proleukin .....	242		
Prolia .....	155	<b>S</b>	
Prolia (bcbs Ri).....	155	Sabril .....	170
		Sabril (s).....	170

Saizen .....	69
Saizen Click.easy .....	69
Sandimmune.....	242
Sandostatin (s).....	171
Sandostatin Lar (s) .....	172
Sandostatin Lar Depot.....	172
Savella.....	173
Savella (s).....	173
Savella Titration Pack .....	173
Serostim.....	174
Serostim (bcbs Ri).....	174
Signifor.....	175
Signifor (s).....	175
Sildenafil .....	167
Simponi .....	176
Simponi (s) .....	176
Simponi Aria .....	177
Simponi Aria (s).....	177
Simulect.....	242
Simvastatin .....	178
Simvastatin (s).....	178
Sirolimus .....	242
Sodium Chloride .....	242
Sodium Chloride 0.45% Viaflex .....	242
Sodium Edecrin.....	242
Somatuline Depot.....	179
Somatuline Depot (s).....	179
Somavert.....	180
Somavert (s) .....	180
Sovaldi.....	181
Sovaldi (bcbs Ri).....	181
Sprycel.....	182
Sprycel (s) .....	182
Stelara.....	183
Stelara (bcbs Ri) .....	183
Stivarga.....	184
Stivarga (bcbs Ri).....	184
Suboxone.....	185
Suboxone (s).....	185
Subsys.....	58
Subutex (s).....	186
Sulfamethoxazole/trimethoprim.....	242
Surmontil.....	77
Sutent.....	187

Sutent (s).....	187
Sylatron.....	188
Sylatron (s).....	188
Sylvant .....	189
Sylvant (s).....	189
Synagis.....	190
Synagis (s).....	190
Synribo.....	191
Synribo (s).....	191

## **T**

Taclonex.....	192
Taclonex (bcbs Ri).....	192
Tacrolimus .....	242
Tafinlar.....	193
Tafinlar (s) .....	193
Talwin .....	76
Tarceva.....	194
Tarceva (s) .....	194
Targretin.....	195
Targretin (s) .....	195
Tasigna.....	196
Tasigna (s).....	196
Taxotere .....	242
Tazicef.....	242
Tazorac.....	197
Tazorac (s) .....	197
Tecfidera .....	198
Tecfidera (s).....	198
Tecfidera Starter Pack.....	198
Teflaro.....	242
Temazepam.....	21
Tetanus Toxoid Adsorbed.....	242
Tev-tropin .....	69
Thalomid.....	199
Thalomid (s).....	199
Thiotepa .....	242
Thymoglobulin.....	242
Ticlopidine Hcl .....	87
Tobi Podhaler.....	101
Tobramycin .....	101
Tobramycin Sulfate.....	242
Tobramycin Sulfate/sodium Chloride.....	242
Topical Retinoids (s).....	200
Toposar .....	242

Topotecan Hcl .....	242	Velcade .....	243
Torisel.....	242	Veletri (s) .....	209
Torseamide.....	242	Ventavis .....	210
Tracleer.....	201	Ventavis (bcbs Ri).....	210
Tracleer (s) .....	201	Verapamil Hcl.....	243
Tranexamic Acid.....	242	Vibativ.....	243
Travasol.....	242	Victrelis.....	211
Treanda.....	242	Victrelis(bcbs Ri) .....	211
Trelstar (s) .....	202	Vidaza (s) .....	212
Trelstar Depot Mixject .....	202	Viekira (s) .....	213
Trelstar La Mixject.....	202	Viekira Pak.....	213
Trelstar Mixject.....	202	Vinblastine Sulfate.....	243
Tretinoin .....	200	Vincasar Pfs .....	243
Tretinoin Microsphere.....	200	Vincristine Sulfate .....	243
Tretinoin Microsphere Pump .....	200	Vinorelbine Tartrate.....	243
Tretin-x.....	200	Virazole.....	243
Trexall .....	242	Voltaren.....	41
Triazolam .....	21	Votrient .....	214
Trihexyphenidyl Hcl .....	82	Votrient (s) .....	214
Trimethobenzamide Hcl.....	78	Vpriv .....	215
Trimipramine Maleate.....	77	Vpriv (bcbs Ri) .....	215
Trisenox.....	242	Vytorin .....	178
Trophamine .....	242		
Twinrix .....	242	<b>X</b>	
Tygacil.....	243	Xalkori .....	216
Tykerb .....	203	Xalkori (bcbs Ri).....	216
Tykerb (s) .....	203	Xeljanz .....	217
Tysabri.....	204	Xeljanz (s).....	217
Tysabri (s) .....	204	Xenazine .....	218
Tyvaso .....	205	Xenazine (s) .....	218
Tyvaso (s).....	205	Xeomin.....	219
Tyvaso Refill.....	205	Xeomin (s) .....	219
Tyvaso Starter .....	205	Xgeva .....	220
Tyzeka .....	206	Xgeva (bcbs Ri) .....	220
Tyzeka (s).....	206	Xifaxan.....	221
		Xifaxan (s) .....	221
<b>U</b>		Xolair .....	222
Universal Criteria .....	207	Xolair (s).....	222
		Xtandi.....	223
<b>V</b>		Xtandi (s) .....	223
Valchlor.....	208	Xyrem .....	224
Valchlor (s).....	208	Xyrem (bcbs Ri).....	224
Valproate Sodium.....	243		
Vancomycin Hcl.....	243	<b>Y</b>	
Vectibix .....	243	Yervoy.....	225

Yervoy (s).....	225
<b>Z</b>	
Zaleplon.....	88
Zaltrap .....	226
Zaltrap (s) .....	226
Zanosar .....	243
Zavesca.....	227
Zavesca (s).....	227
Zelboraf .....	228
Zelboraf (s).....	228
Zemplar .....	243
Ziana.....	200
Zohydro (bcbs Ri) .....	229
Zohydro Er .....	229
Zoledronic Acid.....	161, 231
Zolinza.....	230

Zolinza (s).....	230
Zolpidem Tartrate .....	88
Zolpidem Tartrate Er.....	88
Zometa .....	231
Zometa (s).....	231
Zorbtive.....	233
Zorbtive (bcbs Ri).....	233
Zortress .....	234
Zortress (s) .....	234
Zosyn.....	243
Zydelig .....	235
Zygelig (s).....	235
Zykadia .....	236
Zykadia (s) .....	236
Zytiga .....	237
Zytiga (s).....	237
Zyvox .....	243