

**EFFECTIVE DATE:** 05|23|2011

**POLICY LAST UPDATED:** 03|06|2018

## OVERVIEW

Intravenous (IV) infusion of lidocaine or ketamine has been investigated for the treatment of migraine and chronic daily headache, fibromyalgia, and chronic neuropathic pain. Chronic neuropathic pain disorders include phantom limb pain, postherpetic neuralgia, complex regional pain syndrome (CRPSs), diabetic neuropathy, and pain related to stroke or spinal cord injuries. IV infusion of ketamine has also been investigated for the treatment of depression and obsessive compulsive disorder. For this application, 1 or more courses of IV infusion would be administered over a period of several hours or several days.

## MEDICAL CRITERIA

Not applicable

## PRIOR AUTHORIZATION

Not applicable

## POLICY STATEMENT

### BlueCHiP for Medicare and Commercial Products

Intravenous infusion of anesthetics (e.g., ketamine or lidocaine) for the treatment of chronic pain including, but not limited to chronic neuropathic pain, chronic daily headache, and fibromyalgia, is considered not medically necessary, as the evidence is insufficient to determine the effects of the technology on health outcomes.

## COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate section of the Benefit Booklet, Evidence of Coverage or Subscriber Agreement for services not medically necessary.

## BACKGROUND

Courses of intravenous (IV) anesthetic agents may be given in the inpatient or outpatient setting as part of a pain management program, with the infusion of a subanesthetic dose preceded by a bolus infusion to achieve desired blood levels sooner.

Ketamine is an antagonist of the *N*-methyl-D-aspartate (NMDA) receptor and a dissociative anesthetic. It is the sole anesthetic agent approved for diagnostic and surgical procedures that do not require skeletal muscle relaxation. Respiratory depression may occur with overdosage or too rapid a rate of administration of ketamine; it should be used by or under the direction of physicians experienced in administering general anesthetics. Ketamine is a schedule III controlled substance. Psychological manifestations vary in severity from pleasant dream-like states to hallucinations and delirium, and can be accompanied by confusion, excitement, aggression, or irrational behavior. The occurrence of adverse events (AEs) with IV anesthetics may be reduced by the careful titration of subanesthetic doses. However, the potential benefits of pain control must be carefully weighed against the potential for serious, harmful adverse events

Lidocaine, which prevents neural depolarization through effects on voltage-dependent sodium channels, is also used systemically for the treatment of arrhythmias. Adverse events for lidocaine are common, can be mild to moderate, and include general fatigue, somnolence, dizziness, headache, periorbital and extremity numbness and tingling, nausea, vomiting, tremors, and changes in blood pressure and pulse. Severe adverse

effects may include arrhythmias, seizures, loss of consciousness, confusion, or even death. Lidocaine should only be given intravenously to patients with normal conduction on electrocardiography and normal serum electrolyte concentrations to minimize the risk of cardiac arrhythmias.

IV administration of anesthetic has been reported for various conditions, including chronic pain of neuropathic origin, chronic headache, fibromyalgia, depression, and obsessive compulsive disorders. Chronic daily headache is defined as a headache disorder that occurs more than 15 days a month for at least 3 months. Chronic daily headache includes chronic migraine, new daily persistent headache, hemicranias continua, and chronic tension-type headache.

Neuropathic pain is often disproportionate to the extent of the primary triggering injury and may consist of thermal or mechanical allodynia, dysesthesia, and/or hyperalgesia. Allodynia is pain that occurs from a stimulus that normally does not elicit a painful response (e.g., light touch, warmth). Dysesthesia is a constant or ongoing unpleasant or electrical sensation of pain. Hyperalgesia is an exaggerated response to normally painful stimuli. In the latter, symptoms may continue for a period of time that is longer (e.g.,  $\geq 6$  months) than clinically expected after an illness or injury. It is proposed that chronic neuropathic pain results from peripheral afferent sensitization, neurogenic inflammation, and sympathetic afferent coupling, along with sensitization and functional reorganization of the somatosensory, motor, and autonomic circuits in the central nervous system (CNS). Therefore, treatments focus on reducing activity and desensitizing pain pathways, thought to be mediated through NMDA receptors in the peripheral and central nervous system. Sympathetic ganglion blocks with lidocaine have been used for a number of years to treat sympathetically maintained chronic pain conditions, such as complex regional pain syndrome (previously known as reflex sympathetic dystrophy). Test infusion of an anesthetic has also been used in treatment planning to assess patient responsiveness to determine whether medications, such as oral mexiletine or oral ketamine, may be effective. A course of IV lidocaine or ketamine, usually at subanesthetic doses, has also been examined. This approach for treating chronic neuropathic pain differs from continuous subcutaneous or IV infusion of anesthetics for the management of chronic pain conditions, such as terminal cancer pain, which are not discussed herein.

Fibromyalgia is a chronic state of widespread pain and tenderness. Although fibromyalgia is generally considered to be a disorder of central pain processing or central sensitization, others have proposed that the nerve stimuli causing pain originates mainly in the muscle, causing both widespread pain and pain on movement. There are focal areas of hyperalgesia, or tender points, which tend to occur at muscle tendon junctions. Biochemical changes that have been associated with fibromyalgia include alterations in NMDA receptors, low levels of serotonin, suppression of dopamine-releasing neurons in the limbic system, dysfunction of the hypothalamic-pituitary-adrenal axis, and elevated substance P levels. Fibromyalgia is typically treated with neuropathic pain medications such as pregabalin, non-narcotic pain relievers, or low doses of antidepressants.

Use of IV ketamine has also been reported for treatment-resistant depression, defined as depression that does not respond adequately to appropriate courses of antidepressant medications. Particularly challenging are patients with treatment-resistant depression with suicidal ideation. Several studies are ongoing to test the efficacy of IV ketamine in patients with suicidal ideation who present to the emergency department.

The evidence for intravenous anesthetics in patients who have complex regional pain syndrome (CRPS), fibromyalgia, chronic headache, or other chronic neuropathic pain conditions includes several randomized controlled trials. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, quality of life, medication use, and treatment-related morbidity. Evidence, primarily from outside of the United States, suggests that IV courses of ketamine may provide at least temporary relief to some chronic pain patients. However, the intense treatment protocols, severity of adverse effects, and limited durability raises questions about the overall health benefit of this procedure. Additional clinical trials are

needed to evaluate the long-term safety of repeat courses of IV anesthetics. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have psychiatric disorders (eg, depression, obsessive-compulsive disorder) who receive a course of IV anesthetics (eg, lidocaine, ketamine), the evidence is limited. Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, quality of life, medication use, and treatment-related morbidity. Several trials on the IV infusion of ketamine for the treatment of suicidal ideation in patients with depression are ongoing. The evidence is insufficient to determine the effects of the technology on health outcomes.

### **Regulatory Status**

IV lidocaine is approved by the U.S. Food and Drug Administration (FDA) for systemic use in the acute treatment of arrhythmias and locally as an anesthetic. IV lidocaine for the treatment of chronic pain is an off-label use.

Ketamine hydrochloride injection is FDA-indicated for diagnostic and surgical procedures that do not require skeletal muscle relaxation, for the induction of anesthesia before the administration of other general anesthetic agents, and to supplement low-potency agents, such as nitrous oxide. IV ketamine for the treatment of chronic pain is an off-label use.

### **CODING**

#### **BlueCHiP for Medicare and Commercial Products**

There is no specific CPT code for this service; Claims should be filed using an unlisted CPT code for the treatments noted in this policy.

### **RELATED POLICIES**

None

### **PUBLISHED**

Provider Update, May 2018  
Provider Update, May 2017  
Provider Update, May 2016  
Provider Update, October 2015  
Provider Update, March 2013  
Provider Update, July 2011

### **REFERENCES**

1. Attal N, Gaude V, Brasseur L, et al. Intravenous lidocaine in central pain: a double-blind, placebo-controlled, psychophysical study. *Neurology*. Feb 8 2000; 54(3):564-574. PMID 10680784
2. Baranowski AP, De Courcey J, Bonello E. A trial of intravenous lidocaine on the pain and allodynia of postherpetic neuralgia. *J Pain Symptom Manage*. Jun 1999; 17(6):429-433. PMID 10388248
3. Kvarnstrom A, Karlsten R, Quiding H, et al. The effectiveness of intravenous ketamine and lidocaine on peripheral neuropathic pain. *Acta Anaesthesiol Scand*. Aug 2003; 47(7):868-877. PMID 12859309
4. Medrik-Goldberg T, Lifschitz D, Pud D, et al. Intravenous lidocaine, amantadine, and placebo in the treatment of sciatica: a double-blind, randomized, controlled study. *Reg Anesth Pain Med*. Nov-Dec 1999;24(6):534-540. PMID 10588558

5. Sorensen J, Bengtsson A, Ahlner J, et al. Fibromyalgia--are there different mechanisms in the processing of pain? A double blind crossover comparison of analgesic drugs. *J Rheumatol*. Aug 1997;24(8):1615-1621. PMID 9263160
6. Wallace MS, Ridgeway BM, Leung AY, et al. Concentration-effect relationship of intravenous lidocaine on the allodynia of complex regional pain syndrome types I and II. *Anesthesiology*. Jan 2000;92(1):75-83. PMID 10638902
7. Wu CL, Tella P, Staats PS, et al. Analgesic effects of intravenous lidocaine and morphine on postamputation pain: a randomized double-blind, active placebo-controlled, crossover trial. *Anesthesiology*. Apr 2002; 96(4):841- 848. PMID 11964590
8. Finnerup NB, Biering-Sorensen F, Johannesen IL, et al. Intravenous lidocaine relieves spinal cord injury pain: a randomized controlled trial. *Anesthesiology*. May 2005;102(5):1023-1030. PMID 15851891
9. Vlaimich R, Issy AM, Sakata RK. Effect of intravenous lidocaine associated with amitriptyline on pain relief and plasma serotonin, norepinephrine, and dopamine concentrations in fibromyalgia. *Clin J Pain*. May 2011;27(4):285-288. PMID 21178598
10. Eichenberger U, Neff F, Svetcic G, et al. Chronic phantom limb pain: the effects of calcitonin, ketamine, and their combination on pain and sensory thresholds. *Anesth Analg*. Apr 2008; 106(4):1265-1273, table of contents. PMID 18349204
11. Wertli MM, Kessels AG, Perez RS, et al. Rational pain management in complex regional pain syndrome 1 (CRPS 1)--a network meta-analysis. *Pain Med*. Sep 2014;15(9):1575-1589. PMID 25234478

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

