

EFFECTIVE DATE: 10|01|2015
POLICY LAST UPDATED: 09|05|2017

OVERVIEW

Artificial pancreas device systems link a glucose monitor to an insulin infusion pump that automatically takes action (eg, suspends or adjusts insulin) based on the glucose monitor reading. These devices are proposed to improve glycemic control in patients with insulin-dependent diabetes, in particular control of nocturnal hypoglycemia.

MEDICAL CRITERIA

BlueCHiP for Medicare and Commercial Products

Use of a U.S. Food and Drug Administration (FDA)-approved artificial pancreas device system with a low glucose suspend feature may be considered **medically necessary** in patients with type 1 diabetes who meet all of the following criteria:

- Age 16 and older
- Type 1 diabetes
- Glycated hemoglobin value between 5.8% and 10.0%
- Used insulin pump therapy for more than 6 months
- At least 2 documented nocturnal hypoglycemic events in a 2-week period

PRIOR AUTHORIZATION

Prior authorization is required for BlueCHiP for Medicare and recommended for Commercial products and is obtained via the online tool for participating providers. See the Related Policies section.

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products

Use of an FDA-approved artificial pancreas device system with a low glucose suspend feature may be considered **medically necessary** in patients with type 1 diabetes when medical criteria are met.

Use of an artificial pancreas device system is considered **not medically necessary** in all other situations.

COVERAGE

Benefits may vary by groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable office visit benefits/coverage, Medical Equipment, Medical Supplies and Prosthetic Devices benefits/coverage and Diabetic equipment/supplies benefits/coverage.

BACKGROUND

Tight glucose control in patients with diabetes has been associated with improved health outcomes. The American Diabetes Association recommends a glycated hemoglobin (HbA1c) level below 7% for most patients. However, hypoglycemia, defined as plasma glucose below 70 mg/dL, may place a limit on the ability to achieve tighter glycemic control. Hypoglycemic events in adults range from mild to severe, based on a number of factors including the glucose nadir, presence of symptoms, and whether the episode can be self-treated or requires help for recovery.

Hypoglycemia affects many aspects of cognitive function, including attention, memory, and psychomotor and spatial ability. Severe hypoglycemia can cause serious morbidity affecting the central nervous system (e.g.,

coma, seizure, transient ischemic attack, stroke), heart (e.g., cardiac arrhythmia, myocardial ischemia, infarction), eye (e.g., vitreous hemorrhage, worsening of retinopathy), as well as cause hypothermia and accidents that may lead to injury. Fear of hypoglycemia symptoms can also cause decreased motivation to adhere strictly to intensive insulin treatment regimens.

According to the FDA, an artificial pancreas is a medical device that links a glucose monitor to an insulin infusion pump where the pump automatically reduces and increases subcutaneous insulin delivery according to measured subcutaneous glucose levels using a control algorithm. As control algorithms can vary significantly, there are a variety of artificial pancreas device systems currently under development. These systems span a wide range of designs from low glucose suspend (LGS) device systems to the more complex bihormonal control-to-target systems.

FDA has described 3 main categories of artificial pancreas device systems:

1. **Threshold Suspend Device System**

With threshold suspend device systems, also called low glucose suspend systems, the delivery of insulin is suspended for a set time when 2 glucose levels are below a specified low level indicating hypoglycemia

2. **Control-to-Range System**

With these systems, the patient sets his or her own insulin dosing within a specified range, but the artificial pancreas device system takes over if glucose levels reach outside that range (higher or lower). Patients using this type of system still need to check blood glucose levels and administer insulin as needed.

3. **Control-to-Target System**

With this type of device, the system aims to maintain glucose levels near a target level, such as 100 mg/dL. Control-to-target systems are automated and do not require participation of the user except for calibration of the continuous glucose monitoring system. Several device subtypes are being developed such as those that deliver insulin-only, bi-hormonal systems and hybrid systems.

For individuals who have type 1 diabetes who receive an artificial pancreas device system with a low-glucose suspend feature The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have type 1 diabetes who receive a hybrid closed-loop insulin delivery system, the evidence is insufficient to determine the effects of the technology on health outcomes.

The evidence is insufficient to support use of the FDA-approved artificial pancreas device system for any other clinical indication. No other artificial pancreas device system besides a LGS system is FDA-approved and marketed in the U.S, and therefore, all other types of artificial pancreas devices are considered not medically necessary.

CODING

BlueCHiP for Medicare and Commercial Products

The following codes are covered when medical criteria are met.

S1034 Artificial pancreas device system (eg, low glucose suspend [LGS] feature) including continuous glucose monitor, blood glucose device insulin pump and computer algorithm that communicates with all of the devices

S1036 Transmitter; external, for use with artificial pancreas device system

S1037 Receiver (monitor); external, for use with artificial pancreas device system

RELATED POLICIES

Glucose Monitoring - Continuous

Preauthorization via Web-Based Tool for Durable Medical Equipment (DME)

PUBLISHED

Provider Update, November 2017
Provider Update, September 2016
Provider Update, July 2015
Provider Update, October 2014

REFERENCES

1. Trevitt S, Simpson S, Wood A. Artificial pancreas device systems for the closed-loop control of type 1 diabetes: what systems are in development? *J Diabetes Sci Technol*. May 2016;10(3):714-723. PMID 26589628
2. Food and Drug Administration (FDA). Types of Artificial Pancreas Device Systems. <http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/HomeHealthandConsumer/ConsumerProducts/ArtificialPancreas/ucm259555.htm>. Accessed October 21, 2016.
3. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Artificial Pancreas Device Systems. TEC Assessments. 2013;Volume 28:Tab 14.
4. Bergenstal RM, Klonoff DC, Garg SK, et al. Threshold-based insulin-pump interruption for reduction of hypoglycemia. *N Engl J Med*. Jul 18 2013;369(3):224-232. PMID 23789889
5. Garg S, Brazg RL, Bailey TS, et al. Reduction in duration of hypoglycemia by automatic suspension of insulin delivery: the in-clinic ASPIRE study. *Diabetes Technol Ther*. Mar 2012;14(3):205-209. PMID 22316089
6. Ly TT, Nicholas JA, Retterath A, et al. Effect of sensor-augmented insulin pump therapy and automated insulin suspension vs standard insulin pump therapy on hypoglycemia in patients with type 1 diabetes: a randomized clinical trial. *JAMA*. Sep 25 2013;310(12):1240-1247. PMID 24065010
7. Agrawal P, Zhong A, Welsh JB, et al. Retrospective analysis of the real-world use of the threshold suspend feature of sensor-augmented insulin pumps. *Diabetes Technol Ther*. May 2015;17(5):316-319. PMID 25611577
8. Bergenstal RM, Garg S, Weinzimer SA, et al. Safety of a hybrid closed-loop insulin delivery system in patients with type 1 diabetes. *JAMA*. Oct 4 2016;316(13):1407-1408. PMID 27629148
9. Phillip M, Battelino T, Atlas E, et al. Nocturnal glucose control with an artificial pancreas at a diabetes camp. *N Engl J Med*. Feb 28 2013;368(9):824-833. PMID 23445093
10. Nimri R, Atlas E, Ajzensztejn M, et al. Feasibility study of automated overnight closed-loop glucose control under MD-logic artificial pancreas in patients with type 1 diabetes: the DREAM Project. *Diabetes Technol Ther*. Aug 2012;14(8):728-735. PMID 22853723
11. Nimri R, Muller I, Atlas E, et al. MD-Logic overnight control for 6 weeks of home use in patients with type 1 diabetes: randomized crossover trial. *Diabetes Care*. Nov 2014;37(11):3025-3032. PMID 25078901

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

