

Medical Coverage Policy | Breast Ductal Lavage for Detection of Breast Cancer



EFFECTIVE DATE: 9/1/2001

POLICY LAST UPDATED: 7/10/2014

OVERVIEW

Ductal lavage is a method of collecting breast ductal epithelial cells for cytological analysis. It can be used as a risk assessment tool in women with a higher risk of breast cancer. It is sometimes called the "breast pap Smear".

PRIOR AUTHORIZATION

Not Applicable.

POLICY STATEMENT

BlueCHiP for Medicare and Commercial

Breast ductal lavage is considered not medically necessary as there is insufficient peer-reviewed scientific literature that demonstrates that the procedure/service is effective in comparison to more established techniques for acquisition and evaluation of breast cytology.

Fiberoptic ductoscopy is considered not medically necessary for the detection, diagnosis or treatment of breast cancer as the data is insufficient to permit scientific conclusion regarding the role of breast duct endoscopy in the evaluation and management of patients with known or suspected breast cancer.

MEDICAL CRITERIA

BlueCHiP for Medicare and Commercial

Not Applicable

BACKGROUND

All ductal and lobular breast cancers originate in a single layer of epithelial cells that line the ductal/lobular system of all milk ducts. Ductal lavage enables the retrieval of these cells using a microcatheter inserted into the milk ducts through the nipple orifices. A saline solution is flushed through the catheter into the ducts to wash out cells for cytological examination. The ductal lavage technique is directed at patients identified as being at high risk for breast cancer utilizing the Gail index, a personal history of breast cancer or evidence of BRCA mutation and have no mammographic abnormality.

The gold standard for examination of these cells has been nipple aspiration with cytopathological examination of the specimen. Ductal lavage is designed to harvest an increased number of cells with the ability to gather cells from individual milk ducts. Each breast has 6 to 8 milk ducts. The technique is based upon decades of research indicating that breast cancer originates in the epithelial lining of the milk ducts and involves a series of molecular changes from normal to abnormal to malignant. As long as the abnormal cells are contained within the ducts or lobules, they are termed preinvasive disease. Once they have invaded surrounding tissue, they are considered invasive cancer.

The procedure has been dubbed "breast pap smear" because like the test for cervical cancer, it is a nonsurgical approach to identifying abnormal cells prior to their development into cancer. Mammography is the standard for early detection for breast cancer. However, by the time an abnormality is detected via mammography, the lesion has grown to a size of 1 to 2 cm and may have been present for 6 to 8 years.

The value of cytopathological examination of specimens from the breast, especially from nipple aspiration are well documented. However, no studies utilizing cells acquired via ductal lavage have been reported. There is as yet no consensus among practicing cytopathologists as to the criteria for interpreting specimens obtained by ductal lavage. While the technique may be theoretically interesting and even promising in early studies, it has yet to be the subject of sufficient controlled studies to allow for a determination of its effectiveness, accuracy and safety in both an absolute sense and in comparison to more established techniques for acquisition and evaluation of breast cytology. It is not presently known how the technique will modify management of the high risk patient.

Fiberoptic ductoscopy is a technique that provides direct visual examination of the breast ducts through nipple orifice cannulation and exploration. It has been explored in the following clinical situations:

- Diagnostic technique in women with spontaneous nipple discharge (as potential alternative to surgical exploration)
- A follow-up to atypical cytology from ductal lavage specimen
- Delineation of ductal disease to define margins for surgical resection
- Direct delivery of therapeutic agents

There is minimal published information about how fiberoptic ductoscopy would be used in the management of the patients either in determining the need for other diagnostic tests, such as mammography or ductography, determining the need for biopsy or excision, or determining the extent of surgical excision. Although ductoscopy may be a useful technique for diagnosing DCIS prior to surgery, there is no data reporting on how the results of ductoscopy influences either the decision to undergo biopsy or excision or the extent of the excision. The data is insufficient to permit scientific conclusion regarding the role of breast duct endoscopy in the evaluation and management of patients with known or suspected breast cancer.

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Evidence of Coverage, Subscriber Agreement for applicable "not medically necessary" benefits/coverage.

CODING

BlueCHiP for Medicare and Commercial:

There is no specific CPT code for Breast Ductal Lavage or Fiberoptic Ductoscopy
19499 Unlisted procedure, breast

RELATED POLICIES

Not applicable

PUBLISHED

Provider Update,	Nov 2014
Provider Update,	Jan 2013
Provider Update,	Feb 2012
Provider Update,	Nov 2011
Provider Update,	Oct 2009
Provider Update,	Oct 2008
Policy Update,	Jul 2006

REFERENCES

Wood, ME., Stanley, MA., Crocker, AM., Kingsley, FS., Leiman, G. Ductal lavage of cancerous and unaffected

breasts: procedure success rate and cancer detection. *Acta Cytol.* 2009; 53(4): 410-5.

8. Golshan, M., Iglehart, D. Nipple Discharge. In: *UpToDate*, Basow, DS (Ed), Watham, MA, 2013.

National Cancer Institute (NCI). Breast Cancer Screening PDQ®. Last modified April 2013.

Available at:

<http://www.cancer.gov/cancertopics/pdq/screening/breast/healthprofessional/AllPages>.

Accessed

May 29, 2013.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Breast cancer screening and diagnosis. V1.2013.

Proctor KA, Rowe LR, Bentz JS. Cytologic features of nipple aspirate fluid using an automated non-invasive collection device: a prospective observational study. *BMC Women's Health.* 2005; 5:10.

Sauter ER, Klein-Szanto A, Macgibbon B, Ehya H. Nipple aspirate fluid and ductoscopy to detect breast cancer. *Diagn Cytopathol.* 2010 Apr;38(4):244-51.

Shen KW, Wu J, Lu JS, et al. Fiberoptic ductoscopy for breast cancer patients with nipple discharge. *Surg Endosc.* 2001; 15(11):1340-1345.

Shen KW, Wu J, Lu JS, et al. Fiberoptic ductoscopy for breast cancer patients with nipple discharge. *Surg Endosc.* 2001; 15(11):1340-1345.

Yamamoto D, Senzaki H, Nakagawa H, et al. Detection of chromosomal aneusomy by fluorescence in situ hybridization for patients with nipple discharge. *Cancer.* 2003; 97(3):690-6

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

