



EFFECTIVE DATE: 02|03|2015
POLICY LAST UPDATED: 11|07|2017

OVERVIEW

Bronchial thermoplasty is a potential treatment option for patients with severe persistent asthma. It consists of radiofrequency energy delivered to the distal airways with the aim of decreasing smooth muscle mass believed to be associated with airway inflammation.

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products

Bronchial thermoplasty for the treatment of asthma is considered **not medically necessary** as the evidence is insufficient to determine the effects of the technology on health outcomes.

MEDICAL CRITERIA

Not applicable

BACKGROUND

Bronchial thermoplasty is a proposed treatment option for patients with severe persistent asthma. The therapy is based on the premise that patients with asthma have an increased amount of smooth muscle in the airway and that contraction of this smooth muscle is a major cause of airway constriction. The thermal energy delivered via bronchial thermoplasty aims to reduce the amount of smooth muscle and thereby decrease muscle-mediated bronchoconstriction with the ultimate goal of reducing asthma-related morbidity. Bronchial thermoplasty is intended as a supplemental treatment for patients with severe persistent asthma (i.e., steps 5 and 6 in the stepwise approach to care).

Bronchial thermoplasty procedures are performed on an outpatient basis and each session lasts approximately 1 hour. During the procedure, a standard flexible bronchoscope is placed through the patient's mouth or nose into the most distal targeted airway and a catheter is inserted into the working channel of the bronchoscope. After placement, the electrode array in the top of the catheter is expanded, and radiofrequency energy is delivered from a proprietary controller and used to heat tissue to 65 degrees Centigrade over a 5 mm area. The positioning of the catheter and application of thermal energy is repeated several times in contiguous areas along the accessible length of the airway. At the end of the treatment session, the catheter and bronchoscope are removed. A course of treatment consists of 3 separate procedures in different regions of the lung scheduled about 3 weeks apart.

The uncertain degree of treatment benefit and the presence of substantial adverse events leave a large degree of uncertainty about the impact of bronchial thermoplasty on the net health outcome. As a result, bronchial thermoplasty is considered not medically necessary as a treatment for asthma.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for limitations of benefits/coverage when services are not medically necessary.

CODING

BlueCHIP for Medicare and Commercial Products

The following codes are not medically necessary:

31660 Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; with bronchial thermoplasty, 1 lobe

31661 Bronchoscopy, rigid or flexible, including fluoroscopic guidance, when performed; with bronchial thermoplasty, 2 or more lobes

RELATED POLICIES

Not applicable

PUBLISHED

Provider Update, January 2018

Provider Update, January 2017

Provider Update, April 2015

Provider Update, April 2014

Provider Update, February 2013

Provider Update, March 2012

REFERENCES

1. Centers for Disease Control and Prevention. Asthma FastStats. <http://www.cdc.gov/asthma/faststats.html>. Accessed May 12, 2016.
2. National Heart Lung and Blood Institute. Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma (EPR-3). 2007; <http://www.nhlbi.nih.gov/health-pro/guidelines/current/asthma-guidelines>. Accessed May 12, 2016.
3. Pavord ID, Cox G, Thomson NC, et al. Safety and efficacy of bronchial thermoplasty in symptomatic, severe asthma. *Am J Respir Crit Care Med*. Dec 15 2007;176(12):1185-1191. PMID 17901415
4. Pavord ID, Thomson NC, Niven RM, et al. Safety of bronchial thermoplasty in patients with severe refractory asthma. *Ann Allergy Asthma Immunol*. Nov 2013;111(5):402-407. PMID 24125149
5. Cox G, Thomson NC, Rubin AS, et al. Asthma control during the year after bronchial thermoplasty. *N Engl J Med*. Mar 29 2007;356(13):1327-1337. PMID 17392302
6. Thomson NC, Rubin AS, Niven RM, et al. Long-term (5 year) safety of bronchial thermoplasty: Asthma Intervention Research (AIR) trial. *BMC Pulm Med* 2011;11:8. PMID 21314924
7. Castro M, Rubin AS, Laviolette M, et al. Effectiveness and safety of bronchial thermoplasty in the treatment of severe asthma: a multicenter, randomized, double-blind, sham-controlled clinical trial. *Am J Respir Crit Care Med*. Jan 15 2010;181(2):116-124. PMID 19815809
8. Castro M, Rubin A, Laviolette M, et al. Persistence of effectiveness of bronchial thermoplasty in patients with severe asthma. *Ann Allergy Asthma Immunol*. Jul 2011;107(1):65-70. PMID 21704887
9. Wechsler ME, Laviolette M, Rubin AS, et al. Bronchial thermoplasty: Long-term safety and effectiveness in patients with severe persistent asthma. *J Allergy Clin Immunol*. Dec 2013;132(6):1295-1302. PMID 23998657
10. Zhou JP, Feng Y, Wang Q, et al. Long-term efficacy and safety of bronchial thermoplasty in patients with moderate-to-severe persistent asthma: a systemic review and meta-analysis. *J Asthma*. Feb 2016;53(1):94-100. PMID 26383773

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

