

Medical Coverage Policy | Home Prothrombin Time Monitoring



EFFECTIVE DATE: 04|01|2015

POLICY LAST UPDATED: 01|20|2015

OVERVIEW

Patients who are prescribed chronic warfarin anticoagulation need ongoing monitoring that has generally taken place in a physician's office or anticoagulation clinic. Home prothrombin monitoring with a U.S. Food and Drug Administration (FDA)-approved device is proposed as an alternative to office or laboratory-based testing.

PRIOR AUTHORIZATION

Not applicable.

POLICY STATEMENT

BlueCHiP for Medicare and Commercial

At home monitoring of chronic warfarin therapy may be considered medically necessary in patients who require continuous anticoagulation for chronic medical conditions. These conditions include, but are not limited to, patients with mechanical heart valves and chronic atrial fibrillation.

MEDICAL CRITERIA

Not applicable.

BACKGROUND

Warfarin is an effective anticoagulant for the treatment and prevention of venous and arterial thrombosis. Chronic warfarin therapy is recommended in all patients with mechanical heart valves and in some patients with chronic atrial fibrillation (i.e., patients with risk factors that indicate a higher likelihood of stroke). Patients with mechanical heart valves are frequently prescribed anticoagulants at higher levels than patients given anticoagulants for other indications, which puts them at higher risk of complications from warfarin therapy. Appropriate levels of warfarin anticoagulation are monitored with periodic prothrombin time measurements, as measured by the International Normalized Ratio (INR). For example, an INR result greater than 3 indicates a higher risk of serious hemorrhage, while an INR of 6 indicates an increased risk of developing a serious bleed nearly 7 times that of someone with an INR less than 3. In contrast, an INR less than 2 is associated with an increased risk of stroke. Therefore, monitoring of the prothrombin time is recommended to ensure that the prescribed dosing regimens result in INRs within the therapeutic range. Anticoagulation can be monitored: in the physician's office (usually once a month), at an anticoagulation clinic (usually once every 2 to 3 weeks), or at home.

In order for home prothrombin time monitoring to be effective, patients need to be appropriately trained and able to generate INR test results comparable to laboratory measures. Moreover, the clinical impact of home prothrombin time monitoring is related to improved warfarin management. Specifically, home prothrombin time monitoring permits more frequent monitoring and self-management of warfarin therapy with the ultimate goal of 1) increasing the time that the anticoagulation is within a therapeutic INR range (intermediate health outcome); and 2) decreasing the incidence of thromboembolic or hemorrhagic events (final health outcome). Home self-monitoring is typically associated with some form of self-management of warfarin therapy. In some cases, the patient may be supplied with treatment algorithms and instructed to alter the dose based on the results of self-monitoring. In other cases, the patient may be instructed to provide the results of the self-monitoring (e.g., on the telephone or internet) and receive instructions on warfarin dosage.

In January 2007, the CoaguChek® XS System (patient self-testing) (Roche Diagnostics Corporation) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA determined that this device was substantially equivalent to existing devices, including the CoaguChek SX System (professional, cleared in 2006). Other than a labeling change, the device is identical to the professional version of the CoaguChek XS System. The patient self-testing system is intended for self-monitoring of prothrombin time in patients who are on a stable regimen of anticoagulation medications. Other devices cleared by the FDA for home prothrombin time monitoring include the ProTime® Microcoagulation System (International Technidyne Corporation) and the Alere™ (formerly Hemosense) INRatio® 2 PT/INR Monitoring System.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable coverage/benefits:

- G0248 covered under applicable office visit coverage
- G0249 covered under applicable lab coverage
- G0250 covered under applicable physician services with no co-payment

CODING

BlueCHiP for Medicare and Commercial

The following codes are separately reimbursed services:

G0248, G0249

The following code is a separately reimbursed service and should only be reported by physicians:

G0250

ICD-9 and ICD-10 Diagnosis Codes that may support medical necessity:



Home Prothrombin
ICD 9 Codes that Sup



Home Prothrombin
ICD 10 Codes that Su

The following codes are intended to be used for laboratory services and physician interpretation of laboratory services only and are not for use with the G codes above. Is it not appropriate to use the codes below with home prothrombin time monitoring as the G codes are more specific. Anticoagulant services are intended to describe the outpatient management of warfarin therapy, including ordering, review and interpretation of INR testing, communication with patient and dosage adjustments as appropriate.

85610, 99363, 99364

RELATED POLICIES

Not applicable.

PUBLISHED

Provider Update	Apr	2015
Provider Update	Jun	2014
Provider Update	Apr	2013
Provider Update	Apr	2012
Provider Update	Sep	2011
Provider Update	Oct	2010
Provider Update	Oct	2009

REFERENCES

1. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) for Home PROTHROMBIN Time/International Normalized Ratio (PT/INR) Monitoring for Anticoagulation Management (190.11)
2. Garcia-Alamino JM, Ward AM, Alonso-Coello P et al. Self-monitoring and self-management of oral anticoagulation. *Cochrane Database Syst Rev* 2010; (4):CD003839.
3. Bloomfield HE, Krause A, Greer N et al. Meta-analysis: Effect of patient self-testing and self-management of long-term anticoagulation on major clinical outcomes. *Ann Intern Med* 2011; 154(7):472-82.
4. Heneghan C, Ward A, Perera R et al. Self-monitoring of oral anticoagulation: systematic review and meta-analysis of individual patient data. *Lancet* 2012; 379(813):322-34.
5. Matchar DB, Jacobson A, Dolor R et al. Effect of home testing of international normalized ratio on clinical events. *N Engl J Med* 2010; 363(17):1608-20.
6. Matchar DB, Jacobson AK, Edson RG et al. The impact of patient self-testing of prothrombin time for managing anticoagulation: rationale and design of VA cooperative study #481- the Home INR study (THINRS). *J Thromb Thrombolysis* 2005; 19(3):163-72.
7. Fitzmaurice DA, Murray ET, McCahon D et al. Self management of oral anticoagulation: randomised trial. *BMJ* 2005; 331 (7524):1057.
8. Menendez-Jandula B, Souto JC, Oliver A et al. Comparing self-management of oral anticoagulant therapy with clinic management: a randomized trial. *Ann Intern Med* 2005; 142(1):1-10.
9. Beyth RJ, Quinn L, Landefeld CS. A multicomponent intervention to prevent major bleeding complications in older patients receiving warfarin. *Ann Intern Med*; 2000; 133(9):687-95.
10. Kortke H, Korfer R. International normalized ratio self-management after mechanical heart valve replacement: is an early start advantageous? *Ann Thorac Surg* 2001; 72(1):44-8.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

