

**EFFECTIVE DATE:** 09|01|2015

**POLICY LAST UPDATED:** 06|26|2015

## OVERVIEW

Intensity-modulated radiotherapy (IMRT) may be an integral component in the treatment of cancers of the abdomen and pelvis. IMRT has been proposed as a method of radiotherapy that allows adequate radiotherapy to the tumor while minimizing the radiation dose to surrounding normal tissues and critical structures. IMRT versus 3-dimensional (3D) conformal radiation in chemoradiotherapy for anal cancer shows marked differences in rates of acute toxicity.

## MEDICAL CRITERIA

### BlueCHiP for Medicare and Commercial Products

Not applicable

## PRIOR AUTHORIZATION

Not applicable

## POLICY STATEMENT

### BlueCHiP for Medicare and Commercial Products

Intensity-modulated radiotherapy may be considered **medically necessary** as an approach to delivering radiotherapy for patients with cancer of the anus/anal canal when dosimetric planning with standard 3D conformal radiation predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity and for the treatment of cancer of the abdomen and pelvis including but not limited to the stomach, hepatobiliary tract, pancreas, rectal locations or gynecologic tumors.

IMRT would be considered **not medically necessary** for all other uses in the abdomen and pelvis as there is insufficient peer-reviewed literature that demonstrates that the procedure/service is effective.

## COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage, Subscriber Agreement, or Benefit Booklet for radiology benefit/coverage.

## BACKGROUND

### Radiation Techniques

#### Conventional External-Beam Radiotherapy

Over the past several decades, methods to plan and deliver radiotherapy have evolved in ways that permit more precise targeting of tumors with complex geometries. Most early trials used 2-dimensional treatment planning, based on flat images and radiation beams with cross-sections of uniform intensity that were sequentially aimed at the tumor along 2 or 3 intersecting axes. Collectively, these methods are termed conventional external-beam radiotherapy.

#### Three-Dimensional Conformal Radiation

Treatment planning evolved by using 3-dimensional images, usually from computed tomography (CT) scans, to delineate the boundaries of the tumor and discriminate tumor tissue from adjacent normal tissue and nearby organs at risk for radiation damage. Computer algorithms were developed to estimate cumulative radiation dose delivered to each volume of interest by summing the contribution from each shaped beam. Methods also were developed to position the patient and the radiation portal reproducibly for each fraction

and immobilize the patient, thus maintaining consistent beam axes across treatment sessions. Collectively, these methods are termed 3-dimensional conformal radiotherapy (3D-CRT).

### **Intensity-Modulated Radiotherapy**

IMRT, which uses computer software, CT images, and magnetic resonance imaging, offers better conformality than 3D-CRT because it is able to modulate the intensity of the overlapping radiation beams projected on the target and to use multiple-shaped treatment fields. It uses a device (a multileaf collimator [MLC]) that, coupled to a computer algorithm, allows for “inverse” treatment planning. The radiation oncologist delineates the target on each slice of a CT scan and specifies the target’s prescribed radiation dose, acceptable limits of dose heterogeneity within the target volume, adjacent normal tissue volumes to avoid, and acceptable dose limits within the normal tissues. Based on these parameters and a digitally reconstructed radiographic image of the tumor and surrounding tissues and organs at risk, computer software optimizes the location, shape, and intensities of the beam ports to achieve the treatment plan’s goals.

Increased conformality may permit escalated tumor doses without increasing normal tissue toxicity and thus may improve local tumor control, with decreased exposure to surrounding normal tissues, potentially reducing acute and late radiation toxicities. Better dose homogeneity within the target may also improve local tumor control by avoiding underdosing within the tumor and may decrease toxicity by avoiding overdosing.

Because most tumors move as patients breathe, dosimetry with stationary targets may not accurately reflect doses delivered within target volumes and adjacent tissues in patients. Furthermore, treatment planning and delivery are more complex, time-consuming, and labor-intensive for IMRT than for 3D-CRT. Thus, clinical studies must test whether IMRT improves tumor control or reduces acute and late toxicities when compared with 3D-CRT.

The body of evidence available to assess the role of IMRT in the treatment of cancers of the abdomen and pelvis generally comprises case series, both retrospective and prospective. Only 1 randomized trial has been reported that compared results of whole-pelvic IMRT with whole-pelvic conformal radiotherapy (CRT) for cervical cancer. Reports of case series, including concurrently treated control patients, are emerging. The available results are generally viewed as hypothesis-generating for the design and execution of comparative trials of IMRT versus CRT that evaluate tumor control and survival outcomes in the context of adverse events and safety.

The comparative data on use of IMRT versus 3-dimensional conformal radiation in chemoradiotherapy for anal cancer shows marked differences in rates of acute toxicity. Thus, use of IMRT in cancer of the anus/anal canal may be considered medically necessary.

For other tumors of the abdomen and pelvis, the evidence from treatment planning studies has shown that the use of IMRT decreases radiation doses delivered to normal tissue adjacent to tumor. This potentially lowers the risk of adverse events (acute and late effects of radiation toxicity), although the clinical benefit of reducing the radiation dose to normal tissue using IMRT is theoretical. Due to the limitations in this evidence, this policy underwent clinical vetting. There was support for the use of IMRT in tumors of the abdomen and pelvis when normal tissues would receive unacceptable doses of radiation. The results of the vetting, together with an indirect chain of evidence and the potential to reduce harms, led to the decision that IMRT may be considered medically necessary for the treatment of tumors of the abdomen and pelvis when dosimetric planning with standard 3D conformal radiation predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity.

### **CODING**

#### **BlueCHiP for Medicare and Commercial Products**

**A4648** Tissue marker, implantable, any type, each (Note: This code is not separately reimbursed for institutional providers.)

**Note:** To ensure correct pricing of HCPC code **A4648** for the Calypso 4D localization system, the procedure/clinical notes and the invoice must be submitted.

The following codes are covered for BlueCHiP for Medicare and commercial products when billed with the diagnosis codes listed in the attachments below:

### **Intensity-modulated radiation therapy**

**77301**

**77338**

**77385**

**77386**

**G6015:** Intensity modulated treatment delivery, single or multiple fields/arcs, via narrow spatially and temporally modulated beams, binary, dynamic MLC, per treatment session (effective 1/1/2015)

**G6016:** Compensator-based beam modulation treatment delivery of inverse planned treatment using 3 or more high resolution (milled or cast) compensator, convergent beam modulated fields, per treatment session (effective 1/1/2015)



ICD9: ICD9 IMRT abd  
pelvis 100115.xlsx



ICD10: ICD10 IMRT abd  
and pelvis 100115.xls

### **RELATED POLICIES**

None

### **PUBLISHED**

Provider Update, November 2015

Provider Update, October 2015

Provider Update, August 2014

Provider Update, April 2012

Provider Update, September 2011

Provider Update, January 2010

### **REFERENCES**

1. Veldeman L, Madani I, Hulstaert F et al. Evidence behind use of intensity-modulated radiotherapy: a systematic review of comparative clinical studies. *Lancet Oncol* 2008; 9(4):367-75.
2. Meyer JJ, Czito BG, Willett CG. Intensity-modulated radiation therapy for gastrointestinal tumors. *Curr Oncol Rep* 2008; 10(3):206-11.
3. Randall ME, Ibbott GS. Intensity-modulated radiation therapy for gynecologic cancers: pitfalls, hazards, and cautions to be considered. *Semin Radiat Oncol* 2006; 16(3):138-43.
4. Taylor A, Powell ME. Conformal and intensity-modulated radiotherapy for cervical cancer. *Clin Oncol (R Coll Radiol)* 2008; 20(6):417-25.
5. Staffurth J. A review of the clinical evidence for intensity-modulated radiotherapy. *Clin Oncol (R Coll Radiol)* 2010; 22(8):643-57.

6. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Anal Carcinoma V.1.2015. [http://www.nccn.org/professionals/physician\\_gls/pdf/anal.pdf](http://www.nccn.org/professionals/physician_gls/pdf/anal.pdf). Accessed November 12, 2014.
7. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Gastric Cancer. V.1.2014. [http://www.nccn.org/professionals/physician\\_gls/pdf/gastric.pdf](http://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf). Accessed November 12, 2014.
8. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Hepatobiliary Cancers. V.2.2014. [http://www.nccn.org/professionals/physician\\_gls/PDF/hepatobiliary.pdf](http://www.nccn.org/professionals/physician_gls/PDF/hepatobiliary.pdf). Accessed November 12, 2014.
9. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Pancreatic Adenocarcinoma. V.2.2014 [http://www.nccn.org/professionals/physician\\_gls/pdf/pancreatic.pdf](http://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf). Accessed November 12, 2014
10. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Rectal Cancer. V.1.2015. [http://www.nccn.org/professionals/physician\\_gls/pdf/rectal.pdf](http://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf). Accessed November 14, 2014.
11. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Cervical Cancer. V.2.2015. [http://www.nccn.org/professionals/physician\\_gls/PDF/cervical.pdf](http://www.nccn.org/professionals/physician_gls/PDF/cervical.pdf). Accessed November 12, 2014.
12. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Uterine Neoplasms. V.2.2015. [http://www.nccn.org/professionals/physician\\_gls/PDF/uterine.pdf](http://www.nccn.org/professionals/physician_gls/PDF/uterine.pdf). Accessed November 12, 2014.
13. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology. Ovarian Cancer. V.3.2014. [http://www.nccn.org/professionals/physician\\_gls/pdf/ovarian.pdf](http://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf). Accessed November 14, 2014.
14. Expert Panel on Radiation Oncology-Rectal/Anal C, Hong TS, Pretz JL, et al. ACR Appropriateness Criteria(R)
15. Anal Cancer. *Gastrointest Cancer Res.* Jan 2014;7(1):4-14. PMID 24558509

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge



are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.