

EFFECTIVE DATE: 10|01|2004

POLICY LAST UPDATED: 12|06|2016

OVERVIEW

This policy addresses a variety of minimally invasive techniques that have been investigated over the years as treatment of low back pain related to disc disease. Techniques can be broadly divided into techniques that are designed to remove or ablate disc material, and thus decompress the disc, and those designed to alter the biomechanics of the disc annulus.

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products

Percutaneous annuloplasty (e.g., intradiscal electrothermal annuloplasty, percutaneous intradiscal radiofrequency thermocoagulation, and intradiscal biacuplasty), disc nucleoplasty, laser discectomy, and automated percutaneous and endoscopic discectomies are considered not medically necessary as there is insufficient peer-reviewed scientific literature that demonstrates that the procedure/service is effective.

MEDICAL CRITERIA

Not applicable

BACKGROUND

Intradiscal annuloplasty therapies use energy sources to thermally treat discogenic low back pain arising from annular tears. Thermal annuloplasty techniques are designed to decrease pain arising from the annulus and enhance its structural integrity.

It has been proposed that heat-induced denaturation of collagen fibers in the annular lamellae may stabilize the disc and potentially seal annular fissures and that pain reduction may occur through the thermal coagulation of nociceptors in the outer annulus. With the intradiscal electrothermal annuloplasty procedure (IDET™, Oratec SpineCath System), a navigable catheter with an embedded thermal resistive coil is inserted posterolaterally into the disc annulus or nucleus. The catheter is then snaked through the disc circuitously to return posteriorly. Using indirect radiofrequency energy, electrothermal heat is generated within the thermal resistive coil at a temperature of 90 degrees C; the disc material is heated for up to 20 minutes. Proposed advantages of indirect electrothermal delivery of radiofrequency energy with IDET™ include precise temperature feedback and control and the ability to provide electrothermocoagulation to a broader tissue segment than would be allowed with a direct radiofrequency needle.

Another procedure, referred to as percutaneous intradiscal radiofrequency thermocoagulation (PIRFT), uses direct application of radiofrequency energy. With PIRFT, the radiofrequency probe is placed into the center of the disc, and the device is activated for only 90 seconds at a temperature of 70 degrees C. The procedure is not designed to coagulate, burn, or ablate tissue. The Radionics RF Disc Catheter System has been specifically designed for this purpose.

A more recently developed annuloplasty procedure, referred to as intradiscal biacuplasty (Baylis Medical, Inc., Montreal, Canada) involves the use of 2 cooled radiofrequency electrodes placed on the posterolateral sides

of the intervertebral annulus fibrosus. It is believed that by cooling the probes, a larger area may be treated than could occur with a regular needle probe.

Annuloplasty using a laser-assisted spinal endoscopy kit to coagulate the disc granulation tissue (percutaneous endoscopic laser annuloplasty) has also been described.

Laser energy (laser discectomy) and radiofrequency coblation (nucleoplasty) are being evaluated for decompression of the intervertebral disc. For laser discectomy under fluoroscopic guidance, a needle or catheter is inserted into the disc nucleus, and a laser beam is directed through it to vaporize tissue. For DISC nucleoplasty™, bipolar radiofrequency energy is directed into the disc to ablate tissue.

Patients considered candidates for DISC nucleoplasty™ or laser discectomy include patients with bulging discs and sciatica. A variety of different lasers have been investigated for laser discectomy, including YAG, KTP, holmium, argon, and carbon dioxide lasers. Due to differences in absorption, the energy requirements and the rate of application differ among the lasers. In addition, it is unknown how much disc material must be removed to achieve decompression. Therefore, protocols vary according to the length of treatment, but typically the laser is activated for brief periods only.

The Disc nucleoplasty™ procedure uses bipolar radiofrequency energy in a process referred to as coblation technology. The technique consists of small, multiple electrodes that emit a fraction of the energy required by traditional radiofrequency energy systems. The result is that a portion of nucleus tissue is ablated, not with heat but with a low-temperature plasma field of ionized particles. These particles have sufficient energy to break organic molecular bonds within tissue, creating small channels in the disc. The proposed advantage of this coblation technology is that the procedure provides for a controlled and highly localized ablation, resulting in minimal therapy damage to surrounding tissue.

Traditionally, discectomy and microdiscectomy are performed manually through an open incision. Percutaneous discectomy describes techniques by which disc decompression is accomplished by the physical removal of disc material rather than its ablation. These techniques have been modified by the use of automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Removal of disc herniations under endoscopic visualization is also being investigated. Back pain or radiculopathy related to herniated discs is an extremely common condition and a frequent cause of chronic disability. Although many cases of acute low back pain and radiculopathy will resolve with conservative care, a surgical decompression is often considered when the pain is unimproved after several months and is clearly neuropathic in origin, resulting from irritation of the nerve roots. Open surgical treatment typically consists of discectomy in which the extruding disc material is excised. When performed with an operating microscope, the procedure is known as microdiscectomy.

Minimally invasive options have also been researched, in which some portion of the disc material is removed or ablated, although these techniques are not precisely targeted at the offending extruding disc material.

Traditional discectomy has been modified by automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Endoscopic techniques may be intradiscal or may involve the extraction of non-contained and sequestered disc fragments from inside the spinal canal using an interlaminar or transforaminal approach. Following insertion of the endoscope, the decompression is performed under visual control.

There is limited evidence on the efficacy of intradiscal thermal annuloplasty, consisting of a small number of randomized controlled trials (RCTs) and case series. The two randomized controlled trials on intradiscal electrothermal annuloplasty report different results, with one reporting benefit for IDET and the other reporting no benefit. There is a lack of evidence to support a role for radiofrequency annuloplasty with a single probe. One recent RCT on biacuplasty suggests that this procedure may provide modest benefit in a

proportion of highly selected patients; confirmation of these results in a broader population is needed. Overall, evidence is insufficient to conclude that these procedures improve health outcomes. Therefore, annuloplasty (i.e., IDET™, PIRFT, and biacuplasty) is considered not medically necessary as there is no proven efficacy.

While numerous case series and uncontrolled studies report improvements in pain and functioning following laser discectomy and nucleoplasty, the lack of well-designed and conducted controlled trials limits interpretation of reported data. Questions remain about the safety and efficacy of these treatments. These procedures are considered not medically necessary as there is no proven efficacy.

Automated percutaneous discectomy involves placement of a probe within the intervertebral disc under image guidance with aspiration of disc material using a suction cutting device. There is insufficient evidence obtained from well-designed and executed randomized controlled trials to evaluate the impact of automated percutaneous discectomy on net health outcome. In addition, evidence from small randomized controlled trials does not support the use of these procedures; therefore, automated percutaneous discectomy is considered not medically necessary as there is no proven efficacy.

Endoscopic discectomy involves the percutaneous placement of a working channel under image guidance, followed by visualization of the working space and instruments through an endoscope. The evidence consists of a number of randomized controlled trials. At this time, evidence is considered insufficient to evaluate health outcomes from endoscopic discectomy in U.S. centers. Therefore, it is considered not medically necessary as there is no proven efficacy.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for limitations of benefits/coverage when services are not medically necessary.

CODING

BlueCHiP for Medicare and Commercial Products

The following codes are considered **not medically necessary**:

22526 22527 62287 62380 (New code effective 1/1/2017)
S2348

RELATED POLICIES

None

PUBLISHED

Provider Update, January 2017
Provider Update, April 2015
Provider Update, January 2014
Provider Update, September 2012
Provider Update, August 2010
Provider Update, November 2009
Provider Update, February 2009
Policy Update, February 2008

REFERENCES

1. Manchikanti L, Abdi S, Atluri S et al. An Update of Comprehensive Evidence-Based Guidelines for Interventional Techniques in Chronic Spinal Pain. Part II: Guidance and Recommendations. *Pain Physician* 2013; 16(2 Suppl):S49-S283.
2. Urrutia G, Kovacs F, Nishishinya MB et al. Percutaneous thermocoagulation intradiscal techniques for discogenic low back pain. *Spine (Phila Pa 1976)* 2007; 32(10):1146-54.

3. Helm S, Deer TR, Manchikanti L et al. Effectiveness of thermal annular procedures in treating discogenic low back pain. *Pain Physician* 2012; 15(3):E279-E304.
4. Pauza KJ, Howell S, Dreyfuss P et al. A randomized, placebo-controlled trial of intradiscal electrothermal therapy for the treatment of discogenic low back pain. *Spine J* 2004; 4(1):27-35.
5. Freeman BJ, Fraser RD, Cain CM et al. A randomized, double-blind, controlled trial: intradiscal electrothermal therapy versus placebo for the treatment of chronic discogenic low back pain. *Spine (Phila Pa 1976)* 2005; 30(21):2369-77; discussion 78.
6. Kvarstein G, Mawe L, Indahl A et al. A randomized double-blind controlled trial of intra-annular radiofrequency thermal disc therapy--a 12-month follow-up. *Pain* 2009; 145(3):279-86.
7. Freeman BJ, Mehdian R. Intradiscal electrothermal therapy, percutaneous discectomy, and nucleoplasty: what is the current evidence? *Curr Pain Headache Rep* 2008; 12(1):14-21.
8. Levin JH. Prospective, double-blind, randomized placebo-controlled trials in interventional spine: what the highest quality literature tells us. *Spine J* 2009; 9(8):690-703.
9. Helm S, Hayek SM, Benyamin RM et al. Systematic review of the effectiveness of thermal annular procedures in treating discogenic low back pain. *Pain Physician* 2009; 12(1):207-32.
10. Singh V, Manchikanti L, Benyamin RM et al. Percutaneous lumbar laser disc decompression: a systematic review of current evidence. *Pain Physician* 2009; 12(3):573-88.
11. Singh V, Manchikanti L, Calodney AK et al. Percutaneous lumbar laser disc decompression: an update of current evidence. *Pain Physician* 2013; 16(2 Suppl):SE229-60.
12. Gibson JN, Grant IC, Waddell G. Surgery for lumbar disc prolapse (Cochrane Review). *The Cochrane Library* 2003; Issue 2.
13. Gibson JN, Waddell G. Surgical interventions for lumbar disc prolapse. *Cochrane Database Syst Rev* 2007; (2):CD001350.
14. Hellinger PM. Nd-YAG (104nm) versus diode (940nm)PLDN: a prospective randomized blinded study. In: Brock M SW, Wille C, ed. *Proceedings from the first Interdisciplinary World Congress on Spinal Surgery and Related Disciplines* 2000:555-8.
15. Steffen R, Luetke A, Wittenberg RH et al. A prospective comparative study of chemonucleolysis and laser discectomy. *Orthop Trans* 1996; 20:388.
16. Goupille P, Mulleman D, Mammou S et al. Percutaneous laser disc decompression for the treatment of lumbar disc herniation: a review. *Semin Arthritis Rheum* 2007; 37(1):20-30.
17. Tassi GP. Comparison of results of 500 microdiscectomies and 500 percutaneous laser disc decompression procedures for lumbar disc herniation. *Photomed Laser Surg* 2006; 24(6):694-7.
18. Choy DS. Percutaneous laser disc decompression: an update. *Photomed Laser Surg* 2004; 22(5):393-406.
19. Menchetti PP, Canero G, Bini W. Percutaneous laser discectomy: experience and long term follow-up. *Acta Neurochir Suppl* 2011; 108:117-21
20. Dasenbrock HH, Juraschek SP, Schultz LR et al. The efficacy of minimally invasive discectomy compared with open discectomy: a meta-analysis of prospective randomized controlled trials. *J Neurosurg Spine* 2012; 16(5):452-62.
21. Arts MP, Brand R, van den Akker ME et al. Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial. *JAMA* 2009; 302(2):149-58.
22. Arts MP, Brand R, van den Akker ME et al. Tubular discectomy vs conventional microdiscectomy for the treatment of lumbar disk herniation: 2-year results of a double-blind randomized controlled trial. *Neurosurgery* 2011; 69(1):135-44; discussion 44.
23. Ryang YM, Oertel MF, Mayfrank L et al. Standard open microdiscectomy versus minimal access trocar microdiscectomy: results of a prospective randomized study. *Neurosurgery* 2008; 62(1):174-81; discussion 81-2.
24. Gibson JN, Grant IC, Waddell G. Surgery for lumbar disc prolapse. *Cochrane Database Syst*

Rev 2000; (3):CD001350.

25. Gibson JN, Waddell G. Surgical interventions for lumbar disc prolapse. Cochrane Database Syst Rev 2007; (2):CD001350.
26. Boswell MV, Trescot AM, Datta S et al. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. Pain Physician 2007; 10(1):7-111.
27. Freeman BJ, Mehdian R. Intradiscal electrothermal therapy, percutaneous discectomy, and nucleoplasty: what is the current evidence? Curr Pain Headache Rep 2008; 12(1):14-21.
28. Singh V, Benyamin RM, Datta S et al. Systematic review of percutaneous lumbar mechanical disc decompression utilizing Dekompressor. Pain Physician 2009; 12(3):589-99.
29. Vorobeychik Y, Gordin V, Fuzaylov D et al. Percutaneous mechanical disc decompression using Dekompressor device: an appraisal of the current literature. Pain Med 2012; 13(5):640-6.
30. Manchikanti L, Singh V, Falco FJ et al. An updated review of automated percutaneous mechanical lumbar discectomy for the contained herniated lumbar disc. Pain Physician 2013; 16(2 Suppl):SE151-84.
31. Manchikanti L, Singh V, Calodney AK et al. Percutaneous lumbar mechanical disc decompression utilizing Dekompressor(R): an update of current evidence. Pain Physician 2013; 16(2 suppl):SE1-24
32. Smith N, Masters J, Jensen C et al. Systematic review of microendoscopic discectomy for lumbar disc herniation. Eur Spine J 2013; 22(11):2458-65.
33. Centers for Medicare and Medicaid. NCD for Thermal Intradiscal Procedures (TIPs) (150.11). 2009. Available online at: <http://www.cms.gov/medicare-coverage-database/details/ncd->
34. Kapural L, Vrooman B, Sarwar S, et al. Radiofrequency intradiscal biacuplasty for treatment of discogenic lower back pain: a 12-month follow-up. Pain Med. Mar 2015;16(3):425-431. PMID 25339501
35. Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. Spine J. Jun 1 2015;15(6):1461-1477. PMID 24412033
36. Cong L, Zhu Y, Tu G. A meta-analysis of endoscopic discectomy versus open discectomy for symptomatic lumbar disk herniation. Eur Spine J. Jan 2016;25(1):134-143. PMID 25632840

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

