

**Blue Cross and Blue Shield of Rhode Island Preventive Services for Commercial Members January 2016**

**Note:** See separate policy for Preventive Services for Blue CHIP Members

Coverage for newly recommended preventive health services is made no later than one year after the release date of the recommendation.

Although many USPSTF recommendations are gender specific, BCBSRI does not limit coverage for gender specific services that are medically appropriate.

\* denotes an updated recommendation      \*\*denotes a new recommendation

**Abdominal Aortic Aneurysm Screening: men USPSTF Grade: B (June 2014)\***

USPSTF Recommends one-time screening for abdominal aortic aneurysm by ultrasonography in men ages 65 to 75 years who have ever smoked.

Health Service Code(s)	Diagnosis Code(s)	Frequency
G0369	No Specific Code	1 per lifetime

**Alcohol Misuse Screening and Counseling USPSTF Grade: B (May 2013)**

USPSTF recommends clinicians screen adults age 18 or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.

Health Service Code(s)	Diagnosis Code(s)	Frequency
99408, 99409	No Specific Code	1 per year of either code
99385-99387 99395-99397	No Specific Code	2 per year any combination of codes
99201-99215	ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	

**Anemia Screening USPSTF Grade: B (May 2006)**

USPSTF recommends routine screening for iron deficiency anemia in asymptomatic pregnant women.

Health Service Code(s)	Diagnosis Code(s)	Frequency
80055	No Specific Code	None

**Aspirin to Prevent Cardiovascular Disease USPSTF Grade: A (Mar 2009)**


USPSTF recommends the use of aspirin for men ages 45 to 79 when the potential benefit due to a reduction in myocardial infarctions outweighs the potential harm due to increase in gastrointestinal hemorrhage.

The USPSTF recommends the use of aspirin for women ages 55 to 79 years when the potential benefit of a reduction in ischemic strokes outweighs the potential harm of an increase in gastrointestinal hemorrhage.

Health Service Code(s)	Diagnosis Code(s)	Frequency
Pharmacy benefit with prescription Generic Only	No Specific Code	None

**Bacteriuria Screening: Pregnant Women USPSTF Grade: A (July 2008)**


USPSTF recommends screening for asymptomatic bacteriuria with urine culture in pregnant women at 12 to 16 weeks gestation or at the first prenatal visit, if later.


Health Service Code(s)	Diagnosis Code(s)	Frequency
87086 87088	ICD-9: V22.0 - V23.9 	None


**Blood Pressure Screening USPSTF Grade: A (Dec 2007)**

The USPSTF recommends screening for high blood pressure in adults age 18 years and older.

Health Service Code(s)	Diagnosis Code(s)	Frequency
99385-99387; 99395-99397	None	2 per year any combination of codes
99201-99215	ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	



<b>Blood Pressure Screening Bright Futures/American Academy of Pediatrics</b>		
<b>Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before 3 yrs.</b>		
<b>Health Service Code(s)</b>	<b>Diagnosis Code(s)</b>	<b>Frequency</b>
99381-99384 99391-99394 99201-99215	ICD-9: V20.1-V20.2 ICD-10: , Z00.121, Z00.129, Z76.2	Annual
<b>Psychosocial/Behaviorial Assessments for ages newborn to age 21 Bright Futures/American Academy of Pediatrics</b>		
<b>Health Service Code(s)</b>	<b>Diagnosis Code(s)</b>	<b>Frequency</b>
99381-99384 99391-99394 99201-99215	ICD-9: V20.1-V20.2 ICD-10: , Z00.121, Z00.129, Z76.2	Annual
<b>BRCA Risk Assessment and Genetic Counseling/Testing USPSTF Grade: B (Dec 2013)</b>		
<b>The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes (BRCA1 or BRCA2). Women with positive screening results should receive genetic counseling and, if indicated after counseling, BRCA testing</b>		
<b>Note: BRCA testing effective June 1, 2014 for no cost share.</b>		
<b>Health Service Code(s)</b>	<b>Diagnosis Code (s)</b>	<b>Frequency</b>
99385-99387 99397 99201-99215	No Specific Code ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	2 per year any combination of codes
96040	ICD-9: V16.3, V16.41 ICD-10: Z80.3, Z80.41	None
99401-99404	ICD-9: V16.3, V16.41 ICD-10: Z80.3, Z80.41	3 per year any code
81162, 81211, 81212, 81213, 81214 81215, 81216, 81217	BRCA Testing Requires Preauthorization	None
<b>Breast Cancer Preventive Medications USPSTF Grade: B (Sep 2013)</b>		
<b>The USPSTF recommends that clinicians engage in shared, informed decision making with women who are at increased risk for breast cancer about medications to reduce their risk. For women who are at increased risk for Breast Cancer and at low risk for adverse medication effects, clinicians should offer to prescribe risk reducing medications, such as tamoxifen or raloxifene.</b>		
<b>Health Service Code(s)</b>	<b>Diagnosis Code (s)</b>	<b>Frequency</b>
99385-99387 99395-99397 99201-99215	No Specific Code ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	2 per year any combination of codes
99401-99404	ICD-9: 217, 610 - 610.8, V16.3, V84.01  ICD-10 Codes Breast Cancer Prevention	3 per year any code

Breast Cancer Screening USPSTF Grade: B (Sep 2002)		
The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1 to 2 years for women age 40 years and older. (Plan policy age 40 and older)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
77055	No Specific Code	2 per year any combination of codes/pairs
77055 & 77051		
77056		
77056 & 77051		
77057		
77057 & 77052		
G0202 & 77052		
G0204 & 77051		
G0206		
G0206 & 77051		
G0202		
Breastfeeding support, supplies, and counseling USPSTF Grade: B (Oct 2008)		
The USPSTF recommends interventions during pregnancy and after birth to promote and support breastfeeding		
Note Breast Pumps: Hospital Grade Pumps are not covered as part of preventive services. Please see details of that policy for coverage guidelines		
Electric breast pumps are a contract exclusion and are not covered.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
E0602	No Specific Code Must have a prescription	With each birth
99381-99382; 99391-99392	No Specific Code	2 per year any combination of codes
99201-99215	ICD-9: V20.0, V20.1, V20.2, V20.3, V20.31, V20.32 ICD-10: Z00.110, Z00.111, Z00.121, Z00.129, Z76.1, Z76.2	
99401-99404	ICD-9: V24.1 ICD-10: Z39.1	None
98960	ICD-9: 676.00-676.04, 676.10-676.14, 676.20-676.24, 676.30-676.34, 676.40-676.44, 676.50-676.54, 676.80-676.84, 676.90-676.94 	1 in hospital outpatient clinic & 2 in physician office

Cervical Cancer Screening USPSTF Grade: A (Mar 2012)		
The USPSTF recommends screening for cervical cancer in women ages 21 to 65 years with cytology (Pap smear) every 3 years or, for women ages 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
88141	No Specific Code	1 per year any lab code
88142		
88143		
88147		
88148		
88150		
88152		
88153		
88154		
88164		
88165		
88166		
88167		
88174		
88175		
Exams		
99385-99387; 99395-99397	No Specific Code	2 exams any code every 3 years
99201-99215	ICD-9: V70.0, V72.31, V72.32 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419, Z01.42	
Q0091	ICD-9: V70.0, V72.31, V72.32 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	
Pathology	No Specific Code	2 any pathology code per year
P3000, P3001		
Chlamydia screening		
USPSTF Grade B (Sep 2014)*		
The USPSTF recommends screening for chlamydia in sexually active women age 24 years or younger and in older women who are increased risk for infection.		
Bright Futures American Academy of Pediatrics recommendation for STI/HIV screening		
Health Service Code(s)	Diagnosis Code(s)	Frequency
86631	ICD-9: V73.88	None
86632	V73.98 or	
87110	V22-V23.9	
87270	V70.0, V72.31	
87320	 ICD-10 Codes Chlamydia Screening	
87800		
87810		
87490		
87491		
Cholesterol Abnormalities Screening		
~Men younger than 35 at increased risk for coronary heart disease USPSTF Grade: B (Jun 2008)		
The USPSTF recommends screening men ages 20 to 35 years for lipid disorders if they are at increased risk for coronary artery disease.		
~Men 35 years and older USPSTF Grade: A (Jun 2008)		
The USPSTF strongly recommends screening men age 35 years and older for lipid disorders.		
~Women younger than 45 at increased risk for coronary heart disease USPSTF Grade: B (Jun 2008)		
The USPSTF recommends screening women ages 20 to 45 years for lipid disorders if they are at increased risk for coronary heart disease.		
~Women 45 years and older at increased risk for coronary heart disease USPSTF Grade A (Jun 2008)		
The USPSTF strongly recommends screening women age 45 years and older for lipid disorders if they are at increased risk for coronary heart disease.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
80061, 82465, 83718, 84478	No Specific Code	1 per year

Colorectal Cancer Screening USPSTF Grade: A (Oct 2008)		
<b>Colorectal Cancer Screening USPSTF Grade: A (Oct 2008)</b> <b>The USPSTF recommends screening for colorectal cancer using fecal occult blood testing, sigmoidoscopy or colonoscopy in adults beginning at age 50 years and continuing until age 75 years. The risks and benefits of these screening methods vary. (Plan policy ages 50-75)</b>		
Health Service Code(s)	Diagnosis Code(s)	Frequency
<b>Barium Enema, Double Contrast</b>		
74280	ICD-9: V76.41, V76.50, V76.51 ICD-10 :Z12.10, Z12.11, Z12.12	1 every 5 yrs.
<b>Fecal Occult Blood Test</b>		
82270	None	1 of any code listed per year
82274	ICD-9: V76.41, V76.50, V76.51 ICD-10: Z12.10, Z12.11, Z12.12	
<b>Colorectal Cancer Screening</b>		
Stool-based dna and fecal occult hemoglobin (e.g. kras, hdnrg4 and bump3) Note: This is the		
Cologuard Test		
81528	None	1 every 3 years
<b>Computed Tomographic (CT) Colography</b>		
74263	Requires Preauthorization by eviCore	1 every 5 years
<b>The above services are not part of Health Care Reform Mandate</b>		
<b>Sigmoidoscopy</b>		
45330, 45331, 45333, 45335, 45338, 45346	No Specific Code	1 of any code listed every 5 years
Effective 1/1/2015 CMS considers the following codes invalid for professional providers:. Use alternate code below		
45349, 45350		
Alternate Code:		
G0104		
<b>Colonoscopy</b>		
44388, 44389, 44392, 44394, 44401, 44402, 44403, 44404, 45378, 45380, 45381, 45384, 45385, 45388	No Specific Code	1 occurrence every 10 years
Effective 1/1/2015 CMS considers the following codes invalid for professional providers:. Use alternate code below.		1 occurrence every 5 years with increased risk factors such as a personal or family history in accordance with current American Cancer Society guidelines.
45390, 45398		
Alternate Code:		
G0105		
<b>Preventive when performed for a colorectal cancer screening:</b>		
Anesthesia		
00810-33	No Specific Code	None
Report modifier 33 for preventive colonoscopy		
Pathology		
88305-33	No Specific Code	None
Report modifier 33 for preventive colonoscopy		

Contraception Preventive Services for Women (Aug 2012)		
Note: These are contraceptive services/devices that are covered under the medical benefit.		
Pharmacy: Oral contraceptives (Tier 1) and injections, patches and vaginal ring are covered at the preventive rate		
Health Service Code(s)	Diagnosis Code(s)	Frequency
00851	No Specific Codes	None
11976, 11981, 57170,		
58300, 58301, 58600, 58605		
58611, 58615, 58565, 58670		
58671		
A4264		
J1050, J7297, J7298, J7300, J7301, J7303,		
J7304, J7306, J7307, S4981, S4989		
Critical Congenital Heart Defect Screening Bright Futures/American Academy of Pediatrics		
Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital per the 2011 AAP statement		
"Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease"		
Health Service Code(s)	Diagnosis Code(s)	Frequency
Covered as part of Inpatient Birth	None	None
Dental Caries Screening - Oral Health		
Bright Futures/American Academy of Pediatrics recommends a referral to a dental home if available, otherwise administer oral health risk assessment.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99381-99383 99393	99391 No Specific Code	2 per year any combination of codes
Dental Caries		
Prevention of Dental Caries in Children from Birth Through Age 5 Years USPSTF Grade: B (May 2014)		
The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation starting at age 6 months for children whose water supply is deficient in fluoride.		
The USPSTF recommends the application of fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption in primary care practices. (Plan policy for children under age 7)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
Oral Fluoride Supplementation Pharmacy with Prescription Generic Only 99188	No Specific Code  No Specific Code	  None
Depression Screening		
~Adolescents USPSTF Grade: B (Mar 2009)		
The USPSTF recommends screening adolescents (ages 12-18 years) for major depressive disorder when systems are in place to ensure accurate diagnosis, psychotherapy (cognitive-behavioral or interpersonal), and follow-up.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99384-99385 99394-99395	No Specific Code	2 per year any combination of codes
99201-99215	ICD-9: V70.0, V72.31	
	ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	
~Adults USPSTF Grade: B (Dec 2009)		
The USPSTF recommends screening adults for depression when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up.		
Health Service Code	Diagnosis Code	Frequency
99385-99387 99395-99397	No Specific Code	2 per year any combination of codes
99201-99215	ICD-9: V70.0, V72.31	
	ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	

Developmental and Autism Screening Bright Futures/American Academy of Pediatrics		
HRSA recommends that all children should be screened specifically for Autism Spectrum Disorders (ASD's) during well child doctor visits at: 18 months, 24 months (Plan policy up to 3 years of age for CPT code 96110)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
96110, 96127	No Specific Code	None
Diabetes Screening in Adults with High Blood Pressure USPSTF Grade: B (Jun 2008)		
The USPSTF recommends screening for type 2 diabetes in asymptomatic adults with sustained blood pressure (either treated or untreated) greater than 135/80 mm Hg.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
82947 83036	ICD-9: 401-405.9□	None
	 ICD-10 Codes Diabetes Screening	
Interpersonal and Domestic Violence Screening and Counseling		
Preventive Services for Women (Aug 2012)		
Health Service Code(s)	Diagnosis Code (s)	Frequency
99385-99387 99395-99397	No Specific Code	None
Dyslipidemia Screening Bright Futures/Academy of Pediatrics		
Screening for children at higher risk of lipid disorders		
Health Service Code(s)	Diagnosis Code(s)	Frequency
80061, 82465, 83718, 84478	No Specific Code	None
Falls Prevention in Older Adults		
~Exercise or Physical Therapy USPSTF Grade: B (May 2012)		
The USPSTF recommends exercise or physical therapy to prevent falls in community-dwelling adults age 65 years and older who are at increased risk for falls.		
Health Service Code(s)	Diagnosis Code (s)	Frequency
99387, 99397	None	None
99201-99215	ICD-9: V70.0 ICD-10: Z00.00, Z00.01	2 per year any combination of codes
~Vitamin D USPSTF Grade: B (May 2012)		
The USPSTF recommends vitamin D supplementation to prevent falls in community dwelling adults age 65 years and who are at increased risk for falls.		
Health Service Code	Diagnosis Code	Frequency
Pharmacy Benefit with Prescription Generic Only	None	None
Folic Acid Supplementation USPSTF Grade: A (May 2009)		
The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800ug) of folic acid.		
Health Service Code(s)	Diagnosis Code (s)	Frequency
Pharmacy Benefit with Prescription Generic Only	No Specific Code	None
Gestational Diabetes Mellitus Screening USPSTF Grade: B (Jan 2014)		
The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
82947, 82948, 82950, 82951, 82952	ICD-9: V22.0 - V23.9	None
	 ICD-10 Codes Gestational Diabetes	

Gonorrhea Prophylactic Ocular Topical Medication Newborns USPSTF Grade: A (July, 2011)		
The USPSTF recommends prophylactic ocular topical medication for all newborns for the prevention of gonococcal ophthalmia neonatorum.		
Health Service Code(s)	Diagnosis Code (s)	Frequency
Performed as part of delivery including home birth; is not separately reimbursed	N/A	N/A
Gonorrhea Screening USPSTF Grade: B (Sep 2014)*		
The USPSTF recommends screening for gonorrhea in sexually active women age 24 years or younger and in older women who are at increased risk for infection		
Bright Futures American Academy of Pediatrics recommendation for STI/HIV screening		
Health Service Code(s)	Diagnosis Code (s)	Frequency
87590, 87591, 87592 87850	No Specific Code	None
99385-99387 99395-99397	None	None
99201-99215	ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	2 per year any combination of codes
Healthy Diet and physical activity counseling to prevent cardiovascular disease: adults with cardiovascular risk factors USPSTF Grade B (Aug 2014)*		
The USPSTF recommends offering or referring adults who are overweight or obese and have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
97802, 97803, 97804, S9470, G0108, G0109 S9140, S9141, S9452 S9453, S9460, S9465 S9470	No Specific Code	None
99401-99404	ICD-9: V65.3, 278.00, 278.01, 278.02, V70.0, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z00.00, Z00.01, Z71.3	12 per year in combination with 99411-99412
99411-99412	ICD-9: V65.3, 278.00, 278.01, 278.02, V70.0, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z00.00, Z00.01, Z71.3	12 per year in combination with 99401-99404
Hearing Loss Screening: Newborns USPSTF Grade: B (July 2008)		
The USPSTF recommends screening for hearing loss in all newborn infants. (Plan policy up to age 1 year)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
92586, 92587	No Specific Code	1 per year either code
Hearing Screening Bright Futures/Academy of Pediatrics (Plan policy up to age 21)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
92551, 92558	ICD-9: V20.2 ICD-10: Z00.129	1 per year
Hemoglobinopathies (Sickle Cell) Screening: Newborns USPSTF Grade: A (Sep 2007)		
The USPSTF recommends screening for sickle cell disease in newborns. (Plan policy up to age 1 year)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
84999-32 83020 83021 S3620	No Specific Code	None




<b>Hematocrit or Hemoglobin Testing Bright Futures/Academy of Pediatrics</b>		
Universal screening for anemia should be performed at approximately 12 months of age with determination of Hb concentration and an assessment of risk factors associated with ID/IDA. (Plan policy up to 2 years of age)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
85014	All codes ICD-9: V78.0, V20.2 ICD-10: Z00.121, Z00.129, Z13.0	None
85018		
85025		
85027		
<b>Hepatitis B Virus Test Pregnant Women USPSTF Grade: A (Jun 2009)</b>		
The USPSTF strongly recommends screening for hepatitis B virus infection in pregnant women at their first prenatal visit.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
80055	No Specific Code	None
<b>Hepatitis B Virus Infection Screening for NonPregnant Adolescents and Adults</b>		
USPSTF Grade: B (May 2014)		
The USPSTF recommends screening for hepatitis B virus infection in persons at high risk for infection.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
86704, 86705, 86706, 86707	No Specific Code	None
<b>Hepatitis C Virus Screening: Adults USPSTF Grade: B (June 2013)</b>		
The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection.		
The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
86803, 87520-87522	ICD-9: V73.89 ICD-10: Z11.59	None
G0472	No Specific Code	None
<b>HIV Screening</b>		
~NonPregnant Adults and Adolescents USPSTF Grade: A (Apr 2013)		
The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults ages 15 to 65 years. Younger adolescents and older adults who are at increased risk should also be screened.		
~Pregnant Women USPSTF Grade: A (Apr 2013)		
The USPSTF recommends that clinicians screen all pregnant women for HIV, including those who present in labor who are untested and whose HIV status is unknown.		
<b>Bright Futures/American Academy of Pediatrics</b>		
Health Service Code(s)	Diagnosis Code(s)	Frequency
86689	No Specific Code	2 per year for 86689
86701, 86702, 86703, 87390, 87391, 87534, 87535, 87537, 87538, 87806, G0475		None for other codes
<b>Height, Weight and Body Mass Index Measurements</b>		
Bright Futures/American Academy of Pediatrics		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99381-99384	No Specific Code ICD-9: V20-V20.2 ICD-10: Z00.121, Z00129, Z76.1, Z76.2	None
99391-99394		
99201-99215		


Human Papilloma Virus (HPV) Screening Preventive Services for Women (Aug 2012)		
<b>High-risk human papillomavirus DNA testing in women with normal cytology results.</b>		
<b>(Plan policy ≥ 30 years old)</b>		
Health Service Code(s)	Diagnosis Code(s)	Frequency
87624, 87625, G0476	ICD-9: V73.81, V70.0, V76.2, V72.31, V76.47 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4, Z12.72	1 screening every 3 years
Hypothyroid Screening Newborns USPSTF Grade: A (Mar 2008)		
<b>The USPSTF recommends screening for congenital hypothyroidism in newborns. (Plan policy up to age 1 year)</b>		
Health Service Code(s)	Diagnosis Code(s)	Frequency
84999-32, S3620, 84436, 84437, 84439, 84443	No Specific Code	1 per year any code listed
Immunization Including Administration and Travel Immunization		
Health Service Code(s)	Diagnosis Code(s)	Frequency
<b>Administration Codes:</b>	No Specific Code	None
90460-90474		
<b>Vaccines</b>		
90476, 90477,		
90620, 90621, 90625, 90630		
90632, 90633, 90634,		
90636, 90644,		
90647, 90648,		
90649, 90650, 90651		
90654, 90655, 90656,		
90657, 90658, 90660,		
90661, 90662, 90664,		
90670, 90672,		
90673, 90680,		
90681, 90685,		
90686, 90687,		
90688, 90690, 90691,		
90692, 90696, 90698,		
90700, 90702,		
90707, 90710,		
90713, 90714,		
90715, 90716, 90717,		
90723, 90732, 90733,		
90734, 90736, 90738,		
90740, 90743, 90744,		
90746, 90747, 90748		
Intimate partner violence screening: USPSTF Grade: B (Jan 2013)		
<b>The USPSTF recommends that clinicians screen women of childbearing age for intimate partner violence, such as domestic violence, and provide or refer women who screen positive to intervention services.</b>		
<b>This recommendation applies to women who do not have signs or symptoms of abuse.</b>		
<b>This is part of Well-Visit or E/M Service</b>		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99384-99386	No Specific Code	None
99394-99396		
99201-99215	ICD-9: V70.0, V72.31 ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	2 per year any combination of codes

Health Service Code(s)	Diagnosis Code(s)	Frequency
<b>Iron Supplementation USPSTF Grade: B (May 2006)</b>		
The USPSTF recommends routine iron supplementation for asymptomatic children ages 6 to 12 months who are at increased risk for iron deficiency anemia. (Plan policy up to age 1 year)		
Pharmacy Benefit with Prescription Generic Only	No Specific Code	None
<b>Lead Testing Bright Futures/American Academy of Pediatrics ; CDC</b>		
Pediatricians should screen children at risk. To prevent lead poisoning, lead screening should begin at 9-12 months of age and be considered again at 24 months of age when BLL's peak. The CDC developed explicit guidance to state health departments for developing community screening policies. In communities where universal screening is recommended, pediatricians should follow this recommendation. In communities where targeted screening is recommended pediatricians should determine whether each young patient is at risk and screen when necessary. (Plan policy up to age 7 years)		
83655	No Specific Code	None
<b>Lung Cancer Screening USPSTF Grade: B (Dec 2013)</b>		
The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults ages 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery. (Plan policy ages 55 - 80 years).		
71250, S8032	ICD-9: V15.82, V69.8, V76.0, 305.1 ICD-10: F17.200, F17.201, F17.210, F17.211, Z12.2, Z72.0, Z87.891	1 screening per year
G0297	ICD-10 Z87.891 (ICD-9 V15.82) Screening requires preauthorization eviCore	
G0296 (not separately reimbursed; included in the preventive office visit)	No Specific Code	None
<b>Newborn Blood Screening Bright Futures/American Academy of Pediatrics</b>		
The Recommended Uniform Newborn Screening Panel as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations establish the criteria for coverage of newborn screening procedures and programs. Follow-up must be provided, as appropriate, by the pediatrician.		
84999-32	No Specific Code	None

Obesity Screening and Counseling		
~Adults USPSTF Grade: B (Jun 2012)		
The USPSTF recommends screening all adults for obesity. Clinicians should offer or refer patients with a body mass index of 30 kg/m2 or higher to intensive, multicomponent behavioral interventions.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
97802, 97803, 97804	No Specific Code	None
S9470, G0108		
G0109, S9140, S9141,		
S9452, S9455, S9460		
S9465, S9470		
G0473, G0447	No Specific Code	None
99385-99387	No Specific Code	
99395-99397		
99201-99215	ICD-9: V70.0, V72.31	2 per year any combination of codes
	ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	
99401-99404	ICD-9: V65.3, 278.00, 278.01, 278.02, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z71.3	12 per year in combination with 99411-12
99411-99412	ICD-9: V65.3, 278.00, 278.01, 278.02, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z71.3	12 per year in combination with 99401-99404
~Children USPSTF Grade: B (Jan 2010)		
The USPSTF recommends that clinicians screen children age 6 years and older for obesity and offer them or refer them to comprehensive intensive behavioral interventions to promote improvement in weight status		
Health Service Code(s)	Diagnosis Code(s)	Frequency
97802, 97803, 97804,	No Specific Code	None
S9470, G0108,		
G0109, S9140, S9141, S9452		
S9455, S9460, S9465		
S9470		
G0473, G0447	No Specific Code	None
99383-99385	No Specific Code	
99393-99395	ICD-9: V70.0, V72.31	2 per year any combination of codes
	ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	
99201-99215	ICD-9: V20.1, V20.2 ICD-10: Z00.121, Z00.129, Z76.2,	
99401-99404	ICD-9: V65.3, 278.00, 278.01, 278.02, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z71.3	12 per year in combination with 99411-99412
99411-99412	ICD-9: V65.3, 278.00, 278.01, 278.02, V77.8 ICD-10: E66.01, E66.02, E66.09, E66.1, E66.8, E66.9, Z71.3	12 per year in combination with 99401-99404
Osteoporosis Screening: Women USPSTF Grade: B (Jan 2012)		
The USPSTF recommends screening for osteoporosis in women age 65 years and older and in younger women whose fracture risk is equal to or greater than that of a 65-year-old white woman who has no additional risk factors		
Health Service Code(s)	Diagnosis Code(s)	Frequency
77080	ICD-9: V82.81 ICD-10: Z13.820	1 every 5 years

Phenylketonuria (PKU) Screening Newborns USPSTF Grade: B (Mar 2008)		
The USPSTF recommends screening for phenylketonuria in newborns. (Plan policy up to age 1 year)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
84030, 84999-32, S3620	No Specific code	None
Preeclampsia Prevention: Aspirin USPSTF Grade: B (Sep 2014)**New		
The USPSTF recommends the use of low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation in women who are at high risk for preeclampsia.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
Pharmacy benefit with prescription Generic Only	No Specific Code	None
Preventive		
~Well Baby Exam (Plan Policy birth to 30 months)		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99460, 99461, 99462, 99463	No Specific Code	Newborn Inpatient Codes
99381-99382, 99391-99292		11 visits 0 to 35 months
~Well Baby Exam (Plan policy over 30 Months)		
Health Service Code	Diagnosis Code	Frequency
99382-99287	No Specific Code	2 per year any combination of codes
99392-99297		
Prostate Specific Antigen (PSA) Testing RI State Mandate Not Part of Health Care Reform		
Health Service Code(s)	Diagnosis Code(s)	Frequency
84153	All codes ICD-9: V16,42, V70.0, V70.9, V76.44 ICD-10: Z00.00, Z00.01, Z00.8, Z12.5, Z80.42	1 per year
Rh(D) Incompatibility Screening		
~First Pregnancy Visit USPSTF Grade: A (Feb 2004)		
The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy related care.		
~24 - 28 Weeks Gestations USPSTF Grade: B (Feb 2004)		
The USPSTF recommends repeated Rh (D) antibody testing for all unsensitized Rh (D) -negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
80055, 86901	No Specific Code	None
Sexually Transmitted Infections Counseling USPSTF Grade: B (Sep 2014)*		
The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99384-99387 99394-99397	No Specific Code	2 per year any combination of codes
99201-99215		
	ICD-9: V70.0, V72.31	
	ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	
99401-99404	ICD-9:V65.44, V65.45, V69.2 ICD-10: Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	12 per year in combination with 99411-12
99411-99412	ICD-9:V65.44, V65.45, V69.2 ICD-10: Z71.7, Z71.89, Z72.51, Z72.52, Z72.53	12 per year in combination with 99401-99404

Health Service Code(s)	Diagnosis Code(s)	Frequency
<b>Skin Cancer Behavioral Counseling USPSTF Grade: B (May 2012)</b>		
The USPSTF recommends counseling children, adolescents, and young adults ages 10 to 24 years who have fair skin about minimizing their exposure to ultraviolet radiation to reduce risk for skin cancer.		
99383-99385, 99393-99395	No Specific Code	2 per year any combination of codes
99201-99215	ICD-9: V70.0, V20.32 ICD-10: Z00.00, Z00.01, Z00.111	
<b>Syphilis Infection Screening</b>		
~Non-Pregnant Persons USPSTF Grade: A (July 2004)		
The USPSTF strongly recommends that clinicians screen persons at increased risk for syphilis infection.		
~Pregnant Women USPSTF Grade: A (May 2009)		
The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.		
80055	None	None
86592-86593 86780	ICD-9: V22-V23.9, V69.2, V70.0, V74.5	
		
<b>Tobacco Use and Counseling and Interventions</b>		
~Non-Pregnant Adults USPSTF Grade: A (Apr 2009)		
The USPSTF recommends that clinicians ask all adults about tobacco use and provide tobacco cessation interventions for those who use tobacco products.		
~Pregnant Women USPSTF Grade: A (Apr 2009)		
The USPSTF recommends that clinicians ask all pregnant women about tobacco use and provide augmented, pregnancy-tailored counseling to those who smoke.		
99406, 99407	No Specific Code	None
99385-99387 99395-99397 99201-99215	ICD-9: 70.0, V72.31; ICD-10: Z00.00, Z00.01, Z01.411, Z01.419	2 per year any combination of codes
G0437		None
99411-99412 99441-99443	No Specific Code	None
Smoking Cessation Prescriptions and Over the Counter Drugs - Pharmacy Benefit with Prescription Generic Only unless unavailable		
~Children and Adolescents USPSTF Grade: B (Aug 2013)		
The USPSTF recommends that clinicians provide interventions, including education or brief counseling.		
99383-99385 99393-99395	No Specific Code	None
99201-99215	ICD-9: V20.2 ICD-10: Z00.121, Z00129	
<b>Tuberculin Test Bright Futures/Academy of Pediatrics; The Centers for Disease Control</b>		
Tuberculosis testing per recommendations of the Committee on Infectious Diseases, published in the current edition of AAP Red Book: Report of the Committee on Infectious Diseases. Testing should be performed on recognition of high risk factors ( <a href="#">Plan policy up to age 18 years</a> )		
86580	None	None
99211	ICD-9: V74.1, V01.1 ICD-10: Z11.1, Z20.1	

Urinary tract or other infection screening Preventive Services for Women (Aug 2012)		
For pregnant women part of prenatal care		
Health Service Code(s)	Diagnosis Code(s)	Frequency
81000, 81002, 81008, 87086, 87088	ICD-9: V22-V23.9	None
	 ICD-10 Codes Urinary tract or other infection	
Visual Acuity Screening in Children USPSTF Grade: B (Jan 2011)		
The USPSTF recommends vision screening for all children at least once between the ages of 3 and 5 years to detect the presence of amblyopia or its risk factors.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
(Plan Policy ages 3-21 years)		
99382-99383	No Specific Code	2 per year any combination of codes
99392-99393		
99201-99215	ICD-9: V20.1, V20.2 ICD-10: Z00.121, Z00129, Z76.2	
(Plan policy ages 3-5 years)		
99173	No Specific Code	Once per year
(Plan policy ages 0-12 months)		
99174	No Specific Code	2 per year
(Plan policy ages 1-5 years)		
99174	No Specific Code	Once per year
(99174 NOT covered on same day as 99173)		
Well Visits for Adults		
which includes Well-Woman Visits Preventive Service for Women (Aug 2012)		
Well-Woman Preventive Care Visit annually for adult women to obtain the recommended preventive services that are age and developmentally appropriate, including preconception care and many services necessary for prenatal care.		
Where appropriate, include other preventive services listed in the HRSA guidelines.		
Health Service Code(s)	Diagnosis Code(s)	Frequency
99385-99386	No Specific Code	2 per year any combination of codes
99395-99396		
99201-99215	ICD-9: V70.0 ICD10: Z00.00, Z00.01	