

Medical Coverage Policy | Thermal Capsulorrhaphy as a Treatment of Joint Instability



EFFECTIVE DATE: 02/03/2015

POLICY LAST UPDATED: 02/03/2015

OVERVIEW

Thermal capsulorrhaphy uses thermal energy to restructure collagen in the capsule or ligaments to reduce the capsule size. This procedure has primarily been evaluated for shoulder joint instability but may also be proposed to treat capsular laxity in other joints.

PRIOR AUTHORIZATION

Not applicable.

POLICY STATEMENT

BlueCHiP for Medicare and Commercial

Thermal capsulorrhaphy is considered **not medically necessary** as a treatment of joint instability, including, but not limited to the shoulder, knee, and elbow.

MEDICAL CRITERIA

Not applicable.

BACKGROUND

Shoulder instability may arise from a single traumatic event (i.e., subluxation or dislocation), repeated micro-trauma or constitutional ligament laxity, resulting in deformity and/or damage to the glenohumeral capsule and ligaments. Initial treatment of shoulder subluxation or dislocation is conservative in nature followed by range of motion and strengthening exercises. However, if instability persists, either activity modifications or surgical treatment may be considered.

Surgery consists of inspection of the shoulder joint with repair, reattachment, or tightening of the labrum, ligaments, or capsule, performed either with sutures or sutures attached to absorbable tacks or anchors. While arthroscopic approaches have been investigated over the past decade, their degree of success has been controversial due to a higher rate of recurrent instability compared with open techniques, thought to be related in part to the lack of restoration of capsular tension. Recent reports of arthroscopic techniques have described various suturing techniques for tightening the capsule, which require mastery of technically difficult arthroscopic intra-articular knot-tying.

Thermal capsulorrhaphy has been proposed as a technically simpler arthroscopic technique for tightening the capsule and ligaments. The technique is based on the observation that the use of nonablative levels of radiofrequency thermal energy can alter the collagen in the glenohumeral ligaments and/or capsule, resulting in their shrinkage and a decrease in capsular volume, both thought to restore capsular tension. Thermal capsulorrhaphy may be used in conjunction with arthroscopic repair of torn ligaments or other structures (ie, repair of Bankart or superior labrum anterior and posterior lesion). In addition, thermal capsulorrhaphy has also been investigated as an arthroscopic treatment of glenohumeral laxity, a common injury among overhead athletes, such as baseball players, resulting in internal impingement of the posterior rotator cuff against the glenoid labrum. Internal impingement is often accompanied by posterior rotator cuff tearing and labral injury. Thermal capsulorrhaphy has also been proposed as a sole arthroscopic treatment. For example, the technique may be considered in patients with chronic shoulder pain without recognized instability, based on the theory that the pain may be related to occult or microinstability. This diagnosis may be considered when a diagnostic arthroscopy reveals only lax ligaments and is commonly seen among baseball players. Finally, thermal

capsulorrhaphy may be considered in patients with congenital ligamentous laxity, such as Ehlers-Danlos or Marfan syndrome.

The literature does not support use of thermal capsulorrhaphy. The few available comparative studies do not support that this procedure is an efficacious treatment for shoulder instability. The case series report a high rate of unsatisfactory results and complications, raising the potential for a net harm. Because of the lack of efficacy and potential for harm, this procedure is considered not medically necessary.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for limitations of benefits/coverage when services are not medically necessary.

CODING

BlueCHiP for Medicare and Commercial

As there is no specific CPT code available for the use of thermal capsulorrhaphy in the ankles, knees, hip, wrist, or elbow the following unlisted code should be used.

29999

The following HCPCS code is considered **not medically necessary**:

S2300

RELATED POLICIES

Not applicable.

PUBLISHED

Provider Update Apr 2015

Provider Update Apr 2014

Provider Update Nov 2012

Provider Update Oct 2011

Provider Update Oct 2010

Provider Update Sep 2009

Provider Update Jul 2008

REFERENCES

1. Abrams JS. Thermal capsulorrhaphy for instability of the shoulder: concerns and applications of the heat probe. Instr Course Lect 2001; 50:29-36.
2. Gryler EC, Greis PE, Burks RT et al. Axillary nerve temperatures during radiofrequency capsulorrhaphy of the shoulder. Arthroscopy 2001; 17(6):567-72.
3. Mohtadi NG, Hollinshead RM, Ceponis PJ et al. A multi-centre randomized controlled trial comparing electrothermal arthroscopic capsulorrhaphy versus open inferior capsular shift for patients with shoulder instability: protocol implementation and interim performance: lessons learned from conducting a multi-centre RCT [ISRCTN68224911; NCT00251160]. Trials 2006; 7:4.
4. ClinicalTrials.gov. Electrothermal Arthroscopic Capsulorrhaphy (ETAC) and Open Inferior Capsular Shift in Patients With Shoulder Instability (NCT00251160). Available online at: <http://clinicaltrials.gov/ct2/show/NCT00251160?term=NCT00251160&rank=1>. Last accessed May 2012.
5. Levitz CL, Dugas J, Andrews JR. The use of arthroscopic thermal capsulorrhaphy to treat internal impingement in baseball players. Arthroscopy 2001; 17(6):573-7.
6. Savoie FH, 3rd, Field LD. Thermal versus suture treatment of symptomatic capsular laxity. Clin Sports Med 2000; 19(1):63-75, vi.

7. Chen S, Haen PS, Walton J et al. The effects of thermal capsular shrinkage on the outcomes of arthroscopic stabilization for primary anterior shoulder instability. *Am J Sports Med* 2005; 33(5):705-11.
8. Levy O, Wilson M, Williams H et al. Thermal capsular shrinkage for shoulder instability. Mid-term longitudinal outcome study. *J Bone Joint Surg Br* 2001; 83(5):640-5.
9. D'Alessandro DF, Bradley JP, Fleischli JE et al. Prospective evaluation of thermal capsulorrhaphy for shoulder instability: indications and results, two- to five-year follow-up. *Am J Sports Med* 2004; 32(1):21-33.
10. Levine WN, Bigliani LU, Ahmad CS. Thermal capsulorrhaphy. *Orthopedics* 2004; 27(8):823-6.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

