

Medical Coverage Policy | Transpupillary
Thermotherapy for Treatment of Choroidal
Neovascularization



EFFECTIVE DATE: 02|17|2009
POLICY LAST UPDATED: 10|20|2015

OVERVIEW

This policy documents the coverage determination for transpupillary thermotherapy (TTT) as it relates to the treatment of choroidal neovascularization (CNV) only. Transpupillary thermotherapy is a technique in which low-level heat is delivered through the pupil using a modified diode laser. TTT is designed to gently heat subfoveal choroidal lesions while limiting damage to the overlying retinal pigment epithelium.

MEDICAL CRITERIA

None

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products

Transpupillary thermotherapy for the treatment of choroidal neovascularization is not medically necessary as the clinical literature does not support its use.

COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for the applicable not medically necessary benefits/coverage.

BACKGROUND

Choroidal neovascularization is a common cause of adult-onset blindness, most commonly associated with age-related macular degeneration (AMD). In its earliest stages, AMD is characterized by minimal visual impairment and the presence of large drusen and other pigmentary abnormalities on ophthalmoscopic examination. As AMD progresses, 2 distinctively different forms of degeneration may be observed. The first, called the atrophic, areolar or dry form, evolves slowly. Atrophic AMD is the most common form of degeneration and is often a precursor of the second form, the more devastating exudative neovascular form, also referred to as disciform or wet degeneration. The wet form is distinguished from the atrophic form by serous or hemorrhagic detachment of the retinal pigment epithelium and the development of CNV, sometimes called neovascular membranes. Risk of developing severe irreversible loss of vision is greatly increased by the presence of CNV.

The pattern of CNV, as revealed by fluorescein or indocyanine angiography, is further categorized as classic or occult. For example, classic CNV appears as an initial lacy pattern of hyperfluorescence followed by more irregular patterns as the dye leaks into the subretinal space. Occult CNV lacks the characteristic angiographic pattern, either due to the opacity of coexisting subretinal hemorrhage or, especially in CNV associated with AMD, by a tendency for epithelial cells to proliferate and partially or completely surround the new vessels. Interestingly, lesions consisting only of classic CNV carry a worse visual prognosis than those composed of only occult CNV, suggesting that the proliferative response that obscures new vessels may also favorably alter the clinical course of AMD.

There is ongoing research interest in the use of TTT to treat subfoveal choroidal neovascularization with an “occult” angiographic pattern. TTT is a technique in which heat is delivered to the choroid and retinal

pigment epithelium through the pupil using a modified diode laser. This laser technique contrasts with the laser used in standard photocoagulation therapy in that TTT uses a lower power laser for more prolonged periods of time and is designed to gently heat the choroidal lesion, thus limiting damage to the overlying retinal pigment epithelium.

Transpupillary thermotherapy is a technique in which low-level heat is delivered through the pupil using a modified diode laser. TTT is designed to gently heat subfoveal choroidal lesions while limiting damage to the overlying retinal pigment epithelium. Evidence on TTT is limited. The available studies comparing TTT with sham have not shown a benefit of this procedure. Although trials comparing TTT to photodynamic therapy show similar outcomes for the 2 treatments, there may be an increase in adverse events with TTT. TTT has not been compared with angiogenesis inhibitors. Evidence is insufficient to determine whether TTT is as beneficial as the established alternative; this procedure is considered not medically necessary.

CODING

BlueCHiP for Medicare and Commercial Products

There is no specific code for this service. To report, use the unlisted procedure, posterior segment: **67299**

RELATED POLICIES

None

PUBLISHED

Provider Update, December 2015

Provider Update, October 2014

Provider Update, June 2012

Provider Update, October 2010

Provider Update, April 2009

REFERENCES

1. Reichel E, Berrocal AM, Ip M et al. Transpupillary thermotherapy of occult subfoveal choroidal neovascularization in patients with age-related macular degeneration. *Ophthalmology* 1999; 106(10):1908-14.
2. Newsom RS, McAlister JC, Saeed M et al. Transpupillary thermotherapy (TTT) for the treatment of choroidal neovascularisation. *Br J Ophthalmol* 2001; 85(2):173-8.
3. Gustavsson C, Agardh E. Transpupillary thermotherapy for occult subfoveal choroidal neovascularization: a 1-year, prospective randomized pilot study. *Acta Ophthalmol Scand* 2005; 83(2):148-53.
4. Myint K, Armbrecht AM, Mon S et al. Transpupillary thermotherapy for the treatment of occult CNV in age-related macular degeneration: a prospective randomized controlled pilot study. *Acta Ophthalmol Scand* 2006; 84(3):328-32.
5. Odergren A, Algvere PV, Seregard S et al. A prospective randomised study on low-dose transpupillary thermotherapy versus photodynamic therapy for neovascular age-related macular degeneration. *Br J Ophthalmol* 2008; 92(6):757-61.
6. Odergren A, Algvere PV, Seregard S et al. Vision-related function after low-dose transpupillary thermotherapy versus photodynamic therapy for neovascular age-related macular degeneration. *Acta Ophthalmol* 2010; 88(4):426-30.
7. Tewari HK, Prakash G, Azad RV et al. A pilot trial for comparison of photodynamic therapy and transpupillary thermotherapy for the management of classic subfoveal choroidal neovascularization secondary to age-related macular degeneration. *Indian J Ophthalmol* 2007; 55(4):277-81.

8. Zhang X, Zhu X, Wang D et al. Low-power transpupillary thermotherapy combined with intravitreal triamcinolone acetonide for subfoveal choroidal neovascularization. *Ophthalmic Res* 2007; 39(4):241-2.

9. Nowak MS, Jurowski P, Grzybowski A et al. A prospective study on different methods for the treatment of choroidal neovascularization. The efficacy of verteporfin photodynamic therapy, intravitreal bevacizumab and transpupillary thermotherapy in patients with neovascular age-related macular degeneration. *Med Sci Monit* 2012; 18(6):CR374-80.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

