Medical Coverage Policy | Tumor-Treatment Fields for

Glioblastoma

Blue Cross Blue Shield of Rhode Island

EFFECTIVE DATE: 1/20/2014 **POLICY LAST UPDATED:** 1/29/2015

OVERVIEW

Glioblastoma multiforme is the most common and deadly malignant brain tumor. It has a very poor prognosis and is associated with low quality of life during the course of treatment. Tumor-treatment fields (TTF) therapy, also known as Novocure, is a new, noninvasive technology that is intended to treat glioblastoma using electric fields.

PRIOR AUTHORIZATION

Not Applicable.

POLICY STATEMENT

BlueCHiP for Medicare and Commercial Products:

Tumor-Treatment Fields Therapy for the treatment of Glioblastoma is considered not medically necessary as there is insufficient peer reviewed scientific literature that demonstrates that the procedure/service is effective.

MEDICAL CRITERIA

None

BACKGROUND

Glioblastomas, also known as glioblastoma multiforme (GBM), are the most common form of malignant primary brain tumor in adults, and they comprise approximately 15% of all brain and central nervous system tumors and more than 50% of all tumors that arise from glial cells. The peak incidence for GBM occurs between the ages of 45 and 70 years. GBMs

are grade IV astrocytomas, the most deadly type of glial cell tumor, and are often resistant to standard chemotherapy. According to the National Comprehensive Cancer Network,

GBM is the "deadliest brain tumor with only a third of patients surviving for 1 year and less than 5% living beyond 5 years.

The primary treatment for GBM is debulking surgery to remove as much of the tumor as possible. At that time, some patients may undergo implantation of the tumor cavity with a carmustine (BCNU) -impregnated wafer. (2) Depending on the patient's physical condition, adjuvant radiation therapy, chemotherapy (typically temozolomide), or a combination of the two are sometimes given. After adjuvant therapy, some patients may undergo maintenance therapy with temozolomide. In patients with disease that recurs after these initial therapies, additional debulking surgery may be used if recurrence is localized. Treatment options for recurrent disease include various forms of systemic medications such as bevacizumab, bevacizumab plus chemotherapy (e.g., irinotecan, BCNU/CCNU, temozolomide), temozolomide, nitrosourea, PCV (procarbazine, CCNU, and vincristine), cyclophosphamide, and platinum-based agents. (2) Response rates in recurrent disease are less than 10%, and progression-free survival rates at 6 months are less than 20%. (2, 3)

Tumor-treatment fields (TTF) therapy is a new, noninvasive technology that is intended to treat GBM on an outpatient basis using electrical fields. (3-5) TTF therapy exposes cancer cells to alternating electric fields of low intensity and intermediate frequency, which are purported to both selectively inhibit tumor growth and

reduce tumor angiogenesis. Tumor-treatment fields are proposed to inhibit rapidly dividing tumor cells by two mechanisms, arrest of cell proliferation and destruction of cells while undergoing division.

The NovoTTF-100A[™] System (Novocure Ltd., Haifa, Israel) has been approved by the U.S. Food and Drug Administration (FDA) to deliver TTF therapy. TTF therapy via the NovoTTF-100A[™] System is delivered by a battery-powered, portable device that generates the fields via disposable electrodes that are noninvasively attached to the patient's shaved scalp over the site of the tumor. (3, 4) The device is used by the patient at home on a continuous basis (20–24 hours per day) for the duration of treatment, which can last for several months. Patients can carry the device in a backpack or shoulder pack while carrying out activities of daily living.

Further evidence from high-quality trials is needed to assess the long-term safety and efficacy of TTF therapy. There are currently ongoing clinical trials of the TTF therapy, including an ongoing postmarketing noninferiority study that will provide additional data on outcomes of interest. Based on the small amount of evidence and lack of demonstrated treatment benefit, the use of TTF therapy for glioblastoma is considered not medically necessary.

COVERAGE

Benefits vary between groups/contracts. Please refer to the appropriate Evidence of Coverage or Subscriber Agreement for applicable services not medically necessary coverage.

CODING

BlueCHiP for Medicare and Commercial

The following codes are not medically necessary

A4555: Electrode/transducer for use with electrical stimulation device used for cancer treatment, replacement only

E0766: Electrical stimulation device used for cancer treatment, includes all accessories, any type

RELATED POLICIES

None

PUBLI SHED

Provider Update, Apr 2015 Provider Update, Mar 2014

REFERENCES

1. Novo TTF-100A System

http://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/DeviceApprovalsandClearan ces/Recently-ApprovedDevices/ucm254480.htm

- 2. http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm251669.htm
- National Cancer Institute (NCI) Adult Brain Tumors Treatment (PDQ®). Last modified May 14, 2013. Available online at: http://www.cancer.gov/cancertopics/pdq/treatment/adultbrain/HealthProfessional. Last accessed June 2013.
- National Comprehensive Cancer Network (NCCN). Central nervous system cancers. NCCN Clinical Practice Guidelines in Oncology. Version 2.2013. For additional information visit the NCCN website at: http://www.nccn.org/index.asp. Last accessed June 2013.
- 5. Davies AM, Weinberg U, Palti Y. Tumor treating fields: a new frontier in cancer therapy. Ann N Y Acad Sci 2013.
- Wong ET, Lok E, Swanson KD, et al. Response assessment of NovoTTF-100A versus best physician's choice chemotherapy in recurrent glioblastoma. Cancer Med. Jun 2014;3(3):592-602. PMID 24574359

- Elzinga G, Wong ET. Resolution of cystic enhancement to add-on tumor treating electric fields for recurrent glioblastoma after incomplete response to bevacizumab. Case Rep Neurol. Jan 2014;6(1):109-115. PMID 24847254
- Turner SG, Gergel T, Wu H, et al. The effect of field strength on glioblastoma multiforme response in patients treated with the NovoTTF-100A system. World J Surg Oncol. 2014;12(1):162. PMID 24884522

----- CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

1