

## Medical Coverage Policy | Vertebral Fracture Assessment



**EFFECTIVE DATE:** 10|01|2015  
**POLICY LAST UPDATED:** 09|01|2015

### OVERVIEW

Vertebral fracture assessment (VFA) with densitometry (bone density study) is a technique in which vertebral fractures are assessed at the same time as bone mineral density (BMD), by use of dual-energy X-ray absorptiometry (DXA). The addition of vertebral fractures to BMD may provide additional useful information on a subject's risk of fracture.

This policy addresses VFA **without** densitometry.

### MEDICAL CRITERIA

Not applicable

### PRIOR AUTHORIZATION

Not applicable

### POLICY STATEMENT

#### Commercial Products

Screening for vertebral fractures using dual-energy X-ray absorptiometry (DXA or DEXA), **without bone density study**, is considered not medically necessary due to insufficient peer-reviewed literature that the service is effective.

#### BlueCHiP for Medicare

Screening for vertebral fractures using dual-energy X-ray absorptiometry (DXA or DEXA), **without bone density study**, is covered, but not separately reimbursed.

### COVERAGE

Benefits may vary between groups/contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage, or Subscriber Agreement for limitations of benefits/coverage when services are not medically necessary.

### BACKGROUND

Vertebral fractures are highly prevalent in the elderly population, and epidemiologic studies have found that these fractures are associated with an increased risk of future spine or hip fractures independent of BMD. Only 20% to 30% of vertebral fractures are recognized clinically; the rest are discovered incidentally on lateral spine radiographs. Lateral spine radiographs have not been recommended as a component of risk assessment for osteoporosis because of the cost, radiation exposure, and the fact that the radiograph would require a separate procedure in addition to the BMD study using DXA. However, several densitometers with specialized software are able to perform VFA in conjunction with DXA. The lateral spine scan is performed by using a rotating arm; depending on the densitometer used, the patient can either stay in the supine position after the bone density study or is required to move onto the left decubitus position.

VFA differs from radiologic detection of fractures, as VFA uses a lower radiation exposure and can detect only fractures, while traditional radiograph images can detect other bone and soft tissue abnormalities in addition to spinal fractures. Manufacturers have also referred to this procedure as instant vertebral

assessment, radiographic vertebral assessment, dual-energy vertebral assessment, or lateral vertebral assessment.

For both lateral spine radiographs and images with densitometry, vertebral fractures are assessed visually. While a number of grading systems have been proposed, the semiquantitative system of Genant is commonly used. This system grades the deformities from I to III, with grade I (mild) representing a 20% to 24% reduction in vertebral height, grade II (moderate) representing a 25% to 39% reduction in height, and grade III (severe) representing a 40% or greater reduction in height. The location of the deformity within the vertebrae may also be noted. For example, if only the mid height of the vertebrae is affected, the deformity is defined as an endplate deformity; if both the anterior and mid heights are deformed, it is a wedge deformity; and if the entire vertebrae is deformed, it is classed as a crush deformity. A vertebral deformity of at least 20% loss in height is typically considered a fracture. Accurate interpretation of both lateral spine radiographs and VFA imaging is dependent on radiologic training. Thus, device location and availability of appropriately trained personnel may influence diagnostic accuracy.

There is a lack of direct evidence from screening trials comparing densitometry with and without VFA that use of this technology improves health outcomes. Thus, screening for vertebral fractures using VFA with DXA, without densitometry is considered not medically necessary.

#### **CODING**

The following CPT code is covered, but not separately reimbursed for BlueCHiP for Medicare and is considered not medically necessary for Commercial Products:

**77086**

#### **RELATED POLICIES**

Bone Mineral Density Studies

#### **PUBLISHED**

Provider Update, November 2015

#### **REFERENCES**

1. National Government Services (NGS) Local Coverage Article (LCA) Bone Mass Measurement - Medical Policy Article (A51974)
2. National Osteoporosis Foundation. The Clinician's Guide to Prevention and Treatment of Osteoporosis 2013; <http://nof.org/files/nof/public/content/file/917/upload/481.pdf>. Accessed March, 2015.
3. Domiciano DS, Figueiredo CP, Lopes JB, et al. Vertebral fracture assessment by dual X-ray absorptiometry: a valid tool to detect vertebral fractures in community-dwelling older adults in a population-based survey. *Arthritis Care Res (Hoboken)*. May 2013;65(5):809-815. PMID 23212896
4. Diacinti D, Del Fiacco R, Pisani D, et al. Diagnostic Performance of Vertebral Fracture Assessment by the Lunar iDXA Scanner Compared to Conventional Radiography. *Calcif Tissue Int*. Nov 2012;91(5):335-342. PMID 22965625
5. Diacinti D, Guglielmi G, Pisani D, et al. Vertebral morphometry by dual-energy X-ray absorptiometry (DXA) for osteoporotic vertebral fractures assessment (VFA). *Radiol Med*. Jun 28 2012. PMID 22744340
6. Ferrar L, Jiang G, Clowes J, et al. Comparison of densitometric and radiographic vertebral fracture assessment using the algorithm-based qualitative (ABQ) method in postmenopausal women at low and high risk of fracture. *J Bone Miner Res*. 2007;23(1):103-111.
7. Binkley N, Krueger D, Gangnon R, et al. Lateral vertebral assessment: a valuable technique to detect clinically significant vertebral fractures. *Osteoporos Int*. 2005;16(12):1513-1518.
8. Kanterewicz E, Puigoriol E, Garcia-Barrionuevo J, et al. Prevalence of vertebral fractures and minor vertebral deformities evaluated by DXA-assisted vertebral fracture assessment (VFA) in a

population-based study of postmenopausal women: the FRODOS study. Osteoporos Int. Mar 6 2014. PMID 24599272

9. Mrgan M, Mohammed A, Gram J. Combined Vertebral Assessment and Bone Densitometry Increases the Prevalence and Severity of Osteoporosis in Patients Referred to DXA Scanning. J Clin Densitom. Oct-Dec 2013;16(4):549-553. PMID 23769657
10. Jager PL, Jonkman S, Koolhaas W, et al. Combined vertebral fracture assessment and bone mineral density measurement: a new standard in the diagnosis of osteoporosis in academic populations. Osteoporos Int. 2011;22(4):1059-1068.

[CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS](#)

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

