

EFFECTIVE DATE: 7/1/2007

POLICY LAST UPDATED: 1/28/2015

OVERVIEW

Vertebral axial decompression is a type of lumbar traction that has been investigated as a technique to reduce intradiscal pressure and relieve low back pain associated with herniated lumbar discs or degenerative lumbar disc disease.

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

BlueCHiP for Medicare and Commercial

Vertebral axial decompression is considered not medically necessary as there is insufficient peer-reviewed scientific literature that demonstrates that the procedure is effective.

MEDICAL CRITERIA

None

BACKGROUND

Vertebral axial decompression is a type of lumbar traction in which a pelvic harness is worn by the patient. The specially equipped table on which the patient lies is slowly extended, and a distraction force is applied via the pelvic harness until the desired tension is reached, followed by a gradual decrease of the tension. The cyclic nature of the treatment allows the patient to withstand stronger distraction forces compared to static lumbar traction techniques. An individual session typically includes 15 cycles of tension, and 10 to 15 daily treatments may be administered.

Examples of vertebral decompression therapy devices include, but may not be limited to:

- o Acua-Spina System utilizing Intervertebral Differential Dynamics (IDD Therapy)
- o Decompression Reduction Stabilization (DRS) System
- o DRX-3000
- o DRX9000
- o Lordex Traction Unit
- o SpineMED Decompression Table
- o V DRX 9000
- o VAX-D Table

Medicare determines that vertebral axial decompression does not meet the guidelines of reasonable and necessary as there is insufficient clinical literature to support the use of this device. Therefore, "Medicare does not cover items and services that are not reasonable and necessary for the diagnosis or treatment of an illness or injury or to improve the functioning of a malformed body member."

Evidence for the efficacy of vertebral axial decompression on health outcomes is limited. Because a placebo effect may be expected with any treatment that has pain relief as the principal outcome, randomized trials with validated outcome measures are required. The only sham-controlled randomized trial published to date did not show a benefit of vertebral axial decompression compared with the control group. Therefore,

Vertebral axial decompression is considered not medically necessary as there is insufficient peer-reviewed scientific literature that demonstrates that the procedure is effective.

COVERAGE

Benefits may vary. Please refer to the appropriate Member Certificate, Subscriber Agreement, or Benefit Booklet for applicable "Not medically necessary" service benefits/coverage.

CODING

BlueCHiP for Medicare and Commercial:

The following HCPCS code is not medically necessary for BlueCHiP for Medicare and Commercial Products: S9090 Vertebral axial decompression, per session

It is incorrect coding to file vertebral axial decompression using any other health service code such as chiropractic manipulation, nerve decompression surgery, or physical therapy manipulation.

RELATED POLICIES

None

PUBLISHED

Provider Update,	Apr 2015
Provider Update,	Nov 2014
Provider Update,	Sept 2013
Provider Update,	Jun 2012
Provider Update,	Jul 2011
Provider Update,	Aug 2010
Provider Update,	Jul 2009

REFERENCES

- 1.Schimmel JJ, de Kleuver M, Horsting PP et al. No effect of traction in patients with low back pain: a single centre, single blind, randomized controlled trial of Intervertebral Differential Dynamics Therapy. *Eur Spine J* 2009; 18(12):1843-50.
- 2.Sherry E, Kitchener P, Smart R. A prospective randomized controlled study of VAX-D and TENS for the treatment of chronic low back pain. *Neurol Res* 2001; 23(7):780-4.
- 3.Fritz JM, Lindsay W, Matheson JW et al. Is there a subgroup of patients with low back pain likely to benefit from mechanical traction? Results of a randomized clinical trial and subgrouping analysis. *Spine* 2007; 32(26):E793-800.
- 4.Harte AA, Baxter GD, Gracey JH. The effectiveness of motorised lumbar traction in the management of LBP with lumbo sacral nerve root involvement: a feasibility study. *BMC Musculoskelet Disord* 2007; 8:118.
- 5.Ramos G. Efficacy of vertebral axial decompression on chronic low back pain: study of dosage regimen. *Neurol Res* 2004; 26 (3):320-4.
- 6.Gose EE, Naguszewski WK, Naguszewski RK. Vertebral axial decompression therapy for pain associated with herniated or

degenerated discs or facet syndrome: an outcome study. *Neurol Res* 1998; 20(3):186-90.

7. Ramos G, Martin W. Effects of vertebral axial decompression 7. on intradiscal pressure. *J Neurosurg* 1994; 81(3):350-3.

Beattie PF, Nelson RM, Michener LA et al. Outcomes after a prone lumbar traction protocol for patients with activity-limiting low back pain: a prospective case series study. *Arch Phys Med Rehabil* 2008; 89(2):269-74.

8. Macario A, Richmond C, Auster M et al. Treatment of 94 outpatients with chronic discogenic low back pain with the DRX9000: a retrospective chart review. *Pain Pract* 2008; 8(1):11-7.

9. Centers for Medicare and Medicaid Services. National Coverage Decision for Vertebral Axial Decompression (VAX-D) (160.16). Available online at:

http://www.cms.hhs.gov/mcd/viewncd.asp?ncd_id=160.16&ncd_version=1&basket=ncd%3A160%2E16%3A1%3AVertebral+Axial+Decompression+%28VAX%2DD%29. Last accessed September, 2012.

10. Beattie PF, Nelson RM, Michener LA et al. Outcomes after a prone lumbar traction protocol for patients with activity-limiting low back pain: a prospective case series study. *Arch Phys Med Rehabil* 2008; 89(2):269-74.

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

